



# Utility of Balik's formula for the quantification of pleural effusion by ultrasound in the postoperative period of cardiac surgery

## Utilidad de la fórmula de Balik para la cuantificación del derrame pleural por ultrasonido en el posoperatorio de cirugía cardíaca

Camelia Cruz-Rodríguez,\* Edgar García-Cruz,\* Rodrigo Gopar-Nieto,\* Emmanuel Lazcano-Díaz,\*  
Diana Lizbeth Ordaz-Figueroa,\* José Luis Elizalde-Silva,\* Gustavo Rojas-Velasco,\* Daniel Manzur-Sandoval\*

\*Instituto Nacional de Cardiología Ignacio Chávez. Mexico City, Mexico.

**ABSTRACT. Introduction:** pleural effusion is caused by an imbalance between oncotic and hydrostatic pressure through the pulmonary capillaries or by increased permeability. Ultrasound at the patient's bedside allows for efficient diagnosis in various areas, due to its portability and low cost, enabling the quantification of pleural fluid, determination of its characteristics, and guidance for percutaneous drainage. Post cardiac surgery, large pleural effusions can affect the patient's recovery journey. **Material and methods:** we conducted a cross-sectional study of 26 nonconsecutive adult patients who underwent cardiac surgery in whom pleural effusion was detected by ultrasound in the postoperative period. The pleural effusion volume, quantified by Balik's formula, correlated with the amount of pleural fluid drained. In addition, the characteristics of the fluid were defined to determine any correlation with Light's criteria. **Results:** there was a strong positive correlation between the volume quantified by Balik's formula and the amount of pleural fluid drained. We also found that the characteristics of drained pleural effusion, as determined through ultrasound, had sufficient diagnostic accuracy to differentiate between transudate and exudate compared with Light's criteria. **Conclusions:** there is a strong positive correlation between the fluid volume quantified by ultrasound with Balik's formula and the volume drained in the postoperative period of cardiac surgery, in addition to high diagnostic accuracy in the identification of the fluid as transudate or exudate.

**Keywords:** pleural effusion, lung ultrasound, point-of-care ultrasound.

**RESUMEN. Introducción:** el derrame pleural es causado por un desequilibrio entre la presión oncótica e hidrostática a través de los capilares pulmonares o por un aumento de la permeabilidad. La ecografía a pie de cama del paciente permite un diagnóstico eficiente en diversas áreas, por su portabilidad y bajo costo, posibilitando la cuantificación del líquido pleural, determinación de sus características y orientación para el drenaje percutáneo. Después de una cirugía cardíaca, los derrames pleurales grandes pueden afectar el proceso de recuperación del paciente. **Material y métodos:** realizamos un estudio transversal de 26 pacientes adultos no consecutivos intervenidos de cirugía cardíaca en quienes se detectó derrame pleural mediante ecografía en el posoperatorio. El volumen del derrame pleural, cuantificado por la fórmula de Balik, se correlacionó con la cantidad de líquido pleural drenado. Además, se definieron las características del fluido para determinar cualquier correlación con los criterios de Light. **Resultados:** hubo una fuerte correlación positiva entre el volumen cuantificado por la fórmula de Balik y la cantidad de líquido pleural drenado. También encontramos que las características del derrame pleural drenado, determinadas por ultrasonido, tenían suficiente precisión diagnóstica para diferenciar entre trasudado y exudado en comparación con los criterios de Light. **Conclusiones:** existe una fuerte correlación positiva entre el volumen de líquido cuantificado por ultrasonido con la fórmula de Balik y el volumen drenado en el posoperatorio de cirugía cardíaca, además de una alta precisión diagnóstica en la identificación del líquido como trasudado o exudado.

**Palabras clave:** derrame pleural, ecografía pulmonar, ecografía en el punto de atención.

### Correspondence:

Daniel Manzur-Sandoval, MD

Instituto Nacional de Cardiología Ignacio Chávez. Mexico City

E-mail: drdanielmanzur@gmail.com

Received: VIII-21-2023; accepted: IX-27-2023.

**How to cite:** Cruz-Rodríguez C, García-Cruz E, Gopar-Nieto R, Lazcano-Díaz E, Ordaz-Figueroa DL, Elizalde-Silva JL et al. Utility of Balik's formula for the quantification of pleural effusion by ultrasound in the postoperative period of cardiac surgery. *Neumol Cir Torax.* 2023; 82 (2):72-78. <https://dx.doi.org/10.35366/115393>

## INTRODUCTION


Normally there is approximately 1-10 mL of fluid in the pleural space. This fluid is constantly produced and reabsorbed, and the amount of fluid is maintained by a balance between oncotic and hydrostatic pressure of the parietal and visceral pleura. Disruption in this balance causes the fluid to accumulate in the cavity.<sup>1</sup>

Pleural effusion is caused by an imbalance between oncotic and hydrostatic pressure through the visceral and parietal pleura, an increased permeability, or reduced absorption. It can occur as a result of lung parenchymal disease, infection, malignancy, and inflammatory processes. There are other factors that contribute to the accumulation of pleural fluid in the critical care setting, such as volume overload, renal or hepatic failure, myocardial depression, hypoalbuminemia, infections, and malnutrition.<sup>2</sup> In the postoperative period of cardiac surgery, pleural effusions, especially those that require a secondary drainage procedure during recovery, are associated with significantly worse outcomes including increased mortality, longer in-hospital stay, and higher complication rates. Of patients undergoing coronary artery bypass grafting or heart valve surgery, between 41% and 89% develop pleural effusions in the first seven days after surgery and 10% develop a pleural effusion occupying more than 25% of the hemithorax in the

subsequent month. Causes of pleural effusions after cardiac surgery include diaphragm dysfunction, internal mammary artery harvesting (only in internal mammary artery grafting), and other perioperative complications (e.g., sepsis, congestive heart failure, pulmonary embolism, and chylothorax).<sup>3</sup>

There are various imaging methods to evaluate the lung, pleura, and pleural cavity. While the presence of a pleural effusion is frequently suspected from a chest X-ray, it does not allow the detection of other fluid characteristics or its quantification. Characteristics such as localization, thickening, or fibrosis usually need characterization by axial computed tomography. Ultrasound at the patient's bedside allows for the diagnosis of pleural effusion in various hospital settings (due to its portability, low cost, absence of radiation, and short examination time), enabling the quantification of pleural fluid, determination of some characteristics of the effusion, and guidance for drainage by thoracocentesis or tube thoracostomy, thus improving the success rate of the procedures by up to 97%. Ultrasonographic diagnosis requires the identification of fluid in the pleural space with the typical anechoic (black) image between the diaphragm and the chest wall, which allows even small amounts of fluid to be identified.<sup>4,5</sup>

Regarding the etiology of the pleural effusion, ultrasound does not allow sufficient discrimination in the different etiologies and the compositions of the effusion. The



**Figure 1:**

- A)** Anechoic pleural effusion, suggesting transudate.
  - B)** Non-septated complex pleural effusion.
  - C)** Septated complex pleural effusion.
  - D)** Homogeneously echogenic pleural effusion.
- B-D)** Correspond to exudate. PE = pleural effusion. L = lung (compressive atelectasis).

sonomorphological characteristics may vary depending on its nature, cause, and chronicity. The appearance of the pleural fluid can be divided into four patterns: (1) anechoic (*Figure 1A*), (2) non-septated complex (defined by the presence of particles within the pleural fluid) (*Figure 1B*), (3) septated complex (defined by the presence of septa and fibrin in the pleural fluid) (*Figure 1C*), and (4) homogeneously echogenic («bright» pleural effusion) (*Figure 1D*), the last being the most common in hemothorax and empiema.<sup>6</sup>

### Importance


In chest X-ray, the accuracy to quantify the volume of the effusion is limited. Ultrasound allows for the detection of volume from 5 mL and volume quantification. Some formulas have been described for the quantitation of the effusion.<sup>7,8</sup> Balik's method found a significant positive correlation between the volume quantitation and the drained measurement. In addition to the quantitation of the effusion volume, Balik's method also allowed decision making regarding the realization of the therapeutic drainage, with success in guiding thoracentesis of 100% and no complications such as pneumothorax or bleeding reported.<sup>9</sup>

### Investigation goals

Our primary goal was to determine the correlation between the amount of pleural fluid quantified by ultrasonography using Balik's formula and the amount of fluid drained in the postoperative period of cardiac surgery. Our secondary objective was to determine the diagnostic accuracy of the ultrasonographic characteristics of the fluid to classify it as transudate or exudate.

## MATERIAL AND METHODS

This was a cross-sectional study of 26 nonconsecutive adult patients who were admitted to the critical care unit at the *Instituto Nacional de Cardiología Ignacio Chávez* in Mexico City, Mexico, from 1 May to 30 October 2021, following cardiac surgery, in whom ultrasonographic evaluation was conducted upon arrival in postoperative critical care. For the ultrasonographic evaluation, operator-obtained images were generated using a phased array sector probe at 2-3 MHz, from the patient's right or left side, with sonographic equipment including the following modes: M-mode, 2D mode, color Doppler, pulsed wave Doppler, continuous wave Doppler, and tissue Doppler. The use of sonographic equipment with advanced software technology was not necessary. With the patient in a mild torso elevation of 15°, the transducer was placed at the PLAPS point (intersection between the posterior



**Figure 2:** A) Position of the ultrasound probe at the patient's right or left side. B) Probe marker pointing toward the patient's head.

axillary line and the 7<sup>th</sup> or 8<sup>th</sup> intercostal space, slightly above the diaphragm),<sup>5</sup> with the probe marker pointing toward the patient's head. The diaphragm, liver, and spleen had to be clearly visualized (*Figure 2*). Then, the maximum distance in millimeters was measured between the parietal (from the lung inferior border) and visceral (to the inferior border of the diaphragm) pleura at maximum inspiration (*Figure 3*) using the formula: volume = 20 × separation (in millimeters).

Then, the pleural effusion was classified by ultrasonography as:

1. Transudate (anechoic) or
2. Exudate (non-septated complex, septated complex, and homogeneously echogenic).

Images were processed and analyzed after acquisition. One physician (a critical care physician with training in critical care ultrasonography) acquired the images, and the images were then processed and measured by three different physicians (clinical cardiologists, DMS, ECG, and ELD with training in echocardiography) using imaging software. We reduced bias by blinding the person who analyzed, processed, and measured the images.

The thoracic tubes were placed by the cardiothoracic surgery service after performing the ultrasound making,

without real-time guide, using Argyle TM Thoracic Catheters of 32 Fr when the effusion was echogenic or complex, and Blake drains (usually 24 Fr), when the effusion was anechoic.

The biochemical analysis of the fluid was performed to classify the pleural effusion as exudate according to Light's criteria:<sup>10</sup>

1. The ratio of pleural fluid protein to serum protein is greater than 0.5.
2. The ratio of pleural fluid lactate dehydrogenase (LDH) and serum LDH is greater than 0.6.
3. Pleural fluid LDH is greater than 0.6 or 2/3 times the normal upper limit for serum.

### Statistical analysis

We performed the Shapiro-Wilk test of normality for continuous variables and reported these as a median and interquartile range because all were nonparametric. Comparisons of continuous variables were made using the Kruskal-Wallis test. We report categorical variables as frequencies and percentages and used  $\chi^2$  or Fisher's exact probability tests, as appropriate, to compare expected values. We used Pearson's correlation coefficient to calculate correlation coefficients, with the following r value strength categories: 0.1-0.29 = weak, 0.3-0.49 = medium, and 0.50-1.0 = strong. We used 2 × 2 tables to calculate sensitivity, specificity, predictive values, and likelihood ratios. Statistical analyses were performed using STATA v. 14, and  $p < 0.05$  was considered statistically significant.


## RESULTS

### Demographic and surgical characteristics

Most patients were males (65%) with a mean age of 55 years (range 45-64 years). The most frequent comorbidities were heart failure (65.4%), hypertension (34.6%), and history of myocardial infarction (23.1%). The median of the left ventricular ejection fraction was 45%. The most frequently used drugs in the preoperative period were beta-blockers, anticoagulants, statins, antiplatelet drugs, diuretics, angiotensin-converting enzyme inhibitors, and angiotensin II receptor antagonists (Table 1). The most frequent surgery was aortic valve replacement in 34.6%, followed by coronary artery bypass graft surgery in 15.4% with a mean extracorporeal circulation time of 157 minutes and aortic clamping of 107 minutes (Table 2).

### Characteristics of pleural effusion

The median volume quantified by ultrasound was 600 mL, and the median volume drained was 550 mL. There was a strong positive correlation ( $r = 0.67$ ,  $p < 0.0001$ ). Of the total number of patients, pleural effusion was classified as transudate by ultrasonography in 15 patients (57.7%) and by Light's criteria in 16 patients (61.5%). The most frequent drainage method was tube thoracostomy (84.6%), being more frequent on the left side (53.8%) (Table 3). When the diagnostic test evaluation for the presence of transudate vs. exudate by ultrasound was performed and compared with Light's criteria, the sensitivity and specificity were 81.25% and 80%, respectively. There was a positive predictive value



**Figure 3:**

The measurement is made from the lower border of the lung (parietal pleura) to the abdominal structure (visceral pleura). **A)** Anechoic pleural effusion. **B)** Non-septated complex pleural effusion.



**Table 1:** Baseline characteristics.

Variable	n (%)
Age (years), median [IQR]	55 [45-64]
LVEF (%), median [IQR]	45 [32-54]
Gender	
Men	17 (65.4)
Women	9 (34.6)
Body mass index	
Underweight	2 (7.7)
Normal range	10 (38.5)
Overweight	7 (26.9)
Obese class I	6 (23.1)
Obese class III	1 (3.8)
Heart failure	17 (65.4)
Hypertension	9 (34.6)
Prior myocardial infarction	6 (23.1)
Dyslipidemia	4 (15.4)
Diabetes	4 (15.4)
Stroke	3 (11.5)
Chronic obstructive pulmonary disease	1 (3.8)
Chronic renal disease	1 (3.8)
Peripheral vascular disease	1 (3.8)
Prior medication	
Beta-blockers	11 (42.3)
Anticoagulants	10 (38.5)
Statins	10 (38.5)
Diuretics	9 (34.6)
Antiplatelet drugs	8 (30.8)
Angiotensin-converting enzyme inhibitors	7 (26.9)
Angiotensin II receptor antagonists	6 (23.1)
Aldosterone antagonists	4 (15.4)
Oral hypoglycemic drugs	3 (11.5)
Calcium channel blockers	3 (11.5)
Cardiac glycoside	2 (7.7)
Amiodarone	2 (7.7)
Levothyroxine	1 (3.8)
Allopurinol	1 (3.8)

LVEF = left ventricular ejection fraction. IQR = interquartile range.

of 86.66%, a negative predictive value of 72.73%, and a positive likelihood ratio of 4 (Table 4).

## DISCUSSION

Pleural effusion occurs frequently in patients admitted to intensive care units and varies according to the technique used, from 8% with physical examination to 60% using imaging techniques. Early drainage of clinically significant pleural effusion is associated with improved oxygenation and diagnostic accuracy without increased complications.<sup>11</sup> In postoperative cardiac surgery patients, pleural effusion can cause atelectasis, increased risk of pneumonia and empyema, arrhythmias (such as atrial fibrillation), prolonged

in-hospital stay, need for hemodialysis, and higher mortality.<sup>3</sup> Modifiable associated factors in the management of drains that may contribute to the accumulation of pleural effusion include early removal of chest drains, higher outputs, and removal during or close to mechanical ventilation.<sup>12</sup> Risk factors that have been identified include being female and previous conditions such as heart failure, atrial fibrillation, peripheral vascular disease, and prior use of anticoagulant or antiarrhythmic agents.<sup>13</sup> In addition, dedicated follow-up and treatment of postoperative effusions enhance recovery by 15% measured by improvement in the distance walked one month after cardiac surgery.<sup>14</sup>

**Table 2:** Surgical characteristics.

Variable	n (%)
Surgery	
Aortic valve replacement	9 (34.6)
Coronary artery bypass graft	4 (15.4)
Coronary artery bypass graft + aortic valve replacement	3 (11.5)
Mitral valve replacement	2 (7.7)
Mitral valve replacement + aortic valve replacement	2 (7.7)
Coronary artery bypass graft + mitroaortic valve replacement	2 (7.7)
Coronary artery bypass graft + mitral valve replacement	1 (3.8)
Extracorporeal circulation time (min), median [IQR]	157 [110-214]
Aortic clamping (min), median [IQR]	107 [89-147]

IQR = interquartile range.

**Table 3:** Characteristics of pleural effusion.

Variable	n (%)
Ultrasonographic classification	
Transudate	15 (57.7)
Exudate	11 (42.3)
Light's criteria classification	
Transudate	16 (61.5)
Exudate	10 (38.5)
Drainage method	
Thoracentesis	4 (15.4)
Thoracic tube	22 (84.6)
Localization	
Left lung	14 (53.8)
Right lung	12 (46.2)
Ultrasonically quantified volume (mL), median [IQR]	600 [400-800]
Drained volume (mL), median [IQR]	550 [440-800]

IQR = interquartile range.

**Table 4:** Diagnostic test evaluation for the diagnosis of transudate vs exudate by ultrasound.

	Light's criteria		
	Transudate n (%)	Exudate n (%)	Total n (%)
Ultrasound			
Transudate	13 (81.2)	2 (20.0)	15 (57.7)
Exudate	3 (18.7)	8 (80.0)	11 (42.3)
Total	16 (100.0)	10 (100.0)	26 (100.0)
Diagnostic test evaluation			
Sensitivity		81.25%	
Specificity		80.00%	
PPV		86.66%	
NPV		72.73%	
LR +		4.06	
LR -		0.23	
Accuracy		80.77	

PPV = positive predictive value. NPV = negative predictive value.  
LR = likelihood ratio.

In our study, most of the population was male, with a median age of 55 years. The majority of the patients had a history of heart failure, followed by hypertension and myocardial infarction, which suggests that these patients could have a greater probability of presenting pleural effusion in the postoperative period due to increased hydrostatic pressure.

Among the drugs administered to our patients, a higher proportion of patients were taking anticoagulants and antiplatelet drugs, which could also contribute to the development of pleural effusion. Regarding surgical variables, the median extracorporeal circulation time was 157 minutes, and the aortic clamping time was 107 minutes, which suggest that prolonged extracorporeal circulation pump and aortic clamping times (> 100 minutes) may be a risk factor for developing pleural effusion due to increased bleeding.

The median volume measured by ultrasonography was 600 mL, and one evacuate was 550 mL. When the correlation was determined between the measurement performed by ultrasound using Balik's formula and the volume of liquid drained, there was a strong positive correlation. In the only previous study evaluating the estimation of volume by ultrasound in the postoperative period of cardiac surgery, the maximal distance between mid-height of the diaphragm and visceral pleura, using the formula  $\text{volume} = 16 \times \text{separation}$  (in millimeters), was used, and the authors found a significant positive correlation between the volume estimated and the volume drained. The ultrasonographic characteristics of the effusion were not described.<sup>15</sup>

Although there are studies that have previously compared formulas for the quantitation of pleural effusion,

the equation used by us and proposed by Balik et al.<sup>9</sup> is validated for mechanically ventilated patients in a supine position and with a mild torso elevation of 15°. This is the position in which we usually find our patients in the intensive care unit in the immediate period after cardiac surgery, so we think it is more adequate in this population.

When the characteristics of the pleural effusion were evaluated by ultrasonography, we found that 57% corresponded to transudates and the rest to exudates, according to the echogenicity of the effusion. When the fluid was evacuated, the type of effusion was corroborated by Light's criteria. A total of 61% corresponded to transudate.

Yang et al. evaluated a cohort of 320 patients with pleural effusion, finding high sensitivity, but poor specificity of anechoic effusions for transudative effusions.<sup>16</sup> In a cohort of 126 patients with transudative pleural effusions, Chen et al. found that an anechoic pattern was present in 45% (57/127), while a complex non-septated pattern was seen in 55% (70/127); transudative fluid was never complex septated or homogeneously echogenic.<sup>17</sup> A recent evaluation of 300 pleural effusions in 285 patients demonstrated that the detection of septations or homogenous complexity was 94% specific and carried a 96% positive predictive value for exudative fluid. Furthermore, anechoic fluid did not reliably predict the presence of transudative fluid.<sup>18</sup> Regarding the analysis of the characteristics of the fluid by ultrasound, in our series, high sensitivity, positive predictive value, and positive likelihood ratio (81.25%, 86.66%, and 4.06, respectively) stood out. It is important to mention that this is the first time that these findings have been evaluated in postoperative cardiac surgery patients who had an increased risk of transudate due to increased hydrostatic pressure.

Regarding the exudates, when a septated complex and homogeneously echogenic pleural effusion was identified by ultrasound, it always corresponded to exudate according to Light's criteria.

Complications associated with chest tube placement are reported in the medical literature in up to 40% of cases.<sup>19</sup> In 22 of the patients of our study (84%), the effusion was evacuated by placing a thoracic tube without complications. The complications that occurred were: one patient with tube malposition, one patient with minor bleeding from the insertion site, one patient with insertion site infection, and one patient with re-expansion pulmonary edema.

In up to 84% of the patients, the effusion was evacuated by placing a thoracic tube without complications, even though complications associated with the placement of a chest tube are reported in 20-30% of cases. The most frequent location of pleural effusion was on the left side in up to 53% of cases, as previously reported,<sup>19</sup> which suggests that the anatomical relationship of the heart with the left pleura could be a factor in the patients who presented with a greater effusion on the left side.

### Study limitations

This study was conducted at a single medical center and should be replicated at other centers to assess the protocol's reproducibility. In addition, study outcomes should be interpreted with caution due to the small sample size and because it was performed in a specific subpopulation (postoperative cardiac surgery patients). Thus, the findings may not generalize to all critical care patients. Finally, given that the study was conducted in a cardiovascular critical care unit, where point-of-care ultrasonography evaluation is routine, adequate training is necessary before applying these techniques.

### CONCLUSION

We found a strong positive correlation between the volume of pleural effusion quantified with Balik's formula by ultrasound and the volume drained in postoperative cardiac surgery patients. We also found an adequate diagnostic accuracy of the ultrasound in the identification of the type of effusion compared with Light's criteria.

### REFERENCES

- Mazhar S. An audit to evaluate the quantification of pleural effusions using thoracic ultrasonography in ICU. *Ann Pak Inst Med Sci.* 2019;15(4):191-194. doi: 10.48036/apims.v15i4.242.
- Brogi E, Gargani L, Bignami E, Barbariol F, Marra A, Forfori F, et al. Thoracic ultrasound for pleural effusion in the intensive care unit: a narrative review from diagnosis to treatment. *Critical Care [Internet].* 2017;21(1):325. Available in: <https://ccforum.biomedcentral.com/articles/10.1186/s13054-017-1897-5>
- Schiefenhovel F, Poncette AS, Boyle EM, von Heymann C, Menk M, Vorderwülbecke G, et al. Pleural effusions are associated with adverse outcomes after cardiac surgery: a propensity-matched analysis. *J Cardiothorac Surg.* 2022;17(1):298.
- Ahmed ES, Bakr A, Eid HA, Shaarawy AT, Elsayed WT. Role of ultrasonography in the diagnosis of pleural effusion. *Egypt J Bronchol.* 2017;11(2):120-127. doi: 10.4103/1687-8426.203797.
- Lichtenstein DA. Lung ultrasound in the critically ill. *Ann Intensive Care.* 2014;4(1):1. doi: 10.1186/2110-5820-4-1.
- Ibitoye BO, Idowu BM, Ogunrombi AB, Afolabi BI. Ultrasonographic quantification of pleural effusion: comparison of four formulae. *Ultrasonography.* 2018;37(3):254-260. doi: 10.14366/usg.17050.
- Cotton DW, Lenz R, Kerr B, Ma I. Point of care ultrasound for the general internist: pleural effusions. *Can Journ Gen Int Med [Internet].* 2018;13(2). Available in: <https://cgjim.ca/index.php/csim/article/view/231>
- Dietrich CF, Mathis G, Cui XW, Ignee A, Hocke M, Hirche TO. Ultrasound of the pleurae and lungs. *Ultrason Med Biol.* 2015;41(2):351-365. doi: 10.1016/j.ultrasmedbio.2014.10.002.
- Balik M, Plasil P, Waldauf P, Pazout J, Fric M, Otahal M, et al. Ultrasound estimation of volume of pleural fluid in mechanically ventilated patients. *Intensive Care Med.* 2006;32(2):318-321. doi: 10.1007/s00134-005-0024-2.
- Light RW, Macgregor MI, Luchsinger PC, Ball WC Jr. Pleural effusions: the diagnostic separation of transudates and exudates. *Ann Intern Med.* 1972;77(4):507-513. doi: 10.7326/0003-4819-77-4-507.
- Fysh ETH, Smallbone P, Mattock N, McCloskey C, Litton E, Wibrow B, et al. Clinically significant pleural effusion in intensive care: a prospective multicenter cohort study. *Crit Care Explor.* 2020;2(1):e0070. doi: 10.1097/CCE.000000000000070.
- Brookes JDL, Williams M, Mathew M, Yan T, Bannon P. Pleural effusion post coronary artery bypass surgery: associations and complications. *J Thorac Dis.* 2021;13(2):1083-1089. doi: 10.21037/jtd-20-2082.
- Labidi M, Baillet R, Dionne B, Lacasse Y, Maltais F, Boulet LP. Pleural effusions following cardiac surgery: prevalence, risk factors, and clinical features. *Chest.* 2009;136(6):1604-1611. doi: 10.1378/chest.09-0689.
- Hansen LS, Hjortdal VE, Jakobsen CJ, Heiberg J, Maagaard M, Sloth E. Early, dedicated follow-up and treatment of pleural effusions enhance the recovery rate after open cardiac surgery: results from a randomized, clinical trial. *Eur J Cardiothorac Surg.* 2017;51(1):58-66. doi: 10.1093/ejcts/ezw233.
- Usta E, Mustafi M, Ziemer G. Ultrasound estimation of volume of postoperative pleural effusion in cardiac surgery patients. *Interact Cardiovasc Thorac Surg.* 2010;10(2):204-207. doi: 10.1510/icvts.2009.222273.
- Yang PC, Luh KT, Chang DB, Wu HD, Yu CJ, Kuo SH. Value of sonography in determining the nature of pleural effusion: analysis of 320 cases. *AJR Am J Roentgenol.* 1992;159(1):29-33. doi: 10.2214/ajr.159.1.1609716.
- Chen HJ, Tu CY, Ling SJ, Chen W, Chiu KL, Hsia TC, et al. Sonographic appearances in transudative pleural effusions: not always an anechoic pattern. *Ultrason Med Biol.* 2008;34(3):362-369. doi: 10.1016/j.ultrasmedbio.2007.09.009.
- Shkolnik B, Judson MA, Austin A, Hu K, D'Souza MS, Zumbunn A, et al. Diagnostic accuracy of thoracic ultrasonography to differentiate transudative from exudative pleural effusion. *Chest.* 2020;158(2):692-697. doi: 10.1016/j.chest.2020.02.051.
- Tanner TG, Colvin MO. Pulmonary complications of cardiac surgery. *Lung.* 2020;198(6):889-896. doi: 10.1007/s00408-020-00405-7.

**Ethics approval and consent to participate:** the local institutional research and ethics committees waived approval for this study. Informed consent was obtained from all individual participants included in the study.  
**Consent for publication:** written informed consent for patient information and images to be published was provided by the patient or a legally authorized representative.

**Conflict of interests:** the authors do not have any conflict of interests to declare.