



Gender-adjustment of Moyers dentition analysis for the Nayarit (Mexico) population

Ajuste por género del análisis de dentición de Moyers para la población de Nayarit

Jaime Fabián Gutiérrez Rojo,* Damaris Delgado Sandoval,[§] Alhelí Mendoza Minjarez,^{||} Alma Rosa Rojas García*

ABSTRACT

Moyers percentile tables are one of the diagnostic elements used for early age malocclusions. These tables predict the size of non erupted canines and premolars. **Material and methods:** The sample was composed of 180 male patients study models and 319 female patients study models used for orthodontic pre-treatment. Models met inclusion criteria. So as to match samples, 180 female study models were randomly selected. Descriptive statistics were calculated, percentiles 75 and 50 for each gender, t Student test was used to compare results of tables with adjustment of percentiles 75 and 50 for each gender with Moyers tables results and values of the study models. **Results:** Statistically significant differences were found when comparing Moyers' tables results with those of the Nayarit population. Percentile 75 of Nayarit tables over-estimated values of canines and premolars in both genders. No statistical differences were found with 50 percentile. **Conclusions:** It is recommended to use the 50 percentile of the conducted adjustment in order to predict size of canines and premolars.

Key words: Moyers, mixed dentition analysis.

Palabras clave: Moyers, análisis de dentición mixta.

RESUMEN

Uno de los elementos diagnósticos de las maloclusiones en edades tempranas son las tablas de percentiles de Moyers, éstas predicen el tamaño de caninos y premolares que aún no erupcionan. **Material y métodos:** La muestra fue de 180 modelos de estudio pretratamiento de ortodoncia de sexo masculino que cumplían los criterios de inclusión y 319 de sexo femenino, por lo que se seleccionaron de forma aleatoria 180 modelos de estudio pretratamiento de sexo femenino para igualar las muestras. Se calculó la estadística descriptiva, los percentiles 75 y 50 para cada sexo y se utilizó la prueba de t de Student para comparar los resultados de las tablas con los ajustes de los percentiles 75 y 50 de cada sexo con el resultado de las tablas de Moyers y con los valores de los modelos de estudio. **Resultado:** Se encontraron diferencias estadísticas entre los resultados de las tablas de Moyers y las realizadas para la población de Nayarit. El percentil 75 de las tablas de Nayarit sobrestima el valor de caninos y premolares en ambos sexos, con el percentil 50 no se encontraron diferencias estadísticas. **Conclusiones:** Se recomienda utilizar el percentil 50 del ajuste realizado para predecir el tamaño de caninos y premolares.

INTRODUCTION

Malocclusions are an important factor in the field of social acceptance and interaction as well as in limitations of masticatory functions.¹ Most malocclusions commence during the period of mixed dentition, therefore, mixed dentition analyses are necessary for diagnosis during the dental replacement stage.²

Mixed dentition analyses are used to predict mesio-distal size of canines and premolars. These analyses will determine existing discrepancies between available and required space in the dental arch.³ There are different systems to conduct analysis of existing space during mixed dentition, but most possess as a characteristic the need to have erupted first permanent molars as well as mandibular incisors in order to predict the size of premolars and canines.⁴

Dentition analyses can be classified as follows: those conducted based on formulae, which can be either statistical or of linear regression and based on lower incisors, and those which used X-rays, as well as a combination of both methods.^{5,6}

Moyers' dentition analysis is one of the most commonly used methods, it was developed for Caucasian population.⁷ It has the characteristic of using

* Faculty member of the Dental Academic Unit and Orthodontics Specialty of the Nayarit Autonomous University.

§ Orthodontics Specialist. Private Practice.

|| Student at the Dental School of the Autonomous University of Nayarit.

Received: September 2013. Accepted: November 2013.

This article can be read in its full version in the following page:
<http://www.medigraphic.com/facultadodontologiaunam>

the width of all four lower incisors in order to establish the mesio-distal size of teeth being predicted; this is achieved in two percentile tables divided by gender, one for the upper jaw and one for the lower jaw.⁸

Moyers recommends the use of the 50 percentile, considering it a more precise calculation, and the use of percentile 75 in cases when the clinical operator might be inexperienced, since percentile 75 calculates greater medio-distal values for non erupted canines and premolars. The advantages of Moyers' analysis are the following: it entails minimal system errors, it is easy to rapidly read, it does not require X-rays, it does not require special equipment, it can be conducted either on dental models or on the patient, there are tables for the upper and lower jaw.⁹

Several studies have described mesio-distal size variation in teeth according to patient's ethnicity and gender.¹⁰ Therefore, it might prove necessary to assess dental analyses and if said analyses were not effective, undertake necessary adjustments in order to be able to use them.

The purpose of the present research paper was to assess Moyers' analysis for the Nayarit population and, if it were to be non-effective, undertake necessary adjustments.

MATERIAL AND METHODS

The present research was of a descriptive, non-experimental and cross-section nature, 1,345 study models were examined. Models came from the database of the Orthodontics Specialty, Nayarit Autonomous University. Models spanned the 2002-2012 time frame. The sample was convenience-oriented, that is to say, models that met with inclusion criteria were included; 180 models of pre-orthodontic male patients were used, as well as 319 models of female patients. Out of this female patient group, 180 models were randomly selected so as to match the 180 male models.

Inclusion criteria were as follows: pre-orthodontic treatment models, models were to be manufactured

by only two operators of the Diagnosis Unit, models should exhibit erupted permanent dentition except for third molars. The following pre-treatment models were excluded: those which exhibited fractured teeth, cavities, previous orthodontic treatment, restorations which affected mesio-distal width as well as crowding. All of the aforementioned factors would have prevented mesio-distal measurement of the tooth.

Variables were anterior and posterior segments. The anterior segment was obtained from the sum of the mesiodistal values of lower central and lateral incisors. The upper and lower posterior segment was built with the mean of mesio-distal measurement of canines, first premolar and second premolar of both jaws.

Data gathering was conducted in a recording sheet with the patient's initials, gender, age, impression age, and slots for all teeth to be considered for the present research. All data was processed with Excel 2007 Microsoft Office program; statistics were undertaken with the SPSS, version 18 program.

Materials used were the study models, a Mitutoyo brand digital gauger, thin brushes, data collection sheets, pencil, eraser and a computer. Descriptive statistics were calculated, 75 and 50 percentiles for each gender, t Student test was used to compare table results with 75 and 50 percentile adjustments for each gender with the results of the Moyers tables. Comparison was equally established between adjustment results for tables of the population under study and premolar and canine values of the orthodontic study models.

RESULTS

Descriptive statistics can be observed in *tables I and II*. 75 and 50 percentiles were calculated with the values of canines and premolars of pre-orthodontics study models for both genders (*Tables III and IV*).

Moyers analysis was applied, as well as Nayarit population adjustment. Results were compared and revealed; there were statistically significant

www.medigraphic.org.mx

Table I. Descriptive statistic of female gender population.

	Real premolars and canines		Percentile adjustment 75		Percentile adjustment 50	
	Upper	Lower	Upper	Lower	Upper	Lower
Mean	22.10	21.35	22.6	21.8	22.07	21.25
Standard deviation	1.26	1.23	0.71	0.8	0.67	0.79
Maximum	26	24.7	24.5	23.7	23.8	23.3
Minimum	17.8	17.6	20.9	19.9	20.2	19.1

differences ($p \leq .001$). It could therefore be concluded that this percentile cannot be used for mixed dentition diagnoses.

After this, tables proposed for the present research with the 50 percentile were compared with real values of canines and premolars; no statistically significant differences were found. In females, «p» in upper jaw was 0.615, for males it was 0.118, in the lower jaw «p» values were 0.121 in females and 0.486 in males.

In the tables, minimum value for incisor sum was 20.5 mm in males and 20 mm in females, maximum values were 27 mm for both genders. Moyers' tables results and adjustment were different, due to this, tables were not the same (in adjustment tables lack of ascending continuity could be observed in the values of canines and premolars could be observed, contrary to Moyers tables where value of canines and premolars always increase along with incisor values).

Percentile 75 over-estimated in both genders the value of canines and premolars, for this reason it should not be used in orthodontic diagnosis. On the other hand, percentile 50 was the one to use for dentition analyses, since no statistically significant differences were found.

DISCUSSION

Moyers' percentile tables have been researched all over the world, it was found they were not effective due to genetic differences among populations. For example, in India, size of premolars and canines were under-estimated in the regions of Pondicherry, Mukstar, Ludhiana and Jangam.¹¹⁻¹³ Contrary to this, in other locations such as Sarajevo (Croatia) or Graz (Austria) this mixed dentition analysis over-estimated the values of non erupted canines and premolars.¹⁴ In Nayarit, Moyers' analysis under-estimated value of population's canines and premolars.¹⁵

In India, Ganapati *et al* found that Moyers's analysis was not suitable for use, due to the fact that in the 75 percentile it over-estimated the mesio-distal size of canines and premolars, in the 50 percentile, it under-estimated or over-estimated the size according to gender or whether it was conducted in the upper or lower jaw.² In the present research we concurred with Ganapati's findings when using the 75 percentile on the studied Nayarit population, since statistical differences were found.

Table II. Descriptive statistic of male population.

	Real premolars and canines		Percentile adjustment 75		Percentile adjustment 50	
	Upper	Lower	Upper	Lower	Upper	Lower
Mean	22.05	21.6	22.40	22.12	21.94	21.49
Standard deviation	1.24	1.22	0.64	0.57	0.66	0.57
Maximum	26	25	23.3	23	22.7	22.2
Minimum	19.3	19.1	20.8	20.7	20.2	20

Table III. Percentiles table for males.

Incisors	20.5	21	21.5	22	22.5	23	23.5	24	24.5	25	25.5	26	26.5	27
75 maxillary	22.6	22.7	21.8	22.9	22.7	22.8	23.5	23.6	23.8	24.1	23.7	24.5	24.2	24.8
75 mandibular	21.6	21.8	21.3	22.3	21.9	21.8	22.9	23	23.4	23.2	23	24.2	23.7	24.3
50 maxillary	21.2	21.4	21.3	22	22.1	22.1	22	23.1	23.3	23.7	23.1	23.3	23.3	24.6
50 mandibular	20.2	20.8	20.6	21.5	21.6	21.2	21.7	22.3	22.6	22.8	22.8	23.4	23.1	23.6

Table IV. Percentiles table for females.

Incisors	20	20.5	21	21.5	22	22.5	23	23.5	24	24.5	25	25.5	26	26.5	27
75 maxillary	20.9	21.5	21.7	22.4	21.8	22.2	22.5	22.8	23	22.9	23.2	24.3	23.8	24.3	24.3
75 mandibular	19.9	20.2	20.4	20.9	21.4	21.3	21.8	22.2	22.3	22.5	22.4	23.9	23.4	23.2	23.7
50 maxillary	20.2	21.5	21.1	21.9	21.4	21.5	21.9	22.3	22.5	22.7	22.6	23	23.2	23.7	23.8
50 mandibular	19.1	19.6	19.9	20.5	20.5	20.8	21	21.2	21.8	21.8	21.7	21.9	22.6	23.2	23.3

In the South of Austria, in the city of Graz, it was found that Moyers's tables underestimated the size of the teeth which erupted, after the fact that when sum of incisors reached 23 in maxillary and mandibular value tables, values were over-estimated.¹⁴ In the present study, values for canines and premolars were over-estimated when using the 75 percentile, from the lowest values of incisor sums (20.5 mm in males and 20 mm in females) up to 27 mm values.

Several authors have reported gender dimorphism in the dental size of canines and premolars, this has led to undertaking different dentition analyses for males and females. The present study revealed differences among male and female values, similar to those of Philip in India,¹¹ Diagne in Senegal¹⁶ and Ganapati in India.²

The problem of Moyers's tables' precision can be decreased when they are adjusted to the population to which they are applied.¹⁷ Percentile tables of canines and premolars sizes have been made in India,⁵ Senegal,¹⁶ USA, Austria¹⁴ and Hong Kong.¹⁶ In the percentile tables made up for the present research it can be observed that incisor values as well as percentile values are higher than those published by Moyers.

CONCLUSIONS

Assessment of the Moyers' analysis adjustment to Tepic, Nayarit population revealed that percentile 75 was not effective to calculate the size of canines and premolars in both genders. No statistical differences were found with the results of calculating mesiodistal size of canines and premolars with percentile 50 adjustment, as well as values of canines and premolars in the Nayarit population.

The adjustment of Moyers' analysis in the 50 percentile executed in the present research project, was only effective for the population where it was conducted. Therefore, it is recommended to assess mixed dentition analyses in populations where it was executed, since in different populations, there are differences in mesiodistal sizes of teeth.

REFERENCES

- Scapini A, Feldens C, Machado T, Floriani P. Malocclusion impacts adolescents oral health-related quality life. *Angle Orthod.* 2013; 83 (3): 512-518.
- Ganapati S, Naik V. Evaluation of Moyers mixed dentition analysis children. *Indian J Dent Res.* 2009; 20 (1): 26-30.
- Jarjees H, Awni K, Hasan Y. Validation of Tanaka and Johnston method on Iraqi population. *Al-Rafidain Dent J.* 2009; 9 (1): 17-23.
- Marín G, Pérez M, Califa M, Abdullah E, Al-Shawi A, Abdullah E et al. Validación de la ecuación de Tanaka Johnston en una población de escolares yemenita. *Revista Cubana de Estomatología.* 2009; 46 (4): 23-31.
- Boitor C, Stoica F, Nasser H. Prediction of the mesiodistal size of unerupted canines and premolars for a group of Romanian children: a comparative study. *J Appl Oral Sci.* 2013; 21 (3): 225-230.
- Rakosi T, Jonas I. *Atlas de ortopedia maxilar: diagnóstico.* Barcelona: Editorial Masson; 1992: pp. 219-221.
- Cabello M, Mendoza N, Parés V. Valoración de la exactitud de predicción del tamaño dental mesiodistal de las tablas de probabilidad de Moyers y las ecuaciones de Tanaka/Johnston en una población mexicana. *Revista ADM.* 2004; 61 (5): 176-182.
- Nourallah A, Gesch D, Khordaji M, Splieth C. New regression equations for predicting the size of unerupted canines and premolars in a contemporary population. *Angle Orthod.* 2002; 72 (3): 216-221.
- Moyers R. Manual de ortodoncia. 4a ed. Buenos Aires: Panamericana; 1998: p. 240.
- Oyeyemi V, Da Costa O. Mixed dentition analysis among a Nigerian population. *IJCD.* 2013; 4 (1): 31-36.
- Philip N, Prabhakar M, Arora D, Chopra S. Applicability of the Moyers mixed dentition probability tables and new prediction aids for a contemporary population in India. *Am J Orthod Dentofacial Orthop.* 2010; 138: 339-345.
- Sholapurmath S, Benni D, Mandroli P. Applicability of two mixed dentition analysis in children of Jangam community of Belgaum city. *World J Dent.* 2012; 3 (4): 324-329.
- Legovic M, Novosel A, Legovic V. Regression equations for determining mesiodistal crown diameters of canines and premolars. *Angle Orthod.* 2003; 73 (3): 314-318.
- Droschl H, Golles J, Sager K. Sobre la utilización de las tablas de probabilidades de Moyers. *Revista Española de Ortodoncia.* 1977; 7 (4): 185-201.
- Gutiérrez-Rojo J, Delgado-Sandoval D, Rojas-García A, Rivas-Gutiérrez R, Mendoza-Minjarez A. *Efectividad del análisis de Moyers en la población que asiste a la Clínica de Ortodoncia de la Universidad Autónoma de Nayarit.* Memorias de las Jornadas de Investigación de la Universidad Autónoma de Nayarit: 2011.
- Diagne F, Diop-Ba K, Ibrahima P, Mbow K. Mixed dentition analysis in a Senegalese population: elaboration of prediction tables. *Am J Orthod Dentofacial Orthop.* 2003; 124: 178-183.
- Buwembo W, Luboga S. Moyers method of mixed dentition analysis: a meta-analysis. *Afr Health Sci.* 2004; 4 (1): 63-66.

Mailing address:

Jaime Fabián Gutiérrez Rojo

E-mail: jaimefg79@hotmail.com