

Original article

doi: 10.35366/101859

Incidence and risk factors for acute kidney injury: a prospective, randomized, single-blind clinical trial comparing total knee arthroplasty with and without tourniquet

Incidencia y factores de riesgo para lesión renal aguda: Un ensayo clínico ciego, prospectivo y aleatorizado comparando la artroplastía total de rodilla con y sin torniquete

Colín-Vázquez A,* Galindo-Ávalos J,† Salas-Alemán M,* Bernal-Fortich LD,§
López-Valencia J,¶ Vidal-Cervantes F,¶ Pereyra-Arzate R,|| Rivera-Villa AH**

UMAE «Dr. Victorio de la Fuente Narváez», IMSS.

ABSTRACT. Purpose: Identify the incidence and risk factors for acute kidney injury (AKI) following total knee arthroplasty (TKA) with and without tourniquet. **Material and methods:** 100 patients were randomized into two groups. Postoperative AKI was defined as the postoperative creatinine level ≥ 0.3 mg/dl compared with baseline. Potential variables associated with AKI were analyzed by multivariate logistic regression model to identify the AKI risk factors in TKA patients with and without tourniquet. **Results:** AKI rate was 22%, tourniquet use (OR = 2.66, $p = 0.014$), blood loss > 500 cm³ (OR = 3.99, $p = 0.001$), postoperative Hb < 10 g/dl (OR = 2.68, $p = 0.008$), blood transfusions (OR = 2.86, $p = 0.012$) and diabetes (OR = 2.80, $p = 0.006$) were associated with increased risk of postoperative AKI. **Conclusions:** The use of tourniquet should be indicated with caution and should not be

RESUMEN. Propósito: Identificar la incidencia y factores de riesgo para lesión renal aguda (LRA) después de la artroplastia total de rodilla (ATR) con y sin uso de torniquete. **Material y métodos:** Se dividieron 100 pacientes en dos grupos. Se definió la LRA como una elevación postoperatoria de la creatinina ≥ 0.3 mg/dl comparada con el nivel basal preoperatorio. Las potenciales variables asociadas con la LRA fueron analizadas con un modelo de regresión logística multivariada para identificar los factores de riesgo de LRA en pacientes sometidos a ATR con y sin torniquete. **Resultados:** La incidencia de LRA fue de 22%. El uso de torniquete (OR = 2.66, $p = 0.014$), pérdida sanguínea > 500 cm³ (OR = 3.99, $p = 0.001$), Hb postoperatoria < 10 g/dl (OR = 2.68, $p = 0.008$), transfusión sanguínea (OR = 2.86, $p = 0.012$) y la diabetes (OR = 2.80, $p = 0.006$) fueron asociados a un mayor riesgo postoperatorio

Level of evidence: II

* Traumatólogo y Ortopedista, adscrito al Servicio de Reemplazos Articulares, UMAE «Dr. Victorio de la Fuente Narváez», IMSS.

† Traumatólogo y Ortopedista, adscrito a la Clínica *San Maré Health-Care Group*. ORCID: 0000-0003-2396-0963.

§ Traumatólogo y Ortopedista, adscrito al Nuevo Hospital de Bocagrande, Colombia.

¶ Traumatólogo y Ortopedista, UMAE «Dr. Victorio de la Fuente Narváez» IMSS.

|| Residente primer año de Traumatología y Ortopedia, UMAE «Dr. Victorio de la Fuente Narváez», IMSS-UNAM.

** Traumatólogo y Ortopedista, Jefe de Servicio, Reemplazos Articulares, UMAE «Dr. Victorio de la Fuente Narváez», IMSS.

Correspondence:

Dr. Joel Galindo-Ávalos

Bld. Francisco Medina Ascencio Núm. 2735-9,

Zona Hotelera Norte,

Puerto Vallarta, Jalisco, C.P. 48333.

Tel: 322 252 1711

E-mail: orthofitness2017@gmail.com

Received: 22-04-2020. Accepted: 18-01-2021.

How to cite: Colín-Vázquez A, Galindo-Ávalos J, Salas-Alemán M, Bernal-Fortich LD, López-Valencia J, Vidal-Cervantes F, et al. Incidence and risk factors for acute kidney injury: a prospective, randomized, single-blind clinical trial comparing total knee arthroplasty with and without tourniquet. *Acta Ortop Mex.* 2021; 35(2): 158-162. <https://dx.doi.org/10.35366/101859>



used routinely in patients with other risk factors for the development of acute kidney dysfunction, other measures to achieve trans-surgical hemostasis should be implemented in our environment to reduce the incidence of acute kidney dysfunction related to the use of the tourniquet.

Keywords: Total knee arthroplasty, acute kidney injury, tourniquet, risk factors.

de LRA. **Conclusiones:** El uso de torniquete debe estar indicado con precaución y no debe utilizarse de forma rutinaria en pacientes con otros factores de riesgo para el desarrollo de disfunción renal aguda, otras medidas para lograr la hemostasia transquirúrgica deben implementarse en nuestro entorno para reducir la incidencia de disfunción renal aguda relacionada con el uso del torniquete.

Palabras clave: Artroplastía total de rodilla, lesión renal aguda, torniquete, factores de riesgo.

Introduction

Total knee arthroplasty (TKA), is a procedure commonly performed by orthopedic surgeons across the globe, and it is expected that the demand for primary and revision TKAs will rise exponentially in our country. Only at our hospital, Unidad Médica de Alta Especialidad «Dr. Victorio de la Fuente Narváez», a total of 1,720 primary TKAs and 254 revision TKAs were performed in 2017, which is a 19% increase compared to 2016, when a total of 1,443 primary TKAs and 218 revision TKAs were performed. Acute kidney injury (AKI) is a serious complication

following surgery. It is now recognized that even mild AKI can develop into chronic kidney disease (CKD) in a previously normal kidney and can also cause rapid disease progression in patients with chronic kidney failure, and it is also associated with an increase in hospital length of stay, costs and mortality.¹

The incidence of AKI following elective and emergency orthopedic procedures has been reported to be up to 9%.² The incidence of AKI following elective joint replacement goes from < 2 to 14.6%.^{3,4,5}

The Acute Kidney Injury Network (AKIN) defines AKI as an increase of serum creatinine concentration of >




Figure 1: Flowchart of the population studied.

Table 1: Comorbidities by group.

Group 1 (n = 50)	n (%)
• Arterial high blood pressure	41 (82)
• Diabetes mellitus	16 (32)
• Rheumatoid arthritis	11 (22)
• Chronic kidney disease	3 (6)
• Hypothyroidism	3 (6)
• No comorbidities	4 (8)
Group 2 (n = 50)	n (%)
• Arterial high blood pressure	43 (86)
• Diabetes mellitus	14 (28)
• Rheumatoid arthritis	3 (6)
• Chronic kidney disease	1 (2)
• Hypothyroidism	1 (2)
• Parkinson's disease	1 (2)
• Dyslipidemia	1 (2)
• Mixed cardiomyopathy	1 (2)
• No comorbidities	6 (12)

26.5 μmol ($> 0.3 \text{ mg/dl}$) or 1.5 times its baseline value at 48 hours postoperative, as well as a decrease in urinary output $< 0.5 \text{ ml/kg/h}$.⁶

Tourniquet application is almost indispensable in the orthopedic practice. A tourniquet is often used in total knee arthroplasty resulting in improved visualization of structures, reduced intraoperative bleeding and better cementation.^{7,8,9} However, a series of complications can be related to tourniquet use, and these complications may be related to and excessive tourniquet time or pressure. The local and systemic physiological effects and the anesthetic implications are reviewed. Localized complications result from either tissue compression beneath the cuff or tissue ischemia distal to the tourniquet. Systemic effects are related to the inflation or deflation of the tourniquet.^{10,11,12,13}

The purpose of this study was to evaluate the incidence of AKI following primary TKA, and to identify whether tourniquet application is a risk factor for developing AKI.

Material and methods

ClinicalTrials.gov Identifier: NCT03795805

We conducted the trial in Mexico City, with subjects recruited from the Department of Joint Replacements at Unidad Médica de Alta Especialidad «Dr. Victorio de la Fuente Narváez». The protocol was approved by the Instituto Mexicano del Seguro Social ethics committee. The study was performed in accordance with the protocol, and all subjects provided written informed consent.

Subjects older than 18 years old, who presented at the consult with a diagnosis of knee osteoarthritis, had failed a trial of nonoperative treatment and where schedule to undergo TKA were included. Major exclusion criteria were patients who did not sign informed consent or were schedule to undergo unicompartamental knee arthroplasty or osteotomies.

Patients were randomly assigned to undergo into two groups by computer generated random numbers. Group 1 (G1) underwent TKA with tourniquet application and group 2 (G2) underwent TKA without tourniquet application. All patients in G1 were operated on by the same surgeon (MS-A) as did the patients in G2 (FV-C) (*Figure 1*).

Serum creatinine and hemoglobin levels were obtained as part of preoperative evaluation and at 48 hours following surgery. Blood loss was quantified as both the bleeding during surgery as well as blood collected in the surgical drain. Range of motion was evaluated 48 hours after surgery using a goniometer. Skin-to-skin surgical time was obtained from the anesthetic record sheet. All measures were recorded in a data collection sheet.

A single injection of antibiotic prophylaxis was given. In G1 a pneumatic tourniquet was used after exsanguination using an Esmarch bandage. In G2 a combination of 50 cm^3 normal saline, 0.5 mg epinephrine and 246.25 ropivacaine was injected intra articular to the knee joint 5 minutes before surgery started. Subcutaneous thromboprophylaxis was started in the evening before surgery. The anterior approach was used with a midline incision and a standard parapatellar arthrotomy. Soft tissues were released in a stepwise manner to achieve ligamentous balance in extension. Thus, flexion and extension gaps were approximately equal. Enough laxity was achieved to enable full extension and flexion and anterior translation, but not too loose to cause abnormal anteroposterior motion, impingement, or bearing spin out. Tensioning devices were not routinely used. The proximal tibia was resected with the anterior cruciate ligament sacrificed. The posterior cruciate ligament was preserved with a bone block. Both the femoral and tibial components were cemented. Primary patellar resurfacing was not performed. A surgical drain was placed prior to skin closure.

Postoperatively, physiotherapy (walking exercises aided with a walker and continuous passive motion exercises) was started on day one. Manual lymphatic drainage was performed by the physiotherapist on day two. Surgical drain was removed on day two.

Results

A total of 100 patients were included in the study (50 in each group). Patients were predominantly women (54%), with a mean age of 67.5 years ($41\text{-}86 \pm 8.18$). In G1, patients were predominantly women (58%), with a mean age of 66.82 years (range 41-84, SD 8.93). In G2, half the patients were men and the other half women, with a mean age of 68.48 years (range 47-86, SD 7.34). Comorbidities by group are shown in *Table 1*.

Mean surgical time was 83.7 ± 14.2 , mean surgical blood loss was 528 cm^3 ($528 \pm 119.4 \text{ cm}^3$), differences between groups are shown in *Table 2*. Differences between Hemoglobin levels in each group and differences between range of motion and pain in each group are shown in *Table*

2. All patients started walking with the help of a walker at 48 hours postoperative. Fourteen percent of patients required a blood transfusion, 11 in G1 and 3 in G2 ($p = 0.021$).

Regarding serum creatinine level difference, we found mean preoperative levels of 0.79 ± 0.21 mg/dl and mean postoperative levels of 0.87 ± 0.33 mg/dl ($p = 0.001$), differences between groups are shown in [Table 2](#).

Acute kidney injury rate was found to be 22% when strict definition was applied (elevation of > 0.3 mg/dl compared with baseline). Risk factors identified are shown in [Table 3](#).

Discussion

The main finding of the present study was to describe a 22% incidence of acute kidney dysfunction in patients who underwent total knee arthroplasty, as well as to prove that the use of tourniquet in total knee arthroplasty significantly increases serum creatinine levels 48 hours after surgery, and also increases the risk of suffering acute kidney dysfunction by 2.6 times. Likewise, the use of the tourniquet is related to an increase in total postoperative bleeding, lower postoperative flexion capacity and an increase in the postoperative pain scores. It was observed that, in addition to the use of the tourniquet, the presence of diabetes mellitus, blood loss greater than 500 cm^3 , postoperative Hb less than 10 g/dl , or the need for transfusion, also increases the risk of presenting subsequent acute kidney dysfunction to a total knee arthroplasty.

Hassan and colleagues,¹⁴ in their retrospective study, described an incidence of acute renal dysfunction of 9.7%, compared with 22% of our study, which allowed them

to identify risk factors such as age over 65 years and the presence of arterial high blood pressure, which were not identified in our study, and on the contrary, they did not identified the presence of Diabetes mellitus as a risk factor, which was identified in our study.

Similarly, in a study conducted by Wu et al.,¹⁵ a relatively low incidence of acute renal dysfunction (3.3%) is described in comparison with our study, and, like us, they did not identified arterial high blood pressure, BMI greater than 30, surgical time over 90 minutes, and the patient's gender as risk factors for the development of acute kidney dysfunction, however, unlike our study, they did not identified blood loss $> 500 \text{ cm}^3$, the need for transfusion or the presence of diabetes mellitus to be risk factors for acute kidney dysfunction.

In the study by Kimmel et al.,³ an incidence of acute renal dysfunction secondary to prosthetic joint replacement of 15% was reported, as well as, among the multiple risk factors listed, the use of a pneumatic tourniquet, but an Odds Ratio is not provided. Likewise, Diabetes mellitus is listed as a risk factor for acute kidney dysfunction together with blood transfusions, similar to our study.

Conclusion

The use of tourniquet should be indicated with caution and should not be used routinely in patients with other risk factors for the development of acute kidney dysfunction, other measures to achieve trans-surgical hemostasis should be implemented in our environment to reduce the incidence of acute kidney dysfunction related to the use of the tourniquet. Likewise, factors such as diabetes mellitus,

Table 2: Differences in outcomes between groups.

	G1	G2	p [‡]
Time (min)*	85.2 (15.3)	82.2 (12.9)	0.291
Blood loss (cm ³)*	581 (100.9)	475 (113.5)	< 0.001
Flexion (°)*	59.6 (17.3)	82.2 (12.9)	< 0.001
Extension(°)*	0 (0)	0.2 (1.4)	0.320
Pain (VAS)*	7.02 (1.2)	5.08 (0.6)	< 0.001
		Mean (g/dl)	p [§]
G1			
• Preoperative hemoglobin		14.1	< 0.001
• Postoperative hemoglobin		10.4	
G2			
• Preoperative hemoglobin		14.7	< 0.001
• Postoperative hemoglobin		10.5	
G1			
• Preoperative serum creatinine		0.74	< 0.001
• Postoperative serum creatinine		0.91	
G2			
• Preoperative serum creatinine		0.84	0.472
• Postoperative serum creatinine		0.82	

* Values are expressed as mean (SD). ‡ Student's t-test for independent samples was used. § Student's t-test for paired samples was used. Min = minutes, ° = Degrees, VAS = visual analogue scale.

Table 3: Risk factors for acute kidney injury following total knee arthroplasty.

	AKI (n = 22)	No AKI (n = 78)	OR (IC 95%)	p
Tourniquet application	16	34	2.667 (1.137-6.254)	0.014
Age older than 65 years	17	48	1.831 (0.738-4.541)	0.132
Male	13	33	1.696 (0.798-3.602)	0.125
Surgical time > 90 minutes	5	14	1.254 (0.529-2.972)	0.409
Blood loss > 500 cm ³	17	29	3.991 (1.597-9.978)	0.001
Postoperative hemoglobin < 10 g/dl	13	22	2.683 (1.275-5.645)	0.008
Blood transfusion	7	7	2.867 (1.428-5.756)	0.012
High blood pressure	18	66	0.857 (0.334-2.199)	0.448
Diabetes mellitus	12	18	2.800 (1.360-5.764)	0.006
Rheumatoid arthritis	5	9	1.807 (0.794-4.109)	0.161

AKI = acute kidney injury, TKA = total knee arthroplasty.

systemic arterial hypertension, transurgical bleeding and BMI should be studied due to conflicts in the international literature.

References

- Borthwick E, Ferguson A. Perioperative acute kidney injury: risk factors, recognition, management, and outcomes. *BMJ*. 2010; 341: c3365. doi: 10.1136/bmj.c3365.
- Kateros K, Doulgerakis C, Galanakos SP, Sakellariou VI, Papadakis SA, Macheras GA. Analysis of kidney dysfunction in orthopaedic patients. *BMC Nephrol*. 2012; 13: 101. doi: 10.1186/1471-2369-13-101.
- Kimmel LA, Wilson S, Janardan JD, Liew SM, Walker RG. Incidence of acute kidney injury following total joint arthroplasty: a retrospective review by RIFLE criteria. *Clin Kidney J*. 2014; 7(6): 546-51. doi: 10.1093/ckj/sfu108.
- Pulido L, Parvizi J, Macgibeny M, Sharkey PF, Purtill JJ, Rothman RH, et al. In hospital complications after total joint arthroplasty. *J Arthroplasty*. 2008; 23(6 Suppl 1): 139-45. doi: 10.1016/j.arth.2008.05.011.
- Weingarten TN, Gurrieri C, Jarett PD, Brown DR, Berntson NJ, Calaro RD Jr, et al. Acute kidney injury following total joint arthroplasty: retrospective analysis. *Can J Anaesth*. 2012; 59(12): 1111-8. doi: 10.1007/s12630-012-9797-2.
- Mehta RL, Kellum JA, Shah SV, Molitoris BA, Ronco C, Warnock DG, et al. Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care*. 2007; 11: R31.
- Tai TW, Chang CW, Lai KA, Lin CJ, Yang CY. Effects of tourniquet use on blood loss and soft-tissue damage in total knee arthroplasty: a randomized controlled trial. *J Bone Joint Surg Am*. 2012; 94(24): 2209-15. doi: 10.2106/JBJS.K.00813.
- Alcelik I, Pollock RD, Sukeik M, Bettany-Saltikov J, Armstrong PM, et al. A comparison of outcomes with and without a tourniquet in total knee arthroplasty: a systematic review and meta-analysis of randomized controlled trials. *J Arthroplasty*. 2012; 27: 331-40. doi: 10.1016/j.arth.2011.04.046.
- Tetro AM, Rudan JF. The effects of a pneumatic tourniquet on blood loss in total knee arthroplasty. *Can J Surg*. 2001; 44(1): 33-8.
- Smith TO, Hing CB. Is a tourniquet beneficial in total knee replacement surgery? A meta-analysis and systematic review. *Knee*. 2010; 17: 141-7. doi: 10.1016/j.knee.2009.06.007.
- Kam PC, Kavanagh R, Yoong FF. The arterial tourniquet: pathophysiological consequences and anaesthetic implications. *Anaesthesia*. 2001; 56(6): 534-45.
- Rama KRBS, Apsingi S, Poovali S, Jetti A. Timing of tourniquet release in knee arthroplasty. Meta-analysis of randomized, controlled trials. *J Bone Joint Surg Am*. 2007; 89(4): 699-705. doi: 10.2106/JBJS.F.00497.
- Tai TW, Lin CJ, Jou IM, Chang CW, Lai KA, Yang CY. Tourniquet use in total knee arthroplasty: a meta-analysis. *Knee Surg Sports Traumatol Arthrosc*. 2011; 19(7): 1121-1130. doi: 10.1007/s00167-010-1342-7.
- Hassan BK, Sahlstrom A, Dessau RB. Risk factors for renal dysfunction after total knee joint replacement. *Acta Orthop Belg*. 2015; 81: 647-53.
- Wu KT, Chen CY, Chen B, Wang JW, Lin PC, Yen SH. The incidence and risk factors of acute kidney disease after total knee arthroplasty with early postoperative volume supplement. *Biomed Res Int*. 2018; 2018: 8718545. doi: 10.1155/2018/8718545.