

## ANNALS OF HEPATOLOGY

Volume **4**

Number **3**

July-September **2005**

*Artículo:*

**Budd-Chiari syndrome**

Copyright © 2005:  
Mexican Association of Hepatology

**Otras secciones de  
este sitio:**

-  [Índice de este número](#)
-  [Más revistas](#)
-  [Búsqueda](#)


***Others sections in  
this web site:***

-  [Contents of this number](#)
-  [More journals](#)
-  [Search](#)

Images in Hepatology

## Budd-Chiari syndrome

Nahum Méndez-Sánchez;<sup>1</sup> Norberto C. Chávez-Tapia;<sup>1</sup> Eric López-Méndez;<sup>1</sup> Misael Uribe<sup>1</sup>



**Figure 1.** Hepatic vein selective catheterization and injection through a transjugular approach; showing a small network of collateral veins opacified representing the typical "spider web" appearance of hepatic vein occlusion.

A 29 years old man with Budd-Chiari syndrome. Budd-Chiari syndrome is defined as hepatic venous outflow obstruction at any level from the small hepatic veins to the junction of the inferior vena cava and the right atrium, regardless of the cause of obstruction.<sup>1</sup> It is caused mainly by thrombosis of the hepatic veins. The etiology of Budd-Chiari syndrome caused by hepatic vein thrombosis can be diverse, the hematological diseases associated are polycythemia vera and paroxysmal nocturnal hemoglobinuria. Other acquired risk factors for Budd-Chiari syndrome include antiphospholipid syndrome, abdominal trauma, the use of oral contraceptives, pregnancy, and so on.<sup>2</sup> Budd-Chiari syndrome can be classified as primary (due to intrinsic intraluminal thrombosis or webs) or secondary (due to intraluminal invasion by a parasite or malignant tumour or due to extraluminal compression by an abscess, cyst or solid tumour).<sup>3</sup> According to clinical presentation can be classified as fulminant, acute, sub-acute, or chronic. Therapy for patients with the Budd-Chiari syndrome includes medical management and the relief of hepatic venous outflow tract obstruction in order to prevent necrosis, with liver transplantation in selected patients, especially those with fulminant hepatic failure.<sup>4</sup>

### References

1. Janssen HL, Garcia-Pagan JC, Elias E, Mentha G, Hadengue A, Valla DC. Budd-Chiari syndrome: a review by an expert panel. *J Hepatol* 2003; 38: 364-71.
2. Araki Y, Sakaguchi C, Ishizuka I, et al. Budd-Chiari syndrome: A case with a combination of hepatic vein and superior vena cava occlusion. *World J Gastroenterol* 2005; 11: 3797-9.
3. Bogin V, Marcos A, Shaw-Stiffel T. Budd-Chiari syndrome: in evolution. *Eur J Gastroenterol Hepatol* 2005; 17: 33-5.
4. Menon KV, Shah V, Kamath PS. The Budd-Chiari syndrome. *N Engl J Med* 2004; 350: 578-85.

<sup>1</sup> Liver Unit. Medica sur Clinic & Foundation. Mexico City, Mexico.

Address for correspondence:

Nahum Méndez-Sánchez, M.D., PhD. Departments of Biomedical Research, Gastroenterology & Liver Unit. Medica Sur Clinic & Foundation, Puente de Piedra 150, Col. Toriello Guerra, Tlalpan 14050, Mexico City, Mexico. Phone: (+525) 55606-6222 ext. 4215; Fax: (+525) 55666-4031; E-mail: nmendez@medicasur.org.mx

Manuscript received and accepted: July 18, 2005.