



Concise Review

The future of stem cells in liver diseases

Maurizio Muraca,¹ Ginevra Galbiati,¹ Maria Teresa Vilei,² Aline Sueli Coelho Fabricio,² Maddalena Caruso¹

Abstract

Preliminary experience with clinical hepatocyte transplantation during the past decade has provided proof of concept that cell therapy can be effective for the treatment of some liver diseases. Recent progress in cell biology resulting in the isolation and characterization of hepatic stem cells and progenitor cells further increased the expectation for a new approach to the treatment of genetic and chronic liver disease. Several potential sources have been identified of hepatic stem/progenitor cells exhibiting both differentiation towards the hepatic lineage *in vitro* and hepatic parenchymal repopulation with liver-specific metabolic activity in liver-injured animal models. However, a few of these results proved to be poorly reproducible in different laboratories, and it was recognized that some initial optimistic conclusions were drawn from incorrect interpretation of experimental data or from insufficient knowledge of the mechanisms involved in tissue regeneration. Moreover, only modest results have emerged so far from ongoing clinical experience involving the use of putative stem cells in liver disease. There is much need for a joined effort to concentrate the resources on a specific cell population, in order to better characterize its function, to assess its safety and

to develop better focused clinical trials. In conclusion, while the biological features of stem cells still justify the hope for future clinical applications, hepatic stem cell therapy has still a long way to go from bench to bedside.

Key words: Hematopoietic stem cells, umbilical cord blood, metabolic diseases, liver regeneration, liver cell therapy.

Cell therapy can be defined as «the use of living cells to restore, maintain or enhance the function of tissues and organs».¹ The use of isolated, viable cells has emerged as an experimental therapeutic tool in the past decade, due to progress in cell biology and particularly in techniques for the isolation and culture of cells derived from several organs and tissues. However, experimental cell therapy has a longer tradition in Hepatology, since it has been known for more than 30 years that isolated hepatocytes infused into the portal vein engraft into the liver cords and express normal cell function. Such a therapeutic strategy was put forward as an alternative to orthotopic liver transplantation (OLT), which requires major surgery and is limited by the availability of donors. Indeed, it was shown that significant clinical results can be obtained with the transplantation of isolated hepatocytes corresponding to as little as 1-5% of the total hepatocyte mass.²⁻⁶

The procedure seems relatively safe, provided portal pressure and/or portal flow are monitored during cell infusion in order to prevent vascular thrombosis.⁷ Hepatocyte transplantation has recently been used as an alternative to OLT in patients with liver-based congenital metabolic disorders, such as Crigler-Najjar disease,⁸ α -1-antitrypsin deficiency,⁹ glycogen storage disease type Ia,¹⁰ ornithine transcarbamoylase deficiency^{11,12} and the deficiency of factor VII.¹³ The role of hepatocyte transplantation in the treatment of acute and chronic liver disease is less clear,^{9,14} due to difficulty in organizing large-scale clinical trials. Indeed, the main factor limiting the practice of hepatocyte transplantation is again the availability of liver grafts for cell isolation. Moreover, the metabolic effects of cell transplantation seem to be fading with time, a problem which can partially be solved by repeated hepatocyte infusions¹⁵ but which probably indicates the progressive loss of the terminally-differentiated exogenous cells. In theory, both problems could be solved by replacing the hepatocyte with

¹ Laboratory Medicine, Department of Laboratories, Ospedale Pediatrico Bambino Gesù, Rome, Italy.

² Clinica Medica 1, University of Padova, Italy.

Address for correspondence:

Prof. Maurizio Muraca, MD, Ph D
IRCCS Ospedale Pediatrico Bambino Gesù, Piazza Sant'Onofrio, 4,
00165 Roma, Italy.
Tel. +39 06 6859 2210.
E-mail: muraca@opbg.net

Abbreviations:

SHPCs: Small Hepatocyte-like Progenitor Cells; HSCs: Hematopoietic Stem Cells; BM: Bone Marrow; UCB: Umbilical Cord Blood.

Grants and Financial supports

The financial support of Telethon – Italy (grant no. GGP02051), of the Italian Association for the Study of the Liver (AISF) and of the Regione Veneto (Centro Regionale per la Terapia Cellulare delle Malattie Metaboliche and grant no. 03/03/01) is gratefully acknowledged.

stem or precursor cells, provided they can be isolated from a more affordable source.

Indeed, at present, there is growing interest in the therapeutic use of stem cells.^{16,17} A stem cell has the ability to divide for indefinite periods of time, to self-renew and to give rise to many different cell types. Embryonic stem cells originate from the inner cell mass of the mammalian blastocyst and are totipotent.¹⁸ Adult stem cells are more specialized, being committed to give rise to cells with a particular function within their own specific tissue or organ.¹⁹ Precursor/progenitor cells are defined as cells rapidly dividing and already partially determined towards a specific differentiation pathways.²⁰ However, experimental evidence suggests that some adult stem cells are able to develop into different types of specialized cells (a process also known as transdifferentiation), depending on the microenvironment where they are homed, including the liver.^{19,21-30}

This review will address a series of major issues on hepatic stem cells, including their origin, their role in liver regeneration and fibrosis, and their possible use in the treatment of liver disease.

What is the origin and what are the possible sources of hepatic stem cells?

In mammals, the liver has the unique ability among solid organs to regenerate following parenchymal injury. During fetal development, hepatoblasts give rise both to hepatocytes and to cholangiocytes. The hepatocyte is traditionally considered as the cell responsible for liver regeneration, being able to re-enter the cell cycle, proliferate and differentiate both into hepatocytic and biliary lineage in response to loss of liver mass. However, an intra-hepatic progenitor cell compartment, resident in the canals of Hering and consisting of precursors known as «oval cells» in rodents (or «hepatic progenitor cells» in human), is activated when the replicative capacity of the main epithelial cell compartment is inhibited or exhausted.³¹⁻³⁶ These small round cells express phenotypic markers of both fetal hepatocytes and biliary cells^{35,37-39} and are able to differentiate into hepatocytes, bile ductal cells and intestinal epithelium.⁴⁰

SHPCs also appear to be involved in the process of hepatic regeneration. SHPCs are highly proliferative and can generate mature differentiated hepatocytes *in vitro*.^{41,42} SHPCs express markers such as albumin, alpha-fetoprotein, transferrin, form bile canaliculi and store glycogen.⁴³ SHPCs also appear to be involved in the process of hepatic regeneration after partial hepatectomy (PH) in rats pre-treated with retrorsine, a pyrrolyzidine alkaloid which severely impairs the proliferating capacity of mature hepatocytes.^{41,44,45} In this model, hepatic repopulation takes place mainly by proliferation of SHPCs, exhibiting phenotypic traits in common with fully differentiated hepatocytes, fetal hepatoblast, and oval cells.⁴⁶ Some

reports suggest that SHPCs don't originate from oval cells,⁴² but that they derive from a pre-existing population of retrorsine-resistant hepatocytes.⁴⁷ Indeed, during liver regeneration, the SHPCs lack hepatic cytochrome P450 protein⁴⁸ that is responsible for metabolizing retrorsine to pyrrolic derivatives⁴⁹ and are thus resistant to the toxic effect of the drug. Avril *et al.*⁴⁷ labeled mature hepatocytes using a recombinant retroviral vector harboring the β -galactosidase "LacZ" gene in the retrorsine/PH rat model. During parenchymal regeneration, a similar (4%) proportion of β -galactosidase-positive SHPCs and of mature hepatocytes was observed, suggesting that mature hepatocytes could be the actual progenitors of SHPCs.⁴⁷ However, more recently, using 3-dimensional image analysis in the retrorsine/PH model of liver regeneration, Vig *et al.* observed that oval cells surround and penetrate SHPCs nodules, suggesting that SHPCs nodules can originate from oval cells.⁴³

The origin of hepatic stem or precursor cells is still a matter of debate. Interestingly, oval cells express markers of Hematopoietic Stem Cells (HSCs), such as Thy-1, CD34, CD45, Sca-1, c-Kit and flt-3.^{22,50-54} In particular, Thy-1 is a highly conserved protein. It has been found in the brain and in the hematopoietic system of rat, mouse and humans.⁵³ It is also expressed on stem cells of fetal liver and in bone marrow (BM)-derived cells.⁵⁵ In addition, the normal adult liver contains hematopoietic cells that are phenotypically similar to cells present in the BM.⁵⁶ These observations originated the hypothesis that liver stem cells may arise from a population resident in the BM.²² Petersen *et al.*²² followed the fate of syngeneic BM cells transplanted into lethally irradiated rats whose livers were subsequently injured by 2-acetylaminofluorene and CCl₄, a regimen known to induce oval cell proliferation. Using the *sry* gene as a marker for the Y chromosome, male donor cells were visualized in female recipients. In a separate experiment, dipeptidyl peptidase IV (DPP IV)-positive hepatocytes were identified in the liver of DPP IV-deficient rats transplanted with BM from DPP IV-positive animals.

Lagasse *et al.*²⁹ transplanted fumarylacetoacetate hydrolase (FAH)-deficient mouse, an animal model of Tyrosinemia type I, with BM cells from a non-affected wild-type animal transgenic for the β -galactosidase "LacZ" gene. The liver of the recipient animals was progressively repopulated with hepatocytes harboring both the β -galactosidase and the fumarylacetoacetate hydrolase enzyme. Thus, intravenous injection of adult BM cells in the FAH^{-/-} mouse rescued the mouse and restored the biochemical function of its liver. It was later shown that the correction of the metabolic disorder was not due to transdifferentiation of HSCs but rather to a fusion process, probably involving macrophages derived from the exogenous hematopoietic cell lineage and the recipient hepatocytes.^{57,58} This phenomenon can be demonstrated by cytogenetic analysis in sex-discordant transplantation (*Figure 1*).

However, other Authors have later demonstrated in different models that HSCs can convert in hepatocytes without fusion both *in vitro* and *in vivo*. By co-culturing HSCs with injured liver tissue, Jang *et al.*,⁵⁹ observed production of cells expressing the immunocytochemical and genetic features of hepatocytes, and maintaining the original chromosomal pattern. After two days in culture, about 3% of the cells converted to hepatocytes. A similar finding was observed after transplantation of BM-derived cells from male animals into female animals with liver injury: after 2 days, 8% of hepatocytes incorporated the Y chromosome, while maintaining the original male chromosomal pattern, suggesting differentiation rather than fusion.⁵⁹ Several investigators have claimed *in vitro* differentiation of BM-derived cells into hepatocytes, although the results were often difficult to be reproduced in different laboratories. The group of Catherine Verfaillie isolated from the BM a population of Multipotent Adult Progenitor Cells (MAPCs) with the ability to differentiate in culture into endothelium, neuroectoderm and endoderm. These cells seem to be indeed pluripotent, a feature of embryonic stem cells. Hepatic differentiation of these cells was obtained after a few days in culture with appropriate growth factors.⁶⁰ Lee *et al.*,⁶¹ showed that Mesenchymal Stem Cells (MSCs) isolated from human BM are able to differentiate into functional hepatocyte-like cells. Setting up a novel *in vitro* hepatic differentiation protocol, based on the sequential use of HGF and Oncostatin M, and following cells behavior under hepatogenic conditions at different

times, these Authors showed that the MSCs acquired hepatocyte morphology and expressed genes and functions characteristic of liver parenchymal cells. A modified differentiation medium, containing FGF-4, Oncostatin M, HGF and EGF, was used by Shi *et al.*, to induce BM-derived mononuclear cells of C57BL/6 mice to differentiate into hepatocyte-like cells.⁶² However, in contrast to the above results indicating that oval cells may be derived from the BM, recent work suggests that such cells originate in the liver itself.^{43,57,63}

In summary, experimental evidence suggests that liver parenchymal cells can originate from a specific precursor cell compartment in the liver, from pluripotent stem cells, from transdifferentiation of HSCs or from cell fusion. The occurrence of true transdifferentiation, or reprogramming to the hepatic lineage of an already committed hematopoietic stem cell, is still a matter of debate.³⁰

Additional, potential sources of hepatic stem cells have been identified in the cord blood, in the amniotic fluid and in the placenta. Umbilical Cord Blood (UCB) contains hematopoietic and mesenchymal stem/precursor cells Lee *et al.*^{64,65} The transplantation of UCB has been used for more than 10 years for the treatment of hematologic and genetic diseases.⁶⁶⁻⁷¹ UCB cells are easily accessible, proliferate *in vitro* and have longer telomeres compared to adult cells, indicating higher proliferation capacity.⁷²⁻⁷⁴ Several Authors have investigated the hepatic potential of UCB-derived cells *in vitro* and *in vivo*. Kakinuma *et al.*,⁷⁵ showed that human UCB cells cultured in the presence of a particular combination of growth and differentiation factors (i.e. FGF-1, FGF-2, LIF, SCF and HGF) were able to produce albumin and other hepatocytes specific markers *in vitro*. When inoculated into liver-injured SCID mice, a few functionally differentiated human UCB-derived hepatocytes were found 55 weeks post transplantation. By means of a «two-step» differentiation medium, Lee *et al.*,⁶⁵ induced human UCB-derived (CD3⁺, CD14⁺, CD19⁺, CD38⁺, DC66b⁺, glycophorin A⁺) MSCs to differentiate into hepatocyte-like cells. Sharma *et al.*,⁷⁶ reported that human UCB-derived mononuclear cells generate hepatocyte-like cells after transplantation into NOD-SCID mice with severe hepatocellular damage produced by CCl₄. Noteworthy, all such cells showed some specific human hepatic markers but not a mature hepatocyte phenotype. Moreover, all the donor-derived hepatic cells expressed human albumin and human hepatocyte-specific antigen Hep Par 1 but also expressed the murine cytokeratine CK18, suggesting the occurrence of fusion between human and mouse cells. Newsome *et al.*,⁷⁷ infused an unsorted UCB mononuclear cell preparation into sub-lethally irradiated NOD-SCID mice. These cells were able to engraft into mice livers and differentiate into hepatocytic lineage with no evidence of fusion with mouse hepatocytes.⁷⁷ Piscaglia *et al.*,⁷⁸ transplanted into immunocompetent rats a CD34⁺/CD45⁺ and CD133⁺/CD45⁺ population from human




Figure 1. Distinguishing differentiation from fusion by cytogenetic analysis based on the identification of sex chromosomes.

A. Differentiation of BM-derived cells into liver parenchymal cells originates hepatocytes with the same chromosomal pattern as the parent stem cell (XY), and such identity is maintained also in the case of polyploidy, which is common in the liver (e.g. XYXY in tetraploid cells).

B. By contrast, cell fusion between male hematopoietic cell lineage (XY) and female hepatocytes (XX) results in a XXXY pattern, as it was shown in the Lagasse model.⁵⁸

UBC,⁷⁹ after inducing liver injury by allyl-alcohol,⁸⁰ and observed that the human cell population contributed to hepatic regeneration. Kogler *et al.*,⁸¹ isolated a CD45-negative population from human UCB (denominated «Unrestricted Somatic Stem Cells») exhibiting both pluripotency and a high proliferation capacity *in vitro*. Under appropriate conditions, these cells differentiate into osteoblasts, chondroblasts, adipocytes, hematopoietic and neural cells. When transplanted in the preimmune fetal sheep, they were able to generate albumin-producing human parenchymal liver cells. By analyzing liver parenchymal cells for the coexistence of human and ovine genomes, the Authors conclude that they were the result of differentiation rather than of fusion, although the latter event could not be completely excluded.

Epithelial cells from the amnion express markers of neural progenitors cells,^{82,83} as well as hepatic specific markers such as albumin and alpha-fetoprotein.⁸⁴ Furthermore, cells derived from amniotic fluid have a low immunogenicity, are phenotypically similar to MSCs from the BM⁸⁵ and are able to engraft in different tissues, including the liver.⁸⁶ Interestingly, amniotic mesenchymal cells appear to induce immunological tolerance, a property that might limit the need for immunosuppression in allogeneic transplantation.⁸⁶

Do hepatic stem cells participate in the parenchymal regeneration process associated with acute or chronic liver injury?

Several studies have addressed the ability of stem/progenitor cells to repopulate a diseased liver. In patients with chronic hepatitis C, Tanja Roskams was able to follow the differentiation of hepatic progenitor cells both into hepatocytes and cholangiocytes, suggesting that this stem cell compartment participates in the parenchymal regeneration associated with chronic viral liver disease.⁸⁷ Further studies in acute and chronic liver disease of different etiology also demonstrated differentiation of progenitor cells into hepatocytes.⁸⁸⁻⁹⁵ The activation of the stem/precursor cells compartment seems to be correlated with the severity of the disease.^{96,97} Activation of progenitor cells in chronic liver diseases implies that they form a potential target cell population for hepatocarcinogens,^{87,98-100} a caveat which should be taken into account if such cells are to be considered as a therapeutic tool.¹⁰¹ The differentiation of the progenitor cells into hepatocytes or biliary cells depends on the type of mature epithelial cell that is damaged^{35,88,95} and by the remodeling of the surrounding matrix.^{35,102} However, the factors contributing to the regulation of progenitor cell activation and the components that define the so called «stem cell niche» for adult human liver progenitor cells are poorly characterized.¹⁰³

The contribution of the BM to liver regeneration is less clear. Several reports in different animal models indicate that the participation of BM-derived hepatocytes to hepatic

parenchymal regeneration is insignificant,¹⁰⁴⁻¹⁰⁷ with the notable exception of the previously described work by Jang *et al.*⁵⁹ Theise *et al.* infused CD34⁺ lin⁻ BM cells from male mice into irradiate female mice, and found that only 0.39-1.1% of hepatocytes derived from BM. Following transplantation of BM cells into five irradiate mice, no hepatocytes of BM origin was observed in the liver of recipient animals while in two mice 0.4-2.2% of bile duct cells were positive for the Y chromosome.¹⁰⁸ Wagers *et al.*,¹⁰⁴ generated chimeric animals by transplantation of a single green fluorescent protein (GFP)-marked HSC into irradiate mice. Only one hepatocyte out of 70,000 was found to be GFP(+) in the recipient livers. Similarly, Fuji *et al.*,¹⁰⁵ were unable to identify BM-derived hepatocytes following transplantation of GFP(+) BM cells into GFP(-) hepatectomized mice. Kanazawa and Verma¹⁰⁶ tested three different animal models of liver injury (CCl₄ treatment, albumin-urokinase transgenic mouse and hepatitis B transgenic mouse) and found that only 5/410,000 cells were derived from BM. Finally, Dahlke *et al.*,¹⁰⁷ were unable to demonstrate any contribution of the BM to liver regeneration in a rodent model of CCl₄ liver injury associated with retrorsine administration, in order to inhibit the replication of endogenous hepatocytes. In studies on patients with sex-discordant liver or BM transplantation, the contribution of BM to hepatic parenchyma seems also to be absent or minimal.^{27,28,109-114} The frequency of BM-derived hepatocytes in the different studies was in the range of 0.5-2%,^{28,112,113} 1-8%,²⁷ and 4-7%,¹⁰⁹ respectively. Such discordant but mainly discouraging findings probably originate from very different (and sometimes inappropriate) experimental set-ups. Possible factors influencing liver repopulation with BM-derived hepatic parenchymal cells include the model and timing of liver injury, the route of stem cells administration (systemic vs intraportal) and the selection of the stem/precursor cell population as well as its activation before infusion.

What is the role of stem cells in hepatic fibrogenesis?

Myofibroblasts play a key role in the inflammatory response and in the process of hepatic fibrogenesis, due to their capacity to produce extracellular matrix.¹¹⁵ Working in collaboration with Malcolm Alison and Stuart Forbes and using the Y chromosome as a marker, we were able to identify recipient-derived myofibroblasts in liver grafts following sex-mismatched transplantation.¹¹⁶ These preliminary data were later confirmed in a larger series, suggesting a contribution of the BM to the fibrogenetic process leading to liver cirrhosis and pointing to a possible negative effect of BM-derived cell transplantation in liver disease.¹¹⁷ However, experimental work in rodents indicates that a specific cell population in the BM may actually prevent the fibrotic process in the liver. Systemic infusion of a subpopulation of BM-derived nonhematopoietic

cells, separated using an anti-Liv8 antibody, resulted in the resolution of liver fibrosis induced by CCl_4 treatment and normalized the synthetic function of the liver.¹¹⁸ These reports suggest that the BM may be actively involved in hepatic fibrogenesis, but its role in promoting or preventing the scarring process has yet to be defined.

Can the regenerative potential of stem cells be exploited for the treatment of liver diseases?

The treatment of liver disease with BM-derived hepatic stem cells might have considerable advantages over the use of hepatocytes. BM can be obtained from millions of potential living donors with simple procedures, in contrast to obtaining hepatocytes from the few cadaveric livers rejected for whole organ transplantation but still suitable for cell isolation. It was postulated that a population of stem cells resident in the BM can be released into the circulation, in response to stimuli derived from injured tissue, migrate to injured site and participate in regeneration.¹¹⁹⁻¹²³ We hypothesize that such cells represent the vestiges of a very ancient body repair system present in more primitive life forms. During evolution, with the development of more complex organisms, such a system probably became obsolete and was mostly replaced by more efficient, specific tissue/organ stem/precursor cells. In the light of this hypothesis, it is not surprising that the participation of BM-derived cells to non-hematopoietic tissue has often been described as insignificant. However, we could probably take advantage of the peculiar characteristics of such cells by concentrating them in the injured tissue and providing the optimal conditions to promote their participation in the regenerative process.

With respect to the cell population, about twenty different phenotypes of BM- or UCB-derived cells with potential for hepatic differentiation have been identified using a variety of surface markers.²⁰ It is reasonable to assume that some degree of overlap exists among the different cell populations, and there is much need for a joined effort in order to select a single phenotype including the most significant markers and demonstrating the most convincing and reproducible potential for hepatic differentiation. Clearly, a similar approach should also be applied to isolate intra-hepatic «resident» stem/precursor cells.

Following the pioneering work of Nancy Rolando^{124,125} as well as similar applications in the fields of Cardiology,¹²⁶ several investigators have approached the use of G-CSF as a method to mobilize from the BM stem/precursor cells with the aim to induce hepatic colonization improving parenchymal regeneration both in acute and in chronic liver disease. However, no convincing data have been published so far, suggesting that a significant therapeutic effect has yet to be demonstrated. Meanwhile, studies in laboratory animals have shown that the improvement in hepatic regeneration associated with G-CSF administration was not associated with increased liver repopulation

by BM-derived parenchymal cells, but rather to a more efficient repair process mediated by resident hepatic parenchymal cells.¹²⁷ Probably the most convincing clinical evidence for a possible therapeutic application of liver stem cells was published recently by am Esch II *et al.*¹²⁸ These Authors infused autologous CD133⁺ BM-derived cells into the portal vein following partial portal embolization, and they observed a significant improvement in hepatic regeneration with respect to the control group, which did not receive cell infusion.

Recent work, performed in collaboration between our laboratory and the Laboratory of Surgical Research of the Cedars-Sinai Medical Center in Los Angeles, led to successful isolation and characterization of a putative subpopulation of $\beta 2$ -microglobulin / Thy1⁺ hepatic stem cells both from the liver and from the BM.^{129,130} Selective intraportal infusion of syngeneic $\beta 2$ -microglobulin / Thy1⁺ BM-derived cells, following allogeneic liver transplantation in rats with subtherapeutic immunosuppression, resulted in up to $62 \pm 5.0\%$ repopulation of the transplanted lobes with syngeneic hepatocytes and cholangiocytes. Moreover, the survival of the animals which received cell infusion was doubled with respect to control group, suggesting that graft repopulation with BM-derived cells can rescue liver grafts undergoing rejection. We then used reversible ischemia/reperfusion liver injury to induce engraftment and hepatic parenchymal differentiation of exogenous $\beta 2$ -microglobulin/Thy1⁺ BM-derived cells.¹³¹ Transplantation of BM-derived cells obtained from GFP-transgenic rats into Lewis rats resulted in the presence of up to 20% of GFP(+) hepatocytes in ischemia/reperfusion-injured liver lobes after one month. Infusion of wild-type BM-derived cells into GFP-transgenic rats resulted in the appearance of GFP(-) hepatocytes, suggesting that the main mechanism underlying parenchymal repopulation was differentiation rather than cell fusion. Transplantation of wild-type BM-derived cells into hyperbilirubinemic Gunn rats with deficient bilirubin conjugation after ischemia/reperfusion damage resulted in 30% decrease of serum bilirubin, in the appearance of bilirubin conjugates in bile and in the expression of normal UDP-glucuronyltransferase enzyme, evaluated by PCR analysis. Thus, reversible ischemia/reperfusion injury induced hepatic parenchymal engraftment and differentiation into mature hepatocytes of BM-derived cells, and transplantation of BM-derived cells from non affected animals resulted in the partial correction of hyperbilirubinemia in the Gunn rat, suggesting that this procedure could potentially be used for the treatment of inherited metabolic liver diseases.

Conclusions

The use of isolated viable cells to restore the function of organs and tissues is emerging as a promising therapeutic tool. In the field of Hepatology, multiple sources of hepatic stem/precursor cells have been identified, and pre-

liminary evidence of therapeutic effectiveness has been provided in animal models. However, we have to learn more on the mechanisms of liver regeneration, including the role of stem/precursor cells. Definite (and possibly joined) protocols for the selection of a specific cell population and for *in vitro* expansion/differentiation should be developed, as well as protocols for clinical liver repopulation. The long-term fate of the transplanted cells should also be assessed in animal models, with respect to function, possible extra-hepatic localization, genetic/epigenetic stability and especially tumorigenesis. The risk is that superficial planning, without adequate consideration and knowledge of the underlying pathophysiology, will result in poorly focused clinical trials and possible complications, which could in turn originate skepticism on the development of cell therapy in Hepatology. Even if the biological characteristics of hepatic stem cells still justify the hope for successful future clinical applications, only a more cautious and systematic «bench to bedside» approach will guarantee consistent results.

Acknowledgment

We thank Professor Giuseppe Realdi and Professor Claudio Tiribelli for helpful discussion.

References

1. Sipe JD. Tissue Engineering and Reparative Medicine. In: Sipe JD, Kelley CA, McNicol LA, eds. *Reparative Medicine: Growing Tissues and Organs*. Volume 961, Annals of the New York Academy of Sciences, 2002; 1-9.
2. Lake JR. Hepatocyte transplantation. *N Engl J Med* 1998; 338: 1463-5.
3. Gupta S, Bhargava KK, Novikoff PM. Mechanisms of cell engraftment during liver repopulation with hepatocyte transplantation. *Semin Liver Dis* 1999; 19: 15-26.
4. Strom SC, Chowdhury JR, Fox IJ. Hepatocyte transplantation for the treatment of human disease. *Semin Liver Dis* 1999; 39-48.
5. Grompe M. Liver repopulation for the treatment of metabolic diseases. *J Inher Metab Dis* 2001; 24: 231-44.
6. Horslen SP, Fox IJ. Hepatocyte transplantation. *Transplantation* 2004; 77: 1481-6.
7. Muraca M, Neri D, Parenti A, Feltracco P, Granato A, Vilei MT, Ferraresso C, et al. Intraportal hepatocyte transplantation in the pig. A hemodynamic and histopathological study. *Transplantation* 2002; 73: 890-6.
8. Fox IJ, Chowdhury JR, Kaufman SS, Goertzen TC, Chowdhury NR, Warkentin PI, Dorko K, et al. Treatment of the Crigler-Najjar syndrome type I with hepatocyte transplantation. *N Engl J Med* 1998; 338: 1422-6.
9. Strom SC, Fisher RA, Thompson MT, Sanyal AJ, Cole PE, Ham JM, Posner MP. Hepatocyte transplantation as a bridge to orthotopic liver transplantation in terminal liver failure. *Transplantation* 1997; 63: 559-69.
10. Muraca M, Gerunda G, Neri D, Vilei MT, Granato A, Feltracco P, Meroni M, et al. Hepatocyte transplantation as a treatment for glycogen storage disease type 1a. *Lancet* 2002; 359: 1528.
11. Strom SC, Rubinstein WS, Barranger JA, Towbin RB, Charron M, Miele L, Pisarov LA, et al. Transplantation of human hepatocytes. *Transplantation Proc* 1997; 29: 2103-6.
12. Mitry RR, Dhawan A, Hughes RD, Bansal S, Lehec S, Terry C, Heaton ND, et al. One liver, three recipients: segment IV from split-liver procedures as a source of hepatocytes for cell transplantation. *Transplantation* 2004; 77: 1614-6.
13. Dhawan A, Mitry RR, Hughes RD, Lehec S, Terry C, Bansal S, Arya R, et al. Hepatocyte transplantation for inherited factor VII deficiency. *Transplantation* 2004; 78: 1812-4.
14. Bilir BM, Guinette D, Karrer F, Kumpe DA, Krysl J, Stephens J, et al. Hepatocyte transplantation in acute liver failure. *Liver Transpl* 2000; 6: 41-3.
15. Stephenne X, Najimi M, Sibille C, Nassogne MC, Smets F, Sokal EM. Sustained engraftment and tissue enzyme activity after liver cell transplantation for argininosuccinate lyase deficiency. *Gastroenterology* 2006; 130: 1317-23.
16. Rosenthal N. Prometheus's vulture and the stem-cell promise. *N Engl J Med* 2003; 349: 267-74.
17. Weissman IL. Translating stem and progenitor cell biology to the clinic: barriers and opportunities. *Science* 2000; 287: 1442-6.
18. Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, Jones JM. Embryonic stem cell lines derived from human blastocysts. *Science* 1998; 282: 1145-7.
19. Korbly M, Estrov Z. Adult stem cells for tissue repair - a new therapeutic concept? *N Engl J Med* 2003; 349: 570-82.
20. Laurson J, Selden C, Hodgson HJF. Hepatocyte progenitors in man and in rodents-multiple pathways, multiple candidates. *Int J Path* 2005; 86: 1-18.
21. Ferrari G, Cusella-De Angelis G, Coletta M, Paolucci E, Stornaiuolo A, Cossu G, Mavilio F. Muscle regeneration by bone marrow-derived myogenic progenitors. *Science* 1998; 279: 1528-30.
22. Petersen BE, Bowen WC, Patrene KD, Mars WM, Sullivan AK, Murase N, Boggs SS, et al. Bone marrow as potential source of hepatic oval cells. *Science* 1999; 284: 1168-70.
23. Jackson KA, Mi T, Goodell MA. Hematopoietic potential of stem cells isolated from murine skeletal muscle. *Proc Natl Acad Sci USA* 1999; 96: 14482-6.
24. Bjornson CR, Rietze RL, Reynolds BA, Magli MC, Vescovi AL. Turning brain into blood: a hematopoietic fate adopted by adult neural stem cells *in vivo*. *Science* 1999; 283.
25. Woodbury D, Schwarz EJ, Prockop DJ, Black IB. Adult rat and human bone marrow stromal cells differentiate into neurons. *J Neurosci Res* 2000; 61: 364-70.
26. Mezey E, Chandross KJ. Bone marrow: a possible alternative source of cells in the adult nervous system. *Eur J Pharmacol* 2000; 405: 297-302.
27. Theise ND, Nimmakayalu M, Gardner R, Illei PB, Morgan G, Teperman L, Henegariu O, et al. Liver from bone marrow in humans. *Hepatology* 2000; 32: 11-6.
28. Alison MR, Poulsom R, Jeffery R, Dhillon AP, Quaglia A, Jacob J, Novelli M, et al. Hepatocytes from non-hepatic adult stem cells. *Nature* 2000; 406: 257.
29. Lagasse E, Connors H, Al-Dhalimy M, Reitsma M, Dohse M, Osborne L, Wang X, et al. Purified hematopoietic stem cells can differentiate into hepatocytes *in vivo*. *Nat Med* 2000; 6: 1229-34.
30. Fausto N. Liver Regeneration and Repair: Hepatocytes, Progenitor Cells, and Stem Cells. *Hepatology* 2004; 39: 1477-87.
31. Farber E. Similarities in the sequence of early histological changes induced in the liver of the rat by ethionine, 2-acetylaminofluorene, and 3'-methyl-4-dimethylaminoazobenzene. *Cancer Res* 1956; 16: 142-8.
32. Grisham JW, Hartroft WS. Morphologic identification by electron microscopy of «oval» cells in experimental hepatic degeneration. *Lab Invest* 1961; 10: 317-32.
33. Evarts RP, Nagy P, Nakatsukasa H, Marsden E, Thorgeirsson SS. *In vivo* differentiation of rat liver oval cells into hepatocytes. *Cancer Res* 1989; 49: 1541-7.
34. Yin L, Lynch D, Ilic Z, Sell S. Proliferation and differentiation of ductular progenitor cells and littoral cells during the regeneration of the rat liver to CCl₄/2-AAF injury. *Histol Histopathol* 2002; 17: 65-81.

35. Roskams TA, Libbrecht L, Desmet VJ. Progenitor cells in diseased human liver. *Semin Liver Dis* 2003; 23: 385-96.
36. Libbrecht L, Roskams T. Hepatic progenitor cells in human liver diseases. *Semin Cell Dev Biol* 2002; 13: 389-96.
37. Germain L, Goyette R, Marceau N. Differential cyokeratin and alpha-fetoprotein expression in morphologically distinct epithelial cells emerging at the early stage of rat hepatocarcinogenesis. *Cancer Res* 1985; 45: 673-81.
38. Hixson DC, Faris RA, Yang L, Novikoff P. Antigenic clues to liver development, renewal, and carcinogenesis: a integrated model. In: Sirica, AE eds. *The role of cell types in hepatocarcinogenesis*, CRC Press, Boca Raton, FL. 1992: 151-182.
39. Dabeva MD, Alpini G, Hurston E, Shafritz DA. Models for hepatic progenitor cell activation. *Proc Soc Exp Biol Med* 1993; 204: 242-52.
40. Fausto N, Campbell JS. The role of hepatocytes and oval cells in liver regeneration and repopulation. *Mech Dev* 2003; 120: 117-30.
41. Dabeva MD, Laconi E, Oren R, Petkov PM, Hurston E, Shafritz DA. Liver regeneration and alpha-fetoprotein messenger RNA expression in the retrorsine model for hepatocyte transplantation. *Cancer Res* 1998; 58: 5825-34.
42. Gordon GJ, Coleman WB, Hixson DC, Grisham JW. Liver regeneration in rats with retrorsine-induced hepatocellular injury proceeds through a novel cellular response. *Am J Pathol* 2000; 156: 607-19.
43. Vig P, Russo FP, Edwards RJ, Tadrous PJ, Wright NA, Thomas HC, Alison MR, et al. The sources of parenchymal regeneration after chronic hepatocellular liver injury in mice. *Hepatology* 2006; 43: 316-24.
44. Laconi E, Sarma DS, Pani P. Transplantation of normal hepatocytes modulates the development of chronic liver lesions induced by a pyrrolizidine alkaloid, lasiocarpine. *Carcinogenesis* 1995; 16: 139-42.
45. Laconi E, Oren R, Mukhopadhyay DK, Hurston E, Laconi S, Pani P, Dabeva MD, et al. Long-term, near-total liver replacement by transplantation of isolated hepatocytes in rats treated with retrorsine. *Am J Pathol* 1998; 153: 319-29.
46. Gordon GJ, Coleman WB, Grisham JW. Temporal analysis of hepatocyte differentiation by small hepatocyte-like progenitor cells during liver regeneration in retrorsine-exposed rats. *Am J Pathol* 2000; 157: 771-86.
47. Avril A, Pichard V, Bralet MP, Ferry N. Mature hepatocytes are the source of small hepatocyte-like progenitor cells in the retrorsine model of liver injury. *J Hepatol* 2004; 41: 737-43.
48. Gordon GJ, Coleman WB, Grisham JW. Induction of cytochrome P450 enzymes in the livers of rats treated with the pyrrolizidine alkaloid retrorsine. *Exp Mol Pathol* 2000; 69: 17-26.
49. Mattocks AR. *Chemistry and toxicology of pyrrolizidine alkaloids*. London/Orlando, FL: Academic Press, 1986.
50. Fujio K, Evarts RP, Hu Z, Marsden ER, Thorgeirsson SS. Expression of stem cell factor and its receptor, c-kit, during liver regeneration from putative stem cells in adult rat. *Lab Invest* 1994; 70: 511-6.
51. Kabrun N, Buhring HJ, Choi K, Ullrich A, Risau W, Keller G. Flk-1 expression defines a population of early embryonic hematopoietic precursors. *Development* 1997; 124: 2039-48.
52. Omori M, Evarts RP, Omori N, Hu Z, Marsden ER, Thorgeirsson SS. Expression of alpha-fetoprotein and stem cell factor/c-kit system in bile duct ligated young rats. *Hepatology* 1997; 25: 1115-22.
53. Petersen BE, Goff JP, Greenberger JS, Michalopoulos GK. Hepatic oval cells express the hematopoietic stem cell marker Thy-1 in the rat. *Hepatology* 1998; 27: 433-45.
54. Petersen BE, Grossbard B, Hatch H, Pi L, Deng J, Scott EW. Mouse A6-positive hepatic oval cells also express several hematopoietic stem cell markers. *Hepatology* 2003; 37: 632-40.
55. Craig W, Kay R, Cutler RL, Lansdorp PM. Expression of Thy-1 on human hematopoietic progenitor cells. *J Exp Med* 1993; 177: 1331-42.
56. Taniguchi H, Toyoshima T, Fukao K, Nakauchi H. Presence of hematopoietic stem cells in the adult liver. *Nat Med* 1996; 2: 198-203.
57. Wang X, Ge S, McNamara G, Hao QL, Crooks GM, Nolte JA. Albumin expressing hepatocyte like cells develop in the livers of immune deficient mice that receive transplant of highly purified human hematopoietic stem cells. *Blood* 2003; 101: 4201-8.
58. Vassilopoulos G, Wang PR, Russell DW. Transplanted bone marrow regenerates liver by cell fusion. *Nature* 2003; 422: 901-4.
59. Jang YY, Collector MI, Baylin SB, Diehl AM, Sharkis SJ. Hematopoietic stem cells convert into liver cells within days without fusion. *Nat Cell Biol* 2004; 6: 532-9.
60. Schwartz RE, Reyes M, Koodie L, Jiang Y, Blackstad M, Lund T, Lenvik T, et al. Multipotent adult progenitor cells from bone marrow differentiate into functional hepatocyte-like cells. *J Clin Invest* 2002; 109: 1291-302.
61. Lee KD, Kuo TK, Whang-Peng J, Chung YF, Lin CT, Chou SH, Chen JR, et al. *In vitro* hepatic differentiation of human mesenchymal stem cells. *Hepatology* 2004; 40: 1275-84.
62. Shi XL, Qiu YD, Wu XY, Xie T, Zhu ZH, Chen LL, Li L, et al. *In vitro* differentiation of mouse bone marrow mononuclear cells into hepatocyte-like cells. *Hepatol Res* 2005; 31: 223-31.
63. Menthena A, Deb N, Oertel M, Grozdanov PN, Sandhu J, Shah S, Guha C, et al. Bone marrow progenitors are not the source of expanding oval cells in injured liver. *Stem Cells* 2004; 22: 1049-61.
64. Erices A, Conget P. Mesenchymal progenitor cells in human umbilical cord blood. *Br J Haematol* 2000; 109: 235-42.
65. Lee OK, Kuo TK, Chen WM, Lee KD, Hsieh SL, Chen TH. Isolation of multipotent mesenchymal stem cells from umbilical cord blood. *Blood* 2004; 103: 1669-75.
66. Broxmeyer HE, Douglas GW, Hangoc G, Cooper S, Bard J, English D, Army M, et al. Human umbilical cord blood as a potential source of transplantable hematopoietic stem/progenitor cells. *Proc Natl Acad Sci USA* 1989; 86: 3828-32.
67. Gluckman E, Broxmeyer HA, Auerbach AD, Friedman HS, Douglas GW, Devergie A, Esperou H, et al. Hematopoietic reconstitution in a patient with Fanconi's anemia by means of umbilical-cord blood from an HLA-identical sibling. *N Engl J Med* 1989; 321: 1174-8.
68. Kurtzberg J, Laughlin M, Graham ML, Smith C, Olson JF, Halperin EC, Ciocci G, et al. Placental blood as a source of hematopoietic stem cells for transplantation into unrelated recipients. *N Engl J Med* 1996; 335: 157-66.
69. Gluckman E, Rocha V, Boyer-Chammard A, Locatelli F, Arcese W, Pasquini R, et al. Outcome of cord-blood transplantation from related and unrelated donors. Eurocord Transplant Group and the European Blood and Marrow Transplantation Group. *N Engl J Med* 1997; 337: 373-81.
70. Wagner JE, Kernan NA, Steinbuch M, Broxmeyer HE, Gluckman E. Allogeneic sibling umbilical-cord-blood transplantation in children with malignant and non-malignant disease. *Lancet* 1995; 346: 214-9.
71. Garbuzova-Davis S, Willing AE, Zigova T, Saporta S, Justen EB, Lane JC, Hudson JE, Chen N, Davis CD, Sanberg PR. Intravenous administration of human umbilical cord blood cells in a mouse model of amyotrophic lateral sclerosis: distribution, migration, and differentiation. *J Hematother Stem Cell Res* 2003; 12: 255-70.
72. Smith S, Broxmeyer HE. The influence of oxygen tension on the long-term growth *in vitro* of haematopoietic progenitor cells from human cord blood. *Br J Haematol* 1986; 63: 29-34.
73. Gluckman E. Current status of umbilical cord blood hematopoietic stem cell transplantation. *Exp Hematol* 2000; 28: 1197-205.
74. Salahuddin SZ, Markham PD, Ruscetti FW, Gallo RC. Long-term suspension cultures of human cord blood myeloid cells. *Blood* 1981; 58: 931-8.
75. Kakinuma S, Tanaka Y, Chinzei R, Watanabe M, Shimizu-Saito K, Hara Y, et al. Human umbilical cord blood as a source of transplantable hepatic progenitor cells. *Stem Cells* 2003; 21: 217-27.
76. Sharma AD, Cantz T, Richter R, Eckert K, Henschler R, Wilkens AJR, Arseniev L. Human cord blood stem cells generate human cyokeratin 18-negative hepatocyte-like cells in injured mouse liver. *Am J Pathol* 2005; 167: 555-64.

77. Newsome PN, Johannessen I, Boyle S, Dalakas E, McAulay KA, Samuel K, Rae F, et al. Human cord blood-derived cells can differentiate into hepatocytes in the mouse liver with no evidence of cellular fusion. *Gastroenterology* 2003; 124: 1891-900.
78. Piscaglia AC, Di Campli C, Zocco MA, Di Gioacchino G, Novi M, Rutella S, Bonanno G, et al. Human cordonal stem cell intra-peritoneal injection can represent a rescue therapy after an acute hepatic damage in immunocompetent rats. *Transplant Proc* 2005; 37: 2711-4.
79. Rutella S, Bonanno G, Marone M, De Ritis D, Mariotti A, Voso MT, Scambia G, et al. Identification of a novel subpopulation of human cord blood CD34-CD133-CD7-CD45+lineage- cells capable of lymphoid/NK cell differentiation after in vitro exposure to IL-15. *J Immunol* 2003; 171: 2977-88.
80. Di Campli C, Piscaglia AC, Pierelli L, Rutella S, Bonanno G, Alison MR, Mariotti A, et al. Human umbilical cord stem cell rescue therapy in a murine model of toxic liver injury. *Dig Liver Dis* 2004; 36: 603-13.
81. Kogler G, Sensken S, Airey JA, Trapp T, Muschen M, Feldhahn N, Liedtke S, et al. A new human somatic stem cell from placental cord blood with intrinsic pluripotent differentiation potential. *J Exp Med* 2004; 200: 123-35.
82. Ishii T, Ohsugi K, Nakamura S, Sato K, Hashimoto M, Mikoshiba K, Sakuragawa N. Gene expression of oligodendrocyte markers in human amniotic epithelial cells using neural cell-type-specific expression system. *Neurosci Lett* 1999; 268: 131-4.
83. Sakuragawa N, Thangavel R, Mizuguchi M, Hirasawa M, Kamo I. Expression of markers for both neuronal and glial cells in human amniotic epithelial cells. *Neurosci Lett* 1996; 209: 9-12.
84. Sakuragawa N, Enosawa S, Ishii T, Thangavel R, Tashiro T, Okuyama T, Suzuki S. Human amniotic epithelial cells are promising transgene carriers for allogeneic cells transplantation into liver. *J Hum Genet* 2000; 45: 171.
85. In 't Anker PS, Scherjon SA, Kleijburg-van der Keur C, Noort WA, Claas FH, Willemze R, Fibbe WE, Kanhai HH. Amniotic fluid as a novel source of mesenchymal stem cells for therapeutic transplantation. *Blood* 2003; 102: 1548-9.
86. Bailo M, Soncini M, Vertua E, Bonassi Signoroni P, Sanzone S, Lombardi G, et al. Engraftment potential of human and corion cells derived from placenta. *Transplantation* 2004; 78: 1439-48.
87. Libbrecht L, Desmet V, Van Damme B, Roskams T. The immunohistochemical phenotype of dysplastic foci in human liver: correlation with putative progenitor cells. *J Hepatol* 2000; 33: 76-84.
88. Roskams T, De Vos R, Van Eyken P, Myazaki H, Van Damme B, Desmet V. Hepatic OV-6 expression in human liver disease and rat experiments: evidence for hepatic progenitor cells in man. *J Hepatol* 1998; 29: 455-63.
89. Haque S, Haruna Y, Saito K, Nalesnik MA, Atillasoy E, Thung SN, Gerber MA. Identification of bipotential progenitor cells in human liver regeneration. *Lab Invest* 1996; 75: 699-705.
90. Roskams T, De Vos R, van den Oord JJ, Desmet V. Cells with neuroendocrine features in regenerating human liver. *APMIS Suppl.* 1991; 23: 32-9.
91. Gerber MA, Thung SN, Shen S, Stromeyer FW, Ishak KG. Phenotypic characterization of hepatic proliferation. Antigenic expression by proliferating epithelial cells in fetal liver, massive hepatic necrosis, and nodular transformation of the liver. *Am J Pathol* 1983; 110: 70-4.
92. Lowes KN, Brennan BA, Yeoh GC, Olynyk JK. Oval cell numbers in human chronic liver diseases are directly related to disease severity. *Am J Pathol* 1999; 154: 537-41.
93. Fujita M, Furukawa H, Hattori M, Todo S, Ishida Y, Nagashima K. Sequential observation of liver cell regeneration after massive hepatic necrosis in auxiliary partial orthotopic liver transplantation. *Mod Pathol* 2000; 13: 152-7.
94. Ray MB, Mendenhall CL, French SW, Gartside PS. Bile duct changes in alcoholic liver disease. The Veterans Administration Cooperative Study Group. *Liver* 1993; 13: 36-45.
95. Roskams T, Yang SQ, Koteish A, Durnez A, DeVos R, Huang X, Achten R, et al. Oxidative stress and oval cell accumulation in mice and humans with alcoholic and nonalcoholic fatty liver disease. *Am J Pathol* 2003; 163: 1301-11.
96. Libbrecht L, Desmet V, Van Damme B, Roskams T. Deep intralobular extension of human hepatic 'progenitor cells' correlates with parenchymal inflammation in chronic viral hepatitis: can 'progenitor cells' migrate? *J Pathol* 2000; 192: 373-8.
97. Xiao JC, Ruck P, Adam A, Wang TX, Kaiserling E. Small epithelial cells in human liver cirrhosis exhibit features of hepatic stem-like cells: immunohistochemical, electron microscopic and immunoelectron microscopic findings. *Histopathology* 2003; 42: 141-9.
98. Hsia CC, Evarts RP, Nakatsukasa H, Marsden ER, Thorgeirsson SS. Occurrence of oval-type cells in hepatitis B virus-associated human hepatocarcinogenesis. *Hepatology* 1992; 16: 1327-33.
99. Libbrecht L, Craninx M, Nevens F, Desmet V, Roskams T. Predictive value of liver cell dysplasia for development of hepatocellular carcinoma in patients with non-cirrhotic and cirrhotic chronic viral hepatitis. *Histopathology* 2001; 39: 66-73.
100. Libbrecht L, De Vos R, Cassiman D, Desmet V, Aerts R, Roskams T. Hepatic progenitor cells in hepatocellular adenomas. *Am J Surg Pathol* 2001; 25: 1388-96.
101. Roskams T. Progenitor cell involvement in cirrhotic human liver diseases: from controversy to consensus. *J Hepatol* 2003; 39: 431-4.
102. Desmet V. Organizational principles. In: Arias I, Boyer JL, Chisari F, et al., eds. *The Liver. Biology and pathobiology*. 4th ed. Philadelphia: Lippincott Williams and Wilkins, 2001: 3-15.
103. Roskams T. Acrobatic liver progenitor cells: mythology or truth? Abstract of the 5th meeting Gli incontri della verità: la terapia delle malattie epatiche. 16-17 March 2006, Rome, Italy. pags. 31-34.
104. Wagers AJ, Sherwood RI, Christensen JL, Weissman IL. Little evidence for developmental plasticity of adult hematopoietic stem cells. *Science* 2002; 297: 2256-9.
105. Fuji H, Hirose T, Oe S, Yasuchika K, Azuma H, Fujikawa T, Nagao M, et al. Contribution of bone marrow cells to liver regeneration after partial hepatectomy in mice. *J Hepatol* 2002; 36: 653-9.
106. Kanazawa Y, Verma IM. Little evidence of bone marrow-derived hepatocytes in the replacement of injured liver. *Proc Natl Acad Sci USA* 2003; 100 Suppl 1: 11850-3.
107. Dahlke MH, Popp FC, Bahlmann FH, Aselmann H, Jager MD, Neipp M, Piso P, et al. Liver regeneration in a retrorsine/CCl4-induced acute liver failure model: do bone marrow-derived cells contribute? *J Hepatol* 2003; 39: 365-73.
108. Krause DS, Theise ND, Collector MI, Henegariu O, Hwang S, Gardner R, Neutzel S, et al. Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell. *Cell* 2001; 105: 369-77.
109. Korbiling M, Katz RN, Khanna A, Ruifrok AC, Rondon G, Albitar M, Champlin RE, et al. Hepatocytes and epithelial cells of donor origin in recipients of peripheral-blood stem cells. *N Engl J Med* 2002; 346: 738-46.
110. Fogt F, Beyser KH, Poremba C, Zimmerman RL, Khettry U, Ruschhoff J. Recipient-derived hepatocytes in liver transplants: a rare event in sex-mismatched transplants. *Hepatology* 2002; 36: 173-6.
111. Wu T, Cieply K, Nalesnik MA, Randhawa PS, Sonzogni A, Bellamy C, Abu-Elmagd K, et al. Minimal evidence of transdifferentiation from recipient bone marrow to parenchymal cells in regenerating and long-surviving human allografts. *Am J Transplant* 2003; 3: 173-81.
112. Ng IO, Chan KL, Shek WH, Lee JM, Fong DY, Lo CM, Fan ST. High frequency of chimerism in transplanted livers. *Hepatology* 2003; 38: 89-98.
113. Kleiberger W, Rothamel T, Glockner S, Flemming P, Lehmann U, Kreipe H. High frequency of epithelial chimerism in liver transplants demonstrated by microdissection and STR-analysis. *Hepatology* 2002; 35: 10-6.
114. Quintana-Bustamante O, Alvarez-Barrientos A, Kofman AV, Fabregat I, Bueren JA, Theise ND, Segovia JC. Hematopoietic

- mobilization in mice increases the presence of bone marrow-derived hepatocytes via *in vivo* cell fusion. *Hepatology* 2006; 43: 108-16.
115. Brenner DA, Waterboer T, Choi SK, Lindquist JN, Stefanovic B, Burchardt E, Yamauchi M, et al. New aspects of hepatic fibrosis. *J Hepatol* 2000; 32: 32-8.
116. Granato A, Forbes S, Poulsom R, Muraca M, Quarta M, Rugge M, Alison M. The repopulation of liver allografts with recipient BM derived liver cells varies with time and between recipients. *J Hepatol* 2002; 36: 19 Abstract.
117. Forbes SJ, Russo FP, Rey V, Burra P, Rugge M, Wright NA, Alison MR. A significant proportion of myofibroblasts are of bone marrow origin in human liver fibrosis. *Gastroenterology* 2004; 126: 955-63.
118. Sakaida I, Terai S, Yamamoto N, Aoyama K, Ishikawa T, Nishina H, Okita K. Transplantation of bone marrow cells reduces CCl₄-induced liver fibrosis in mice. *Hepatology* 2004; 40: 1304-11.
119. Lapidot T, Petit I. Current understanding of stem cell mobilization: the roles of chemokines, proteolytic enzymes, adhesion molecules, cytokines, and stromal cells. *Exp Hematol* 2002; 30: 973-81.
120. Hatch HM, Zheng D, Jorgensen ML, Petersen BE. SDF-1 α /CXCR4: a mechanism for hepatic oval cell activation and bone marrow stem cell recruitment to the injured liver of rats. *Cloning Stem Cells* 2002; 4: 339-51.
121. Kollet O, Shivtiel S, Chen YQ, Suriawinata J, Thung SN, Dabeva MD, Kahn J, et al. HGF, SDF-1, and MMP-9 are involved in stress-induced human CD34⁺ stem cell recruitment to the liver. *J Clin Invest* 2003; 112: 160-9.
122. Dalakas E, Newsome PN, Harrison DJ, Plevris JN. Hematopoietic stem cell trafficking in liver injury. *FASEB J* 2005; 19: 1225-31.
123. Kucia M, Ratajczak J, Ratajczak MZ. Are bone marrow stem cells plastic or heterogeneous—that is the question. *Exp Hematol* 2005; 33: 613-23.
124. Rolando N, Clapperton M, Wade J, Panetsos G, Mufti G, Williams R. Granulocyte colony-stimulating factor improves function of neutrophils from patients with acute liver failure. *Eur J Gastroenterol Hepatol* 2000; 12: 1135-40.
125. Rolando N, Clapperton M, Wade J, Wendon J. Administering granulocyte colony-stimulating factor to acute liver failure patients corrects neutrophil defects. *Eur J Gastroenterol Hepatol* 2000; 12: 1323-8.
126. Orlic D, Kajstura J, Cimenti S, Limana F, Jakoniuk I, Quaini F, Nadal-Ginard B, et al. Mobilized bone marrow cells repair the infarcted heart, improving function and survival. *Proc Natl Acad Sci USA* 2001; 98: 10344-9.
127. Yannaki E, Athanasiou E, Xagorari A, Constantinou V, Batsis I, Kaloyannidis P, Proya E, et al. G-CSF-primed hematopoietic stem cells or G-CSF *per se* accelerate recovery and improve survival after liver injury, predominantly by promoting endogenous repair programs. *Exp Hematol* 2005; 33: 108-19.
128. am Esch JS 2nd, Knoefel WT, Klein M, Ghodsizad A, Fuerst G, Poll LW, Piechaczek C, et al. Portal application of autologous CD133⁺ bone marrow cells to the liver: a novel concept to support hepatic regeneration. *Stem Cells* 2005; 23: 463-70.
129. Avital I, Feraresso C, Aoki T, Hui T, Rozga J, Demetriou A, Muraca M. Bone marrow-derived liver stem cell and mature hepatocyte engraftment in livers undergoing rejection. *Surgery* 2002; 132: 384-90.
130. Avital I, Inderbitzin D, Aoki T, Tyan DB, Cohen AH, Feraresso C, Rozga J, et al. Isolation, characterization, and transplantation of bone marrow-derived hepatocyte stem cells. *Biochem Biophys Res Commun* 2001; 288:156-64.
131. Feraresso C, Granato A, Vilei MT, Quarta M, Cozzi E, Lanza C, Ruge M, et al. Bone Marrow-derived hepatic stem cells (BMSC) repopulate rat livers after the induction of ischemia/reperfusion injury and are able to correct a congenital metabolic disorder. *Abstract of the 40th Annual Meeting of the European Association for the study of the liver*, April 13-17, 2005, Paris, France, pag. 260.