



Images in Hepatology

Fused PET/CT Images in Hepatocarcinoma

Ivan F. Vega-González;¹ Ernesto Roldán-Valadez;¹ Guillermo Valdiviezo-Cárdenas¹




Figure 1.

¹ Unidad PET/CT. Fundación Clínica Médica Sur

Address for correspondence:

Ivan F. Vega-González

Unidad PET/CT. Puente de Piedra 150 Col. Toriello Guerra 14050
México, D.F. México.

Manuscript received and accepted: 31 August 2006

75-years old hispanic female with history of liver cirrhosis. It has been reported that the ^{18}F -FDG PET scan of the liver can depict false negative results particularly with history of liver cirrhosis (1, 2). In this case ^{18}F -FDG PET/CT depicted a large tumoral mass in segment VI (15.6 x 12.4 cm), there is enlargement of the liver with medial displacement of the gall-bladder. Our first differential diagnosis was hepatocarcinoma, pathologic analysis of liver biopsy confirmed the diagnosis.

Figures A-C, sagittal, coronal and axial planes PET/CT fused images show a big lesion in segment VI of the liver with increased uptake of ^{18}F -FDG, the maximum SUV was 14.5. The lesion presented two central zones of no uptake related with central necrosis. Figures D-F, oblique volumetric-3D fused PET/CT images depicting different aspects of the lesion.

References

1. Khan MA, Combs CS, Brunt EM, Lowe VJ, Wolverson MK, Solomon H, Collins BT, et al. Positron emission tomography scanning in the evaluation of hepatocellular carcinoma. *J Hepatol* 2000;32:792-797.
2. Son HB, Han CJ, Kim J, Jeong SH, Kim YC, Lee JO, Choi CY, et al. Evaluation of various hepatic lesions with positron emission tomography. *Taehan Kan Hakhoe Chi* 2002;8:472-480.