

## Images in Hepatology

## A 25 years old women after Kasai operation for biliary atresia

Héctor A. Baptista-González;<sup>1</sup> Guadalupe Ponciano-Rodríguez;<sup>2</sup> Misael Uribe;<sup>1</sup> Nahum Méndez-Sánchez<sup>1</sup>

Biliary atresia is a rare disease characterized by a biliary obstruction of unknown origin that presents in the neonatal period.<sup>1</sup> We present the magnetic resonance cholangiography (MRC) from a 25 years old women who was treated in the neonatal period with Kasai operation. The MRC reveals a choledochal cyst

and shows multiple saccular dilatation of both left and right hepatic ducts.

### References

1. Chardot C. Biliary atresia. *Orphanet J Rare Dis* 2006; 1: 28.



<sup>1</sup> Liver Unit. Medica Sur Clinic & Foundation. Mexico City, Mexico.

<sup>2</sup> Faculty of Medicine, National Autonomous University of Mexico, Mexico City, Mexico.

Address for correspondence:

Nahum Méndez-Sánchez, M.D., PhD.

Departments of Biomedical Research, Gastroenterology & Liver Unit. Medica Sur Clinic & Foundation, Puente de Piedra 150, Col. Toriello Guerra, Tlalpan 14050, Mexico City, Mexico.

Phone: (+525) 55606-6222 ext. 4215; Fax: (+525) 55666-4031;

E-mail: nmendez@medicasur.org.mx

Manuscript received and accepted: 2 August and 15 September 2008