

## Protective and curative effects of polyphenolic extracts from *Ichnocarpus frutescense* leaves on experimental hepatotoxicity by carbon tetrachloride and tamoxifen

Chidambaram Kumarappan,<sup>\*,†</sup> Madhavan Vijayakumar,<sup>‡</sup> Ellappan Thilagam,<sup>§</sup> Manikam Balamurugan,<sup>||</sup> Madheswaran Thiagarajan,<sup>¶</sup> Siddan Senthil,<sup>§</sup> Sunil C. Das,<sup>\*\*</sup> Subhash C. Mandal<sup>\*</sup>

<sup>\*</sup>Division of Pharmacognosy, Pharmacognosy and Phytotherapy Research Laboratory, Department of Pharmaceutical Technology, Faculty of Engineering and Technology, Jadavpur University, Kolkata 700 032, India.

<sup>†</sup>Department of Life sciences, School of Pharmacy & Health Sciences, International Medical University, Kuala Lumpur, Malaysia).

<sup>‡</sup>Ethnopharmacology Laboratory, National Botanical Research Institute, India).

<sup>§</sup>Department of Phytochemistry and Pharmacognosy, J.K.K.M.M.R.F. College of Pharmacy, B. Komarapalayam, India)

<sup>||</sup>Department of Pharmaceutics, University of Nizwa, Birkat Al Mouz, Nizwa, Sultanate of Oman.

<sup>¶</sup>Thiagarajan M, Clinical Pharmacy Lab, College of Pharmacy, Yeungnam University, South Korea).

<sup>\*\*</sup>Department of Plant Molecular Biology and Biotechnology, Entomology Research Institute, Loyola College (Autonomous), India).

### ABSTRACT

The aim of this study was to investigate prophylactic and curative effect of polyphenolic extract of *Ichnocarpus frutescense* against carbon tetrachloride and tamoxifen induced hepatotoxicity in rats. Carbon tetrachloride and tamoxifen caused liver damage in rats manifested by significant rise in serum enzymes levels. Models of carbon tetrachloride and tamoxifen intoxication elicited significant declines in the reduced glutathione concomitant with significant elevations in malondialdehyde levels. The oral administration of polyphenolic extract to carbon tetrachloride and tamoxifen intoxicated rats, produced significant increments in the reduced glutathione concomitant with significant decrements in malondialdehyde and liver transaminases levels. Prophylactic and curative treatments with the polyphenolic extract generally resulted in a relatively good protection against both carbon tetrachloride and tamoxifen intoxicated rats. The histopathological changes of liver sections showed an improved histological appearance. The extract inhibits CYP monooxygenases aminopyrine-*N*-demethylase and aniline hydroxylase, suggesting a plausible hepatoprotective mechanism. However prophylactic treatment with the polyphenolic extract exhibited a higher activity compared to curative treatment. The normalization of phenobarbitone induced sleeping time suggests the restoration of liver CYP enzymes. The study shows that hepatoprotective activity of polyphenol extract is by regulating the levels of hepatic microsomal drug metabolising enzymes. These results supported the use of this plant for the treatment of hepatitis in oriental traditional medicine.

**Key words.** *Ichnocarpus frutescens*. Hepatoprotection. Carbon tetrachloride. Tamoxifen. Polyphenolic extract. Phase I enzymes.

### INTRODUCTION

The liver is an important organ actively involved in metabolic and detoxification functions and is a frequent and primary target for several toxicants.<sup>1</sup> Ingestion of toxicants leads to liver cell necrosis in human as well as in experimental animals.<sup>2,3</sup> Drugs

continue to be removed from the market with disturbing regularity because of the late discovery of hepatotoxicity. Now, the incidence of acute exposure to xenobiotics has a significant trend to increase and becomes a hazard for the people. Tamoxifen citrate, is an anti-estrogenic, non-steroidal compound widely used for adjuvant therapy in breast cancer.<sup>4</sup> This strong hepatotoxic and hepatocarcinogenic effect of tamoxifen in rats raises the issues bearing on the prophylactic chronic administration to healthy women.<sup>5</sup> Tamoxifen is metabolized by human CYP 3A4 and 2D6 and is also an effective inducer of CYP 2B2, 2B1, and 3A in rat liver at doses comparable with the therapeutic doses used in humans.<sup>6,7</sup>

Carbon tetrachloride is highly volatile and is relatively stable in the environment. Carbon tetrachloride

Correspondence and reprint request: Dr. CT. Kumarappan, M.Pharm, PhD International Medical University No. 126, Jalan Jalil Perkasa 19, Bukit Jalil, 57000 Kuala Lumpur, Malaysia, HP:016-2242696, E-mail: ctkumarrx@gmail.com.

Manuscript received: August 07, 2010.  
Manuscript accepted: November 23, 2010.

de produces acute systemic toxicity following ingestion or inhalation. Symptoms of hepatic toxicity (5 ppm or greater) may appear after a delay of one to four days following acute exposure. Since  $\text{CCl}_4$  toxicity results from its bioactivation mainly by CYP2E1.<sup>8-10</sup> Toxicant-or drug-induced liver injury is a potential complication of nearly every medication. We needed to rule out the possibility that the observed hepatic toxicity was due to toxicant or therapeutic drugs for treatment. It is well known that a number of non-toxic herbs are having activities like membrane stabilizing, hepatoprotective, anti-oxidative and CYP2E1 inhibitory effects. Several herbal formulation prevented hepatotoxicity significantly and improved the disease outcome as well as patient compliance without any toxicity or side effects.<sup>11</sup>

Several endogenous protective mechanisms have been evolved to limit reactive oxygen species and the damage caused by them.<sup>12</sup> It was obvious that, certain toxicants (may be drug overdose or harmful substances) have been characterized by their ability to induce liver injury following the cleavage by CYP to form free radicals.<sup>13-14</sup> The body has developed mechanisms to remove free radicals. However, since this protection may not be complete, or when the formation of reactive oxygen species is excessive, additional protective mechanisms of dietary antioxidants may be of significance. Therefore, many natural and artificial agents possessing anti-oxidative properties have been proposed to prevent and treat hepatopathies induced by oxidative stress.<sup>15</sup> There is an increasing evidence for the hepatoprotective role of flavonoids, polyhydroxy organic compounds and terpenoids particularly from vegetables, fruits and some herbs.<sup>16-19</sup>

Leaves of *Ichnocarpus frutescens* (L.) R.Br. (*Apocynaceae*) boiled in gingelly oil are applied for headaches, fevers and wounds.<sup>20</sup> Survey of literature revealed that different pentacyclic triterpenoids and flavonoids have been isolated<sup>21-22</sup>. The utilization of decoction of leaves of *I. frutescens* in the treatment of jaundice, and diabetes is noteworthy.<sup>23-24</sup> Some of the active constituents of this plant, such as triterpenoids and flavonoids were shown to present hepatoprotective, antioxidant and related biological activities.<sup>25</sup> Our earlier report revealed the anti-inflammatory, free radical scavenging, anti-cancers and anti-diabetic activities of the polyphenolic extract of *I. frutescens*.<sup>26-28</sup> In spite of the reported use, no systematic experimental studies have been carried out to assess the hepatoprotective activity of *I. frutescens*. Therefore, the polyphenolic extract of *I. frutescens* leaves was investigated for its hepatoprotective effect to assess ethnomedical uses.

## MATERIALS AND METHODS

### Chemicals

Tamoxifen citrate was purchased from Sigma (St. Louis, USA). Phenobarbitone sodium was obtained from Nicholas Piramal India Limited (Mumbai, India). Assay kits aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase,  $\gamma$ -glutamyl transferase, total bilirubin and the total proteins were purchased from Span Diagnostic Ltd (Mumbai, India). Reduced glutathione, thiobarbituric acid and 5, 5'-dithiobis (2-nitrobenzoic acid) were purchased from SISCO Research Laboratories Pvt. Ltd (Mumbai, India). Carbon tetrachloride and dimethyl sulfoxide were obtained from Merck (Mumbai, India). All other chemicals were of high purity and obtained from commercial sources.

### Plant material

Fresh leaves of *Ichnocarpus frutescens* (L.) R.Br. (*Apocynaceae*) were collected from the delta region of the Cauvery river, Thiruchirappalli, India, during February 2005 and were authenticated at the Botanical Survey of India, Central National Herbarium Howrah, India (Ref No: CNH/I -1/87/2005-TECH/1326). An authentic voucher specimen was deposited at the Herbarium of the Division of Pharmacognosy, Department of Pharmaceutical Technology, Jadavpur University, Kolkata, India.

### Preparation of the polyphenolic extract (PPE)

Leaves were air dried under room temperature without exposure to sunlight and then coarsely powdered. Powdered leaves (200 g) were macerated with 70 % aqueous/ethanol (500 mL) by stirring at room temperature for 7 days. The extract was decanted, filtered under vacuum, concentrated in a rotary vacuum evaporator (Buchi, India), and then lyophilized (Lyophilization Systems India Pvt Limited, Hyderabad). The powder extract (yielded 23 % w/v) was used for the present study.

### Animals and ethical approval

Wistar rats of either sex, weighing about 200-250 g body wt were used in the present study (M/s Ghosh Enterprises, Kolkata, India). Animals were collected from breeding colony and acclimatized to the laboratory condition for two weeks. They were housed in

macrolon cages under standard laboratory conditions (light period 7.00 am to 7.00 pm,  $21 \pm 2$  °C, and relative humidity 55-70%). The animals were fed with commercial diet from Hindustan Lever Ltd (Bangalore, India) and had free access to water (ad libitum) during the experiments. Experiments were performed complied with the rulings of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) New Delhi, India (Registration No: 0367/01/C/CPCSEA) and the study was permitted by the institutional animal ethical committee (IAEC) of the Jadavpur University, Kolkata, India.

#### Carbon tetrachloride-induced hepatotoxicity

After acclimatization, the rats were divided into four groups of six rats each. Liquid paraffin was injected (i.p) to Group I (vehicle group) at a dose of 1 ml/kg body wt (i.p), for 16 days. Concentrated carbon tetrachloride in liquid paraffin (1:2) was injected (i.p) to Group-II (control group) at a dose of 1ml/kg body wt, every 72 h for 16 days (i.e 1,4,7,10,13 and 16 days). Group III (prophylactic group) was treated with the polyphenolic extract (200 mg/kg body wt) for 7 days before carbon tetrachloride injection (1 ml/kg body wt (i.p) in liquid paraffin every 72 h for 16 days) and continued for 16 days. Group IV (curative group) was simultaneously treated with the polyphenolic extract (200 mg/kg body wt) orally along with carbon tetrachloride injection (1 ml/kg body wt, (i.p) in liquid paraffin at every 72 h for 16 days) and continued for 16 days. The details of the groups and treatments are given in Table 1. All animals were sacrificed 24 h after the last dose of carbon tetrachloride injection<sup>29</sup>.

#### Tamoxifen-induced hepatotoxicity

The experiment was performed as described by El-Beshbishy with slight modifications<sup>30</sup>. Rats were randomly divided into four groups of 6 animals each. Group I served as normal untreated, which received a normal saline (5 ml/ kg body wt). Group II (tamoxifen- intoxicated rats) rats were treated with tamoxifen at a dose of 45 mg/kg/day (i.p.) for 7 successive days. Group III rats were treated orally with the polyphenolic extract (200 mg/ kg body wt daily) for 10 days before and 7 days after tamoxifen intoxication (dose identical to Group II) for 18 days as protection against (prophylactic group) liver injury induced by tamoxifen. Group IV rats were simultaneously treated with the polyphenolic extract along with tamoxifen intoxication (dose identical to Group II) for 18 days as protective against (curative treatment) liver injury induced by tamoxifen.

#### Determination of serum biochemical parameters

At the end of each experiment, the animals were fasted, anesthetized (ketamine 60 mg/kg, i.p.) and killed by cervical decapitation. Blood was obtained by direct cardiac puncture for serum biochemical estimations. Blood samples were kept at room temperature for 1 h and centrifuged at 3000 rpm for 30 min to obtain the serum, which was kept at -20° C until the assay. The levels of AST, ALT, ALP,  $\gamma$ -GT, bilirubin and the total proteins were measured spectrophotometrically (Spekol 1200, Japan) using the Span Diagnostic Kits (Mumbai, India). The abdomen was excised and their livers were removed immediately by dissection, washed in ice-cold isotonic saline and blotted between two filter papers. Their livers were wrap-

**Table 1.** Effects of the polyphenolic extract of *I. frutescens* on serum biochemical parameters in carbon tetrachloride-induced hepatotoxic rat liver.

Parameters	Liquid Paraffin (1 mL/kg)	CCl <sub>4</sub> (1 mL/kg)	PPE (P) (200 mg/kg)	PPE (C) (200 mg/kg)
AST (U/L)	137.33 $\pm$ 2.45	267.16 $\pm$ 3.39	181.50 $\pm$ 3.04**	209.83 $\pm$ 2.27**
ALT(U/L)	40.67 $\pm$ 2.29	181.0 $\pm$ 5.79	109.33 $\pm$ 2.84**	152.50 $\pm$ 3.50**
ALP(U/L)	182.16 $\pm$ 3.	361.5 $\pm$ 3.24	220.33 $\pm$ 2.48**	254.83 $\pm$ 2.68**
Total protein (g/dL)	8.30 $\pm$ 0.14	5.75 $\pm$ 0.19	6.90 $\pm$ 0.18**	6.53 $\pm$ 0.15*
Total bilirubin (mg/dL)	0.29 $\pm$ 0.04	0.74 $\pm$ 0.02	0.54 $\pm$ 0.03**	0.65 $\pm$ 0.02*
$\gamma$ -GTP(U/L)	7.77 $\pm$ 0.31	27.50 $\pm$ 0.50	9.68 $\pm$ 0.37**	6.25 $\pm$ 0.23**

PPE: Polyphenolic extract. P: Prophylactic. C: Curative. Values are expressed as mean  $\pm$  SEM, (n = 6). \*Significance difference from CCl<sub>4</sub> treated group at p < 0.05. \*\* Significance difference from CCl<sub>4</sub> treated group at p < 0.01.

ped in aluminium foil and stored at - 80 °C. A 10 % (w/v) liver homogenate was prepared in ice-cold 0.1 M potassium phosphate buffer, pH 7.5 using Teflon homogenizer (Bio-Lab, India). The fresh aliquot from their livers homogenate was used for the estimation of glutathione and lipid peroxidation.<sup>31,32</sup>

### Histopathology

For histopathological study, liver from each animal was removed after dissection and preserved in 10% formalin. Then representative blocks of liver tissues from each lobe were taken and possessed for paraffin embedding using the standard microtechnique. The paraffin sections, 6 micron thickness were cut and stained with hematoxylin and eosin (H&E) and observed under the light microscope (Zeiss Photomicroscope, India).

### Effect of phenobarbitone-induced sleep time

The antihepatotoxic activity was assessed by determining one functional parameter, i.e., barbiturate sleep time against phenobarbital-induced sleep of rats poisoned with carbon tetrachloride. The time of lost reflex induced by short acting barbiturate is significantly prolonged if any hepatic damage and this can be used as a measure for the function of the liver drug metabolizing enzymes.<sup>33-34</sup> Eighteen rats were selected and divided into three groups of six each. Group A was kept as normal and sleep time was determined after injecting phenobarbital (40 mg/kg i.p.). Group B was administered with carbon tetrachloride (1 mL/kg i.p). Group C was given orally a dose of the polyphenolic extract (200 mg/kg). After 24 h a dose of carbon tetrachloride was given to B and C groups, duration of sleep and sleep latency was determined against sodium phenobarbital (40 mg/ kg body wt.) after 2 h of carbon tetrachloride injection to rats.

### Inhibition of phase I enzymes

The enzymes were induced in rats by oral administration of phenobarbital sodium (80 mg/ kg body wt) for 5 days<sup>35</sup>. The animals were sacrificed after an overnight fasting by ether anesthesia and the liver was excised. A 10 % homogenate was used for the assay. Various concentrations of the extract (50-500 µg/ml) were added to the reaction mixture and incubated for 2 h at 37 °C. *p*-aminophenol formed during the enzyme action reacts with phenol in alkaline me-

dium to form a blue colored product, which was measured at 630 nm. The percentage inhibitions of aniline hydroxylase and aminopyrene-N-demethylase were calculated by comparing the absorbance of control to that of the drug treated samples.

### Statistical analysis

The experimental data were expressed as means  $\pm$  SEM. The significances of difference among the various treated groups and the control group were analyzed by one-way ANOVA followed by Dunnett's multiple comparison test using the Graphad Instat Software (San Diego, USA).  $p < 0.05$  was considered as statistically significant.

## RESULTS

### Effects of carbon tetrachloride-induced hepatotoxicity

Effect on carbon tetrachloride induced hepatotoxicity on rats revealed severe hepatocellular damages evident by the marked elevation in the serum of the levels of AST, ALT, ALP,  $\gamma$ -GT and bilirubin content when compared with the carbon tetrachloride treated rats (Table 1). The different biochemical parameters were found to recover to nearly normal in prophylactic treated and curative treated rats with the polyphenolic extract when compared to carbon tetrachloride. A significant increase was observed in the melondialdehyde concentration of the liver tissue after carbon tetrachloride administration. On the contrary, a marked fall was observed in the reduced glutathione level. Both treatments (prophylactic and curative) with the polyphenolic extract when given orally, prevented progression of carbon tetrachloride-induced chronic liver injury and lipid peroxidation and level of reduced glutathione was also recouped (Table 3). Analysis of variance ( $p < 0.01$ ) showed maximum recouplement in the serum biomarkers, lipid peroxidation and reduced glutathione level with the extract at a dose of 200 mg/kg in the prophylactic study.

### Effect on tamoxifen-induced hepatotoxicity

The tamoxifen-treated group manifested a severe liver damage with elevation of AST, ALT, ALP,  $\gamma$ -GT levels. On the contrary, the total protein level was decreased as compared to the tamoxifen treated rats. The higher concentration of bilirubin and the

**Table 2.** Effects of the polyphenolic extract of *I. frutescens* on serum biochemical parameters in tamoxifen-induced hepatotoxic rat liver.

Parameters	Normal Saline (5 mL/kg)	TAM (45 mg/kg)	PPE (P) (200 mg/kg)	PPE (C) (200 mg/kg)
AST (U/L)	129.66 ± 2.76	224.0 ± 2.05	181.16 ± 4.10**	203.60 ± 3.65**
ALT(U/L)	41.36 ± 1.57	189.0 ± 1.98	77.83 ± 2.61**	98.66 ± 3.21**
ALP(U/L)	151.55 ± 3.44	184.30 ± 2.51	175.86 ± 2.29**	177.65 ± 3.11**
Total protein (g/dL)	8.05 ± 0.19	4.96 ± 0.25 6	86 ± 0.26**	6.18 ± 0.18**
Total bilirubin (mg/dL)	0.32 ± 0.01	0.66 ± 0.02	0.57 ± 0.01*	0.63 ± 0.02#
γ-GTP(U/L)	8.11 ± 0.26	20.37 ± 0.44	11.33 ± 0.27**	15.83 ± 0.22**

PPE: Polyphenolic extract. TAM: Tamoxifen. P: Prophylactic. C: Curative. Values are expressed as mean ± SEM, (n = 6).# Non significant difference from the tamoxifen-treated group. \*Significantly different from the tamoxifen treated group at p < 0.05.\*\* Significantly different from the tamoxifen-treated group at p < 0.01.

**Table 3.** Effect of polyphenolic extract of *I. frutescens* on reduced glutathione and lipid peroxidation level in carbon tetrachloride-induced hepatotoxic rat livers.

Treatment	Dose	Reduced glutathione (μg/mg protein)	Lipid peroxidation (mol MDA/mg protein)
Liquid paraffin	1 mL/kg	15.49 ± 0.51	1.57 ± 0.03
CCl <sub>4</sub>	1 mL/kg	10.24 ± 0.35	1.97 ± 0.49
PPE (C)	200 mg/kg	13.34 ± 0.38**	1.61 ± 0.02**
PPE (C)	200 mg/kg	12.48 ± 0.31**	1.49 ± 0.25**

PPE: Polyphenolic extract. P: Prophylactic. C: Curative. Values are expressed as mean ± SEM, (n = 6).\*\* Significantly different from the CCl<sub>4</sub> treated group at p < 0.01.

**Table 4.** Effects of the polyphenolic extract of *I. frutescens* on reduced glutathione and lipid peroxidation level in tamoxifen-induced hepatotoxic rat livers.

Treatment	Dose	Reduced glutathione (μg/mg protein)	Lipid peroxidation (mol MDA/mg protein)
Normal saline	5 mL/kg	16.93 ± 0.34	1.61 ± 0.01
Tamoxifen	45 mL/kg	11.61 ± 0.22	3.37 ± 0.13
PPE (C)	200 mg/kg	13.54 ± 0.24**	2.44 ± 0.15**
PPE (C)	200 mg/kg	13.18 ± 0.22**	2.84 ± 0.07**

PPE: Polyphenolic extract. P: Prophylactic. C: Curative. Values are expressed as mean ± SEM, (n = 6). \*\* Significantly different from the TAM treated group at p < 0.01.


lower of total protein confirm the depth and intensity of liver necrosis. The results on tamoxifen-intoxicated rats are shown in Table 2. These higher levels were markedly suppressed by prophylactic and curative treatments of the polyphenolic extract. But significant reverse changes in all of the parameters were observed in the prophylactic treatment (200 mg/kg) when compared to the curative group. The latter group showed a tendency towards the normal. Intraperitoneal administration of tamoxifen (45 mg/kg for 7 days) caused a significant increase of lipid peroxidation. In addition, the level of reduced glutathione was drastically reduced in tamoxifen-intoxi-

cated rats. Oral treatment with the polyphenolic extract significantly (p < 0.01), prevented elevated lipid peroxidation (Table 4). Pre-administration of the polyphenolic extract significantly increased the level of reduced glutathione in the liver when compared to the post treated tamoxifen intoxicated groups.


### Histopathology

Figure 1 exhibits the normal anatomy of the rat liver treated with liquid paraffin. Figure 2 & 3 shows an abnormal anatomy with loss of hepatic lobules, degeneration of fatty cells and cloudy swelling







**Figure 1.** Section of rat liver showing no histological abnormalities (staining with haematoxylin and eosin)  $\times 10$




**Figure 2.** Liver section of carbon tetrachloride (1 mL/kg) treated rats showing the liver cells having high degree of damage characterized by ballooning degeneration and hydrophobic degeneration (haematoxylin and eosin)  $\times 10$ .



**Figure 3.** Liver section of tamoxifen citrate (45 mg/kg) treated rats showing wide spaces of sinusoids with severe degeneration of the hepatocytes (haematoxylin and eosin)  $\times 100$ .



**Figure 4.** Section of rat liver from the carbon tetrachloride intoxicated group treated with the polyphenolic extract (200 mg/kg) showing reduced necrotic area, reduced sinusoidal dilation and increased number of hepatocytes (haematoxylin and eosin)  $\times 10$ .



**Figure 5.** Section of rat liver from the tamoxifen intoxicated group treated with the polyphenolic extract (200 mg/kg) showing reduced vacuolation, reduced space formation and increased number of hepatocytes with normal nucleus (haematoxylin and eosin)  $\times 100$ .

of the liver treated with carbon tetrachloride and tamoxifen respectively; pictures were similar to that was reported earlier.<sup>36</sup> Kupffer and Sinusoidal cells demonstrated an arrest in distribution. On the other hand, no changes or necrosis in fatty cells were observed and the arrangements of the hepatocytes were almost normal in the rats treated with the polyphenolic extract in carbon tetra chloride (Figure 4) and tamoxifen (Figure 5) intoxicated rats.


#### EFFECT ON DRUG METABOLIZING ENZYMES

Administration of the polyphenolic extract (200 mg/kg) 1 h before carbon tetrachloride injection significantly shortened the phenobarbital-induced sleep

**Table 5.** Effects of the polyphenolic extract of *I. frutescens* on Phenobarbital-induced sleep time in carbon tetrachloride-induced intoxicated rats.

Treatment	Sleep Latency (min)	Duration of sleep (min)
Control	3.83 ± 0.57	84.33 ± 2.60
CCl <sub>4</sub> + Phenobarbital	3.42 ± 0.18	187.66 ± 2.49
CCl <sub>4</sub> + Phenobarbital+ PPE	2.43 ± 0.17	106.0 ± 2.47

Dose: CCl<sub>4</sub> - 1 ml/kg; Phenobarbital - 40 mg/kg; PPE - 200 mg/kg. PPE - polyphenolic extract; Values are expressed as mean ± SEM, (n = 6). \*\* Significantly different from control group at p < 0.01.



**Figure 6.** Inhibition of aniline hydroxylase and aminopyrene-N-demethylase enzyme activities by polyphenolic extract. Values were shown as mean ± SEM of triplicate experiments. Statistical significance of the treatment was done by using Student's t-test. Results were expressed as percentage inhibition of CYP enzymes. Concentration needed for 50% inhibition (IC<sub>50</sub>) of aniline hydroxylase was 479.05 µg/ml while concentration needed for aminopyrene-N-demethylase was 344.97 µg/ml.

time (Table 5). Aniline hydroxylase and aminopyrene-N-demethylase, which are phase I enzymes and are present in the CYP, becomes implicated in the activation of carcinogens to their ultimate form. The polyphenolic extract inhibits the activities of both enzymes in a concentration-dependent manner (Figure 6) and the concentration needed for 50 % inhibition was found to be 479.05 µg/mL and 344.97 µg/mL for aniline hydroxylase and aminopyrene-N-demethylase, respectively.

## DISCUSSION

Carbon tetrachloride is an extensively studied liver toxicant, and its metabolites trichloro methyl radical (CCl<sub>3</sub> •) and trichloro methyl peroxy radical (CCl<sub>3</sub>O<sub>2</sub> •) are known to be involved in the patho-

genesis of liver.<sup>37-39</sup> These free radicals may bind directly to microsomal lipids and other cellular macromolecules contributing to the breakdown of membrane structure and disrupting cell energy processes and reactions. Prophylactic administration of the polyphenolic extract at a dose of 200 mg/kg for 7 days was effective to ameliorating morphological and biochemical indices of carbon tetrachloride induced hepatotoxicity. Simultaneous treatment with the polyphenolic extract for 18 days was also effective in mitigating a few biochemical signs of carbon tetrachloride-induced liver damage. The prophylactic treatment mediated suppression of the increased AST, ALT, ALP, γ-GT activities suggesting a possible protection against liver injury, and restoration of the level of total serum protein confirming thus its hepatoprotective action. A significant increase in hepatic lipid peroxidation was observed after carbon tetrachloride exposure. Alterations in the structure of the endoplasmic reticulum and other membranes, and loss of the metabolic enzyme activation may lead to liver damage.<sup>40</sup>

The pretreatment of carbon tetrachloride-intoxicated rats with polyphenolic extract elicited the strongest effect in inducing a highly significant decline in TBARS and a highly significant increase in the level of reduced glutathione in comparison to tamoxifen-intoxicated rats. Results suggest that the polyphenolic extract reduces the oxidative stress in rats with carbon tetrachloride and tamoxifen treatment. Taken together, the current study shows that prophylactic administration of the polyphenolic extract enhances the protection against hepatic damage caused by free radicals and lipid peroxidation, and strengthens the detoxification by carbon tetrachloride and tamoxifen in rats. The ability of a hepatoprotective drug to reduce the injurious effects or to preserve the hepatic physiological mechanisms showed the index of its protective effects. Although serum enzyme levels are not a direct measure of hepatic injury, they show the functional status of the liver. Lowering of the enzyme level is a definite indi-

cation of hepatoprotective action of the polyphenolic extract.

On the other hand, liver injury caused by the administration of tamoxifen, shows an increased activities of AST, ALT, ALP,  $\gamma$ -GT. Tamoxifen citrate is an anti-estrogenic drug widely used for the treatment of breast cancer.<sup>41</sup> It was obvious that tamoxifen in toxic doses lead to oxidative liver damage as it has been elucidated to be a hepatocarcinogen in rats.<sup>42</sup> It may be more toxic to liver because it has much higher affinity to hepatic tissues than to any other tissues. The status of oxidative stress after tamoxifen administration accompanies the decreased hepatic glutathione content and the increased lipid peroxidation. One week pre-treatment with the polyphenolic extract significantly improved serum biochemical parameters. These results show that the polyphenolic extract significantly inhibited lipid peroxidation in liver induced by tamoxifen. The decrease in antioxidant defense systems in animals renders them more susceptible to the hepatotoxicity. In the tamoxifen model prophylactic treatment with the polyphenolic extract also reduced the degree of hepatocellular injury as evidenced by the improved serum biochemical parameters and malondialdehyde levels in tamoxifen-treated rat livers.

Oral administration of the polyphenolic extract shortened the phenobarbital-induced sleep time in carbon tetrachloride-treated rats, since barbiturates are metabolized exclusively in the liver<sup>34</sup>. Prolonged sleep time after exposure of rats to carbon tetrachloride in rats shows an impaired metabolism of these hypnotics possibly due to carbon tetrachloride-induced liver damage. The treatment of animals with carbon tetrachloride caused damage to the microsomal drug-metabolizing enzymes in the hepatocytes leading to substantial decrease of the hepatic drug metabolizing capacity, which reflects a prolongation of the phenobarbital-induced sleep time.

CYP is involved in several biological interactions involving hydroxylation, and oxidative deamination, demethylation, dehydrogenation.<sup>43,44</sup> While some of the biological reactions of CYP are highly essential for the removal of xenobiotics from the body, these reactions are also involved in the activation of xenobiotics to active metabolites, which can be highly injurious to the body. Hence CYP can be considered as double-edged weapon and its control is highly needed in the physiological balance of the body. Tamoxifen has been found to be metabolized by liver primarily into three metabolites, tamoxifen-N-oxide, N-desmethyl and 4-hydroxytamoxifen, formed by CYP. The N-demethylation was demonstrated to be

catalyzed by CYP3A4 in rat and human liver. The polyphenolic extract has also been found to inhibit enzymes responsible for the activation of tamoxifen, such as, aniline hydroxylase and aminopyrene-N-demethylase. The inhibition of enzymes activity in vitro could be due to the direct inactivation of the enzymes or in vivo inhibition of enzyme expression induced by phenobarbitone.

The polyphenolic extract may interfere with the CYP and ultimately hinder the formation of free radicals and metabolites (4-hydroxy tamoxifen). The ability of the polyphenolic extract of *I. frutescens* to reduce the prolongation of phenobarbital-induced sleep in carbon tetrachloride-exposed rats is further supportive for the anti-hepatotoxic potential of the polyphenolic extract. Despite being an inhibitor of phase I enzymes, the polyphenolic extract does not cause harmful bioactivation. In vivo studies have showed that the constituents of *I. frutescens* decrease the hepatotoxicity of carbon tetrachloride and tamoxifen by favourably influencing hepatic metabolism. The probable reason for this favourable effect is inhibition of phase I enzymes, which results in the rapid clearance of the potentially toxic metabolite.

Medicinal practitioners have prescribed Ayurveda and drug from herbal origin as a system of medicine in India over centuries. Treatment options for common liver diseases such as cirrhosis, fatty liver and chronic hepatitis are problematic. Physician and patients are in need of effective therapeutic agents with a low incident of side effects. In India, more than 87 plants are used in different combinations in the preparation of 33 patented herbal formulations for liver disorders. Plants traditionally used in the alleviation of liver dysfunctions might therefore provide a useful source of new hepatoprotective compounds for development as pharmaceutical entities or as simple adjuncts to existing clinical therapies.

Our results showed that polyphenolic extract of *I. frutescens* (both prophylactic and curative) could protect the experimental animals from the deleterious effects of carbon tetrachloride and tamoxifen on liver. Moreover, the use of polyphenolic extract as prophylactic is more effective than its curative against carbon tetrachloride and tamoxifen induced liver injury. The polyphenolic extract alleviates the symptoms of liver damage as evidenced by biochemical and histopathological studies. Further clinical studies will justify the use of polyphenolic extract of *I. frutescens* as an adjuvant to any kind of patients with tamoxifen and carbontetra chloride intoxication.



## ACKNOWLEDGEMENTS

The authors are grateful to All India Council of Technical Education (AICTE), Government of India, New Delhi, India, for financial support.

## Abbreviations

- **PPE:** Polyphenolic extract.
- **CPCSEA:** Committee for the Purpose of Control and Supervision of Experiments on Animals.
- **SEM:** Standard error mean.
- **ANOVA:** Analysis of variance.
- **CCl<sub>4</sub>:** Carbon tetrachloride.
- **AST:** Aspartate aminotransferase.
- **ALT:** Alanine aminotransferase.
- **TBARS:** Thiobarbituric acid reactive substances.
- **ALP:** Alkaline phosphatase.
- **γ-GT:** Gamma glutamyl transferase.
- **AICTE:** All India Council of Technical Education.
- **AICTE, bw:** body weight, **ip:** intraperitoneal, **w/v:** weight/volume, **g:** gram, **kg:** kilogram, **ml:** milliliter.
- **TAM:** Tamoxifen.
- **MDA:** Malondialdehyde.
- **h:** hour.

## REFERENCES

1. Meyer SA, Kulkarni AP. Introduction to biochemical toxicology, Hepatotoxicity. In: Hodgson E, Smart RC (eds.). 3rd. Ed. A. New York: John Wiley & Sons Inc; 2001, p. 487-90.
2. Tomboloni A, Cingolani M. Fatal accidental ingestion of carbon tetrachloride: a postmortem distribution study. *J Forensic Sci* 1996; 41(1): 166-8.
3. Tesheke CB, Vierke W, Goldermann L. Carbon tetrachloride levels and serum activities of liver enzymes following acute carbon tetrachloride intoxication. *Toxicol Lett* 1983; 17(1-2): 175-80.
4. Jordan VC. Tamoxifen: a most unlikely pioneering medicine. *Natl Rev Drug Discov* 2003; 2: 205-13.
5. Williams GM, Iatropoulos MJ, Djordjevic MV, Kaltenberg OP. The triphenylethylene drug tamoxifen is a strong liver carcinogen in the rat. *Carcinogenesis* 1993; 14: 315-17.
6. Dehal SS, Kupfer D. CYP2D6 catalyzes tamoxifen 4-hydroxylation in human liver. *Cancer Res* 1997; 57: 3402-6.
7. Fowler JS. Carbon tetrachloride metabolism in the rabbit. *Br J Pharmacol* 1969; 37: 733-7.
8. Emile FN, Dragan YP, Colin RJ, Jordan VC, Pitot, HC. Effects of tamoxifen administration on the expression of xenobiotic metabolizing enzymes in rat liver. *Cancer Res* 1995; 55: 1780-6.
9. Wong FW, Chan WY, Lee SS. Resistance to carbon tetrachloride induced hepatotoxicity in mice which lack CYP 2E1 expression. *Toxicol App Pharmacol* 1998; 153: 109-18.
10. Boll M, Weber LW, Becker E, Stampfl A. Mechanism of Carbon Tetrachloride-Induced Hepatotoxicity. Hepatocellular Damage by Reactive Carbon Tetrachloride Metabolites. *Z Naturforsch* 2001; 56: 649-59.
11. Mitra SK, Venkataranganna MV, Sundaram R, Gopumadhavan S. Protective effect of HD-03, a herbal formulation, against various hepatotoxic agents in rats. *J Ethnopharmacol* 1998; 63(3): 181-6.
12. Sies H. Strategies of antioxidant defence. *Eur J Biochem* 1993; 215(2): 213-19.
13. Letteron P, Labbe G, Degott C, Berson A, Fromenty B, Delaforge M, Larrey D, Pessayre D. Mechanism for the protective effects of silymarin against carbon tetrachloride-induced lipid peroxidation and hepatotoxicity in mice. Evidence that silymarin acts both as an inhibitor of metabolic activation and as a chain-breaking antioxidant. *Biochem Pharmacol* 1990; 39: 2027-34.
14. Rosen GM, Britigan BE, Halpern HJ, Pou S. Spin trapping free radicals in biological systems; in Free Radicals, Biology and Detection by Spin Trapping. Oxford Press New York: Oxford; 1999, p. 394.
15. Lieber CS. Role of oxidative stress and antioxidant therapy in alcoholic and nonalcoholic liver diseases. *Adv Pharmacol* 1997; 38: 601-28.
16. Bass NM. Is there any use for nontraditional or alternative therapies in patients with chronic liver disease? *Curr Gastro Rep* 1999; 1(1): 50-6.
17. Ross JA, Kasum CM. Dietary flavonoids: bioavailability, metabolic effects, and safety. *Ann Rev Nutr* 2002; 22: 19-34.
18. Breinholt V, Lauridsen ST, Dragsted LO. Differential effects of dietary flavonoids on drug metabolizing and antioxidant enzymes in female rat. *Xenobiotica* 1999; 29(12): 1227-40.
19. Hikino H, Kiso Y. Natural products for liver diseases. In: Economical and medicinal plant research. Vol. 2. London: Academic Press; 1988, p. 39-72.
20. Anonymous. Wealth of India. Raw Materials, Council of Scientific and Industrial Research (CSIR). Vol. V. New Delhi: CSIR; 1976, p. 162-3.
21. Lakshmi DKM, Venkata Rao E, Venkata Rao D. Triterpenoid constituents of *Ichnocarpus frutescens*. *Indian Drugs* 1985; 22: 552-3.
22. Singh RP, Singh RP. Flavonoids of the flowers of *Ichnocarpus frutescens*. *J Indian Chem Soc* 1987; 14: 715-56.
23. Parinitha M, Harish GU, Vivek NC, Mahesh T, Shivanna MB. Ethno-botanical wealth of Bhadra wild life sanctuary in Karnataka. *Indian J Trad Knowl* 2004; 3(1): 37-50.
24. Bhandary MJ, Chandrashekar KR, Kaveriappa KM. Medical Ethnobotany of the Siddis of Uttara Kannada District, Karnataka, India. *J Ethnopharm* 1995; 47(3): 149-58.
25. Kang SY, Sung SH, Park JH, Kim YC. Hepatoprotective activity of scopoletin, a constituent of *Solanum lyratum*. *Arch Pharm Res* 1998; 21(6): 718-22.
26. Kumarappan, CT, Chandra R, Mandal SC. Anti-inflammatory activity of *Ichnocarpus frutescens*. *Pharmacologyonline* 2006; 3: 201-16.
27. Kumarappan, CT, Mandal SC. Antitumor activity of polyphenolic extract of *Ichnocarpus frutescens*. *Exp Oncol* 2007; 29(2): 94-101.
28. Kumarappan, CT, Mandal SC. Polyphenolic extract of *Ichnocarpus frutescens* attenuates diabetic complications in streptozotocin treated diabetic rats. *Renal Failure* 2008; 30(3): 307-22.
29. Mukherjee S, Sur A, Maiti BR. Hepatoprotective effect of *Swertia chirata* on rat. *Ind J Exp Biol* 1997; 35(4): 384-8.
30. El-Beshbishy H. The effect of dimethyl dimethoxy biphenyl dicarboxylate (DDB) against tamoxifen induced liver injury

- in rats: DDB use is curative or protective. *J Biochem Mol Biol* 2005; 38(3): 70-7.
31. Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys* 1959; 82(1): 70-7.
  32. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxidation in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979; 95(2): 351-8.
  33. Brauer RW, Root MA. The effect of carbon tetrachloride induced liver injury upon the acetylcholine hydrolyzing activity of blood plasma of the rat. *J Pharmacol Exp Ther* 1946; 88(2): 109 - 118.
  34. Vogel G. New natural products and plants with drugs, pharmacological, biological or therapeutical activity. H. Wagner and P. Wolff (eds.). Heidelberg, Berlin: Springer Verlag; 1977, p. 240-65.
  35. Mazel P. Experiments illustrating drug metabolism in vitro. In: La Du, BN, Mandel HG, Way EL (eds.). Fundamentals of drug metabolism and drug disposition. Baltimore: Williams and Wilkins; 1971, p. 546-50.
  36. Agarwal AK, Mehendale HM. Chlordecone potentiation of carbon tetrachloride hepatotoxicity in ovariectomized rats. *Toxicol* 1984; 29(4): 315-23.
  37. Pencil SD, Brattin WJ, Glende EA, Recknagel RI. Carbon tetra chloride dependent inhibition of lipid secretion by isolated hepatocytes characterization and requirement for bioactivation. *Biochem Pharmacol* 1984; 33(15): 2419-23.
  38. Recknagel RO, Glonde EA, Dolak JA, Walter RL. Mechanism of carbon tetrachloride toxicity. *Pharmacol Therap* 1989; 43(1): 139-54.
  39. Farber JL, Gerson RJ. Mechanisms of cell injury with hepatotoxic chemicals. *Pharmacol Rev* 1984; 36(2): 71S-75S.
  40. Azri S, Mata HP, Reid LL, Gandolfi AJ, Brendel K. Further examination of the selective toxicity of CCl<sub>4</sub> in rat liver slices. *Toxicol Appl Pharmacol* 1992; 112(1): 81-6.
  41. Desai P, Nallani S, Sane R, Moore L, Goodwin B, Buckley D, Buckley A. Induction of cytochrome P450 3A4 in primary human hepatocytes and activation of the human pregnane X receptor by tamoxifen and 4- Hydroxytamoxifen. *Drug Metab Disp* 2002; 30(5): 608-12.
  42. Athoupa M, Hirsimaki P, Parssinen R, Mantyla E. Alterations of drug metabolizing and antioxidant enzyme activities during tamoxifen induced hepatocarcinogenesis in rats. *Carcinogenesis* 1994; 15(5): 863-8.
  43. Aguiar M, Masse R, Gibbs BF. Regulation of cytochrome P450 by posttranslational modification. *Drug Metab Rev* 2005; 37: 379-404.
  44. Cooper DY. Discovery of the function of the heme protein P-450: a systematic approach to scientific research. *Life Sci* 1973; 13: 1151-61.