

Fatty liver and abdominal fat relationships with high C-reactive protein in adults without coronary heart disease

Guillermo C. Cardoso-Saldaña,* Aida X. Medina-Urrutia,* Carlos Posadas-Romero,*
Juan G. Juárez-Rojas,* Esteban Jorge-Galarza,* Gilberto Vargas-Alarcón,** Rosalinda Posadas-Sánchez*

* Departamento de Endocrinología. ** Departamento de Biología Molecular.
Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico.

ABSTRACT

Background and rationale. Fatty liver (FL) and abdominal visceral fat (AVF) are strongly associated with systemic inflammation, however, it has not been defined if each one is independently involved, and if the insulin resistance is associated. To investigate if FL, AVF and insulin resistance are independently or additively associated with the high-sensitivity C-reactive protein (hs-CRP) in subjects without coronary artery disease we included 491 men and 553 women. **Material and methods.** All had anthropometric and plasma biochemical measurements, FL and AVF assessments by computed tomography. **Results.** The FL prevalence was 35.6% in men and 28.0% in women, $p < 0.01$. The prevalence of obesity, metabolic syndrome and homeostasis model assessment of insulin resistance (HOMA-IR) was significantly higher in FL compared to non FL subjects. FL and AVF accounted for 21 and 17%, respectively, to hs-CRP plasma levels. FL, AVF $\geq P_{75}$ and HOMA-IR $\geq P_{75}$ were independently and additively associated with plasma hs-CRP. The risk of hs-CRP ≥ 3 mg/L increased progressively in men from 1.36 (0.5-3.86) through 3.58 (1.32-9.7) in those with 1 or 3 factors respectively. In women from 2.25 (1.2-4.2) to 4.67 (2.3-9.4), respectively. In conclusion, both the FL and hs-CRP ≥ 3 mg/L occur in 1 of every 3 non CAD subjects. In men, FL and AVF $\geq P_{75}$ were associated with 3.6 times the risk of hs-CRP ≥ 3 mg/L, while in women, these factors were independently and additively associated with a 4.7 times higher risk of systemic inflammation.

Key words. Visceral adiposity. Hepatic fat. Insulin resistance. Systemic inflammation.

INTRODUCTION

Fatty liver (FL) disease is the most common hepatic condition in adult population. Depending on the ethnic group, it affects from 24 to 42% of the overall population and from 70 to 90% of obese subjects.¹ It is characterized by the hepatic triglyceride (TG) infiltration in individuals with less than 20 g of alcohol consumption per day and without other causes of liver disease.² Several studies have shown that FL is associated with cardiometabolic risk factors.³⁻⁵

A recent study showed that FL identified by computed tomography was associated independ-

ently of visceral fat, with diabetes, hypertension, impaired fasting glucose, metabolic syndrome, high TG levels, as well as low concentrations of serum high-density lipoprotein cholesterol (HDL-C) and adiponectin.⁶ In another study that used the hyperinsulinemic-euglycemic clamp, increased insulin resistance in liver, adipose tissue and muscles was observed in obese subjects with FL compared to obese subjects without fatty liver.⁷ In addition, recent studies have shown association of FL with coronary artery disease (CAD) even after adjusting for traditional risk factors.^{8,9} Altogether, these studies suggest that increased intrahepatic fat contributes to the pathogenesis of metabolic abnormalities and CAD. Nevertheless, it is unclear how the FL is associated with the development of CAD. One possible mechanism is the chronic inflammation state manifested by increased serum markers of systemic inflammation, among which C-reactive protein is one of the most widely studied for having a clear association with atherosclerosis.¹⁰

Correspondence and reprint request: Guillermo C. Cardoso-Saldaña, Ph.D.
Juan Badiano No. 1, Col. Sección XVI. C.P. 14080, Mexico City, Mexico.
Tel.: (5552) 5573-2911, ext.: 1272. Fax: (5552) 5573-4687
E-mail: gccardosos@yahoo.com

*Manuscript received: October 14, 2014.
Manuscript accepted: December 20, 2014.*

Several reports have shown the association of FL with C-reactive protein,¹¹⁻¹⁴ however, the possible involvement of abdominal visceral fat (AVF) and insulin resistance, two pro-inflammatory factors closely related to FL, has not been considered in those studies. The main objective of this study was to investigate whether fatty liver is associated with the C-reactive protein concentration, independently of AVF, and insulin resistance, in men and women. Furthermore, we investigated whether the coexistence of increased AVF and insulin resistance increases that association.

MATERIAL AND METHODS

The Genetics of Atherosclerotic Disease Study (GEA) was designed at the National Institute of Cardiology Ignacio Chávez, Mexico, to examine the genetic basis of CAD and to investigate the relationship between traditional and emerging risk factors with atherosclerosis in the adult Mexican mestizo population. The GEA study included 1,500 patients with premature CAD and a control group of 1,600 individuals with no personal or family history of premature CAD. The present study included 1,044 individuals of the control group, aged 30 to 75 years, without diabetes mellitus, alcohol consumption less than 20 g/day, no acute or chronic inflammatory processes, and no history or clinical evidence of renal failure or liver disease.² Participants were recruited from blood bank-donors, and through brochures posted in social service centers. The GEA study was approved by the Ethics committee of the National Institute of Cardiology Ignacio Chávez and conducted according to the Declaration of Helsinki. All participants signed an informed consent form.

Subjects

Questionnaires were applied to participants to obtain information regarding demographics, personal history of cardiovascular risk factors, physical activity, alcohol consumption, and medications. Weight was measured in kilograms (kg) and height was measured in centimeters (cm), using a calibrated scale and a wall-mounted height rod. Body mass index (BMI) was calculated using the weight (kg)/height (m²) formula. Waist circumference was measured with a fiberglass tape, at a midpoint between the lower margin of the last rib and the iliac crest with a participant breathing out gently. After at least 5 min of rest, systolic and diastolic blood pres-

ures were measured three times in a sitting position. The average of the last two consecutive measurements was used for the analysis.

Samples

After 10 h of fasting and 20 minutes in a sitting position, venous blood samples were collected. Glucose, total cholesterol, TG and HDL-C concentrations were determined in plasma with enzymatic methods (Roche Diagnostics GmbH, Mannheim, Germany) using a Hitachi 902 auto analyzer (Hitachi LTD, Tokyo, Japan).¹⁵ Low-density lipoprotein cholesterol (LDL-C) was calculated based on the Friedewald's equation.¹⁶ The reproducibility and precision of lipids and lipoproteins measurements were periodically assessed by the Lipid Standardization Program of the Center for Disease Control and Prevention (LSP-CDC, Atlanta, GA, USA). The intra and inter-assay coefficients of variation were less than 3%. The serum insulin concentration was determined by radioimmunoassay (Human insulin RIA kit; Millipore, Cat. HI-14K St. Charles, Missouri, USA). The intra and inter-assay coefficients of variation were 2.1% and 6.8%, respectively. Insulin resistance was estimated using the homeostasis model assessment of insulin resistance (HOMA-IR) (Insulin IU/mL X Glucose mmol/22.5).¹⁷ High-sensitivity C-reactive protein (hs-CRP) concentration was determined by immunonephelometry (Cardiophase hsCRP, SIEMENS Healthcare Diagnostics Products GmbH, Marburg, Germany) on a BN ProSpec nephelometer according to the manufacturer's procedures. Inter and intra-assay coefficients of variation were less than 6%.

Tomography

Computed axial tomography (CAT) is a validated method to quantify AVF and abdominal subcutaneous fat (ASF) and to identify the presence of steatosis.¹⁸⁻²⁰ In this study, these measurements were performed using a 64-slice CAT scanner (Somatom Sensation, 64, Forchheim, Germany). Liver attenuation in CAT was calculated as the average of five measurements made in regions of interest of 1.0 cm² in both lobes. Spleen attenuation measured in several regions of interest was used as an internal control for the standardization of hepatic attenuation. The presence of hepatic steatosis was defined as the liver/spleen attenuation ratio less than 1.0 Hounsfield units.²¹ To measure abdominal fat, a single topographic slice at the level of L4-L5 inter-vertebral

disc space was performed. AVF and ASF areas were separated by manual tracing along the abdominal muscle wall. Total abdominal fat and AVF were quantified in cm^3 and the ASF was calculated by subtracting the AVF area from total abdominal fat.^{15,16}

Hypertension was defined as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg and/or use of antihypertensive medications. Overweight was defined as a BMI from 25.0 to 29.9 kg/m^2 and obesity as a BMI ≥ 30 kg/m^2 . The presence of abdominal obesity was considered when waist circumference values were ≥ 80 cm in women and ≥ 90 cm in men.²⁵ Fasting blood glucose from 100 to 125 mg/dL was considered impaired fasting glucose, whereas values ≥ 126 mg/dL or treatment with hypoglycemic agents were used to determine the presence of diabetes mellitus. Metabolic syndrome was defined based on the presence of 3 or more of the following factors:²³

- Central obesity.
- Triglycerides ≥ 150 mg/dL.
- HDL cholesterol < 40 mg/dL in men and < 50 mg/dL in women.
- Fasting glucose ≥ 100 mg/dL.
- Systolic blood pressure ≥ 130 mmHg and/or diastolic blood pressure ≥ 85 mmHg.

According to its association with increased cardiovascular risk, hs-CRP was considered high when it was ≥ 3 mg/dL.²⁴ Physical activity was measured using the Baecke questionnaire.²⁵ Total activity was obtained from the sum of the work exercise and leisure time activities. This questionnaire, has been validated in adult populations and provides reliable information.

In order to set the cut-off points to other coronary risk factors in our study population, a group of individuals with BMI < 30 , no hypertension, hypertriglyceridemia, hypoalphalipoproteinemia or diabetes mellitus was selected. The 75th percentile (P_{75}) for risk factors in the sub-sample was as follows: insulin = 15.20 $\mu\text{U}/\text{mL}$ in men, and 16.97 $\mu\text{U}/\text{mL}$ in women; HOMA-IR = 3.38 in men, and 3.66 in women; ASF = 221.7 cm^3 in men and 335.5 cm^3 in women; AVF = 151.5 cm^3 in men and 122.0 cm^3 in women.

Statistical analysis

Continuous variables with normal distribution were expressed as mean \pm SD, and categorical

variables were expressed as percentage. Variables with normal distribution were compared using the Student's *t*-test and variables with asymmetrical distribution with the Mann-Whitney U test. Chi-square test was applied to compare the prevalence of coronary risk factors between the groups. The independence of the associations was assessed with a multivariate linear or logistic regression analysis [standardized β coefficient or odds ratio (OR)] and a 95% confidence interval (95% CI). Two-tailed *p* values < 0.05 were considered statistically significant. All analyses were performed using the statistical package SPSS version 15.0 (SPSS Chicago, II).

RESULTS

The prevalence of FL in the study population was 31.6% (35.6% in men and 28.0% in women, $p < 0.01$). In subjects with FL compared to participants without FL, the prevalence of abdominal obesity, defined by waist circumference (93 *vs.* 70%, $p < 0.001$ in men and 94 *vs.* 80%, $p < 0.001$ in women) or AVF $\geq P_{75}$ (86 *vs.* 59%, $p < 0.001$ in men and 81 *vs.* 43%, $p < 0.001$ in women), was markedly increased. The high prevalence of abdominal obesity was associated with high rates of metabolic syndrome (55.3 *vs.* 38.1%, $p < 0.001$ in men and 48.4 *vs.* 23.2%, $p < 0.001$ in women) and insulin resistance (86.8 *vs.* 53.1%, $p < 0.001$ in men and 86.7 *vs.* 45.2%, $p < 0.001$ in women), in participants with and without FL, although significantly higher in subjects of both genders with intra-hepatic fat excess. Compared to subjects with non FL, all adiposity measurements were significantly higher in the subjects with FL (Table 1).

Overall, FL was associated with a more adverse metabolic profile (Table 1), characterized in both genders by higher blood pressure, triglycerides, insulin, HOMA-IR and hs-CRP values, as well as by significantly higher prevalence of impaired fasting glucose, increased triglycerides and low HDL-C. No significant differences were observed in age or the prevalence of increased LDL-C, hypertension, and the percentage of subjects under pharmacological treatment for these conditions in both groups. Figure 1 shows that compared to subjects without risk factors, subjects with FL, AVF $\geq P_{75}$ or HOMA-IR $\geq P_{75}$, particularly those of female gender, had a higher prevalence of hs-CRP ≥ 3 mg/L. The independence of the associations of FL, AVF $\geq P_{75}$ and HOMA-IR $\geq P_{75}$ with the natural logarithm of hs-CRP (Ln hs-CRP) as a continuous variable, was determined by a multivariate linear regression analysis in each gen-

Table 1. Clinical and biochemical characteristics of study subjects by gender and fatty liver.

	Men			Women		
	FL (-)	FL (+)	p*	FL (-)	FL (+)	p*
n (%)	316 (64.4)	175 (35.6)	-	398 (72.0)	155 (28.0)	-
Age, years	53.5 ± 10.3	51.9 ± 8.8	ns	53.21 ± 9.2	52.3 ± 8.5	ns
Waist, cm	95.5 ± 9.8	102.8 ± 9.4	< 0.001	88.5 ± 10.6	96.4 ± 11.5	< 0.001
BMI, kg/m ²	27.2 ± 3.5	30.0 ± 3.7	< 0.001	27.4 ± 4.4	30.6 ± 4.6	< 0.001
Adipose subcutaneous fat, cm ³	226.1 ± 87.7	296.3 ± 103.7	< 0.001	316.9 ± 105.8	359.4 ± 106.3	< 0.001
Adipose visceral fat, cm ³	161.9 ± 65.8	199.3 ± 59.0	< 0.001	121.8 ± 51.7	163.4 ± 51.9	< 0.001
Systolic blood pressure, mmHg	120.1 ± 16.8	122.2 ± 15.5	ns	112.0 ± 17.8	115.8 ± 15.1	0.023
Diastolic blood pressure, mmHg	74.3 ± 9.7	76.3 ± 9.5	0.03	69.1 ± 8.8	71.4 ± 8.9	0.006
Triglycerides, mg/dL	154.5 (113.0-210.8)	171.8 (126.2-236.3)	0.013	125.0 (98.0-166.2)	161.5 (126.4-211.5)	< 0.001
Insulin, UI/mL	14.6 (10.6-20.2)	22.5 (17.5-28.5)	< 0.001	14.9 (11.4-19.9)	23.1 (17.9-31.9)	< 0.001
HOMA-IR	3.2 (2.3-4.5)	5.5 (3.9-6.9)	< 0.001	3.2 (2.3-4.4)	5.3 (3.8-7.5)	< 0.001
hs-CRP, mg/L	1.15 (0.63-1.97)	1.65 (0.89-3.21)	< 0.001	1.42 (0.76-3.0)	2.93 (1.5-4.5)	< 0.001
Glucose ≥ 100, mg/dL, n (%)	39 (12.3)	50 (28.6)	< 0.001	28 (7.0)	40 (25.8)	< 0.001
TG > 150 mg/dL, n (%)	162 (21.3)	111 (63.4)	0.006	132 (33.2)	86 (55.5)	< 0.001
Hypoalphalipoproteinemia, n (%)	148 (46.8)	101 (57.7)	0.013	173 (43.5)	108 (69.7)	< 0.001
LDL-C > 130 mg/dL, n (%)	113 (32.8)	63 (36.0)	ns	125 (31.4)	53 (34.2)	ns
Hypertension, n (%)	69 (21.8)	48 (27.4)	ns	75 (18.8)	33 (21.3)	ns
Lipid therapy, n (%)	19 (6.0)	13 (7.4)	ns	23 (5.8)	12 (7.7)	ns
Antihypertensive therapy, n (%)	37 (11.7)	30 (17.1)	ns	61 (15.4)	25 (16.1)	ns

Values are expressed as mean ± SD, median (interquartile range), or n(%). FL: fatty liver. BMI: body mass index. HOMA-IR: homeostasis model assessment of insulin resistance. hs-CRP: high-sensitive C-reactive protein. TG: triglycerides. Hypoalphalipoproteinemia: < 40 mg/dL in men and < 50 mg/dL in women. LDL-C: low density lipoprotein cholesterol; hypertension: systolic ≥ 140 or diastolic ≥ 90. p* Student's t-test or Mann-Whitney U test.

Table 2. Association between fatty liver, adipose visceral fat ≥ P₇₅ and HOMA-IR ≥ P₇₅ with the natural logarithm of hs-CRP in multivariate linear regression analyses.

	Fatty liver β coefficients (95%CI)	AVF ≥ P ₇₅ β coefficients (95%CI)	HOMA-IR ≥ P ₇₅ β coefficients (95%CI)
Men			
Model 1	0.21 (0.12 - 0.30)	0.20 (0.12 - 0.30)	0.16 (0.07 - 0.25)
Model 2	0.19 (0.10 - 0.30)	0.16 (0.06 - 0.30)	0.09 (-0.01 - 0.20)
Model 3	0.17 (0.06 - 0.30)	0.12 (0.02 - 0.20)	0.01 (-0.09 - 0.13)
Women			
Model 1	0.27 (0.20 - 0.40)	0.31 (0.25 - 0.43)	0.26 (0.19 - 0.30)
Model 2	0.19 (0.10 - 0.30)	0.25 (0.17 - 0.38)	0.18 (0.08 - 0.30)
Model 3	0.13 (0.04 - 0.30)	0.19 (0.10 - 0.32)	0.08 (-0.02 - 0.21)

AVF: adipose visceral fat. HOMA-IR: homeostasis model assessment of insulin resistance. Model 1: unadjusted. Model 2: adjusted by age, hypertension, glucose. HDL-C: high density lipoprotein cholesterol, triglycerides, total kilocalories and physical activity. Model 3: model 2 plus AVF and HOMA-IR.

der using unadjusted and adjusted models (Table 2). All three factors were related to increased hs-CRP concentrations in both genders for model 1. After adjustment for traditional risk factors, model 2, the association of hs-CRP with FL and $AVF \geq P_{75}$ remained significant whereas the significance of the HOMA-IR association was lost in men. In model 3, the analysis of the association of ln hs-CRP with FL was further adjusted for $AVF \geq P_{75}$ and $HOMA-IR \geq P_{75}$, while the association of the protein with $AVF \geq P_{75}$ was adjusted for FL and $HOMA-IR \geq P_{75}$. In this full adjusted model, the presence of FL was independently and significantly associated with an increase in ln hs-CRP concentrations in men (β [95% CI]: 0.17 [0.06-0.30]; $p < 0.001$) which correspond to a 21% higher average hs-CRP level, and in women (β [95% CI]: 0.13 [0.04-0.30]; $p < 0.01$) which correspond to a 17% higher average hs-CRP level. The $AVF \geq P_{75}$ association with ln hs-CRP in both genders was also significant and independent; how-

ever, ln hs-CRP was more strongly associated in women than in men ($p < 0.01$) (Table 2).

Other analyses were conducted to individually and jointly examine the association of FL, $AVF \geq P_{75}$ and $HOMA-IR \geq P_{75}$ with hs-CRP as a categorical variable ($hs-CRP \geq 3$ mg/L). Figure 2 shows that increasing the number of cardiometabolic risk factors from zero to three raises the prevalence of elevated hs-CRP from 10.8 to 27.7% ($p < 0.01$) in men, and from 15.4 to 52.2% in women ($p < 0.001$). In addition the association of FL, AVF and HOMA-IR with $hs-CRP \geq 3$ mg/L in men, was statistically significant after adjust for traditional risk factors, only when combining all three cardiometabolic factors (OR [95% CI]:3.58 [1.32-9.7], $p < 0.01$) (Table 3). In women, the presence of a single factor was enough to raise the prevalence of $hs-CRP \geq 3$ mg/L, which showed gradual increments (OR [95% CI]: 2.25 [1.20-4.2]) to 4.67 (2.3-9.4) as the number of risk factors increased.




Figure 1. Prevalence of $hs-CRP \geq 3$ mg/L in subjects without (-) or with (+) fatty liver (FL), adipose visceral fat ($AVF \geq P_{75}$ (AVF : men = 151.5 cm³, women = 122 cm³) or $HOMA-IR \geq P_{75}$ (men = 3.38, women = 3.66).




Figure 2. Prevalence of $hs-CRP \geq 3$ mg/L associated with the presence of fatty liver, adipose visceral fat $\geq P_{75}$ ($AVF \geq P_{75}$: men = 151.5 cm³, women = 122 cm³) or $HOMA-IR \geq P_{75}$ (men = 3.38, women = 3.66) and combination of two or three cardiometabolic risk factors. Trend in men $p = 0.005$, in women $p < 0.001$.

Table 3. Association of fatty liver, adipose visceral fat $\geq P_{75}$ and $HOMA-IR \geq P_{75}$, with high $hs-CRP \geq 3$ mg/L in multivariate logistic regression analyses.

Risk factors (n)*	Men		Women	
	Unadjusted, OR (95% CI)	Ajusted, [†] OR(95% CI)	Unadjusted, OR(95% IC)	Ajusted, [†] OR(95% IC)
0**	Reference	Reference	Reference	Reference
1	1.26 (0.5 - 3.0)	1.36 (0.5 - 3.86)	2.25 (1.27 - 3.96)	2.25 (1.20 - 4.2)
2	2.04 (0.92 - 4.5)	2.03 (0.75 - 5.5)	3.29 (1.93 - 5.61)	2.61 (1.4 - 4.9)
3	3.16 (1.44 - 6.43)	3.58 (1.32 - 9.7)	6.20 (3.54 - 10.9)	4.67 (2.3 - 9.4)

* Factors: fatty liver, abdominal visceral fat $\geq P_{75}$ and $HOMA-IR \geq P_{75}$. ** Without fatty liver, abdominal visceral fat $\geq P_{75}$, $HOMA-IR \geq P_{75}$. [†] Adjusted by age, hypertension, glucose, HDL-C: high density lipoprotein cholesterol, triglycerides, total kilocalories and physical activity.

DISCUSSION

This study analyzes for the first time the influence of AVF, FL and HOMA-IR on hs-CRP in Mexican mestizo population. Both adipose stores and HOMA-IR had an independent and additive effect on the inflammation degree. FL and increased AVF were associated with hs-CRP concentration in both men and women independently of traditional risk factors and insulin resistance. The effect of FL was more evident in men while AVF was in women. These results suggest that, independent of insulin resistance, both adipose stores can predict an increased systemic inflammation. The combination of FL, AVF $\geq P_{75}$ and HOMA-IR $\geq P_{75}$ was associated with a gradual increase in the risk of hs-CRP ≥ 3 mg/L, which was 3.58 and 4.67 times higher in men and women in which all three factors were present, respectively.

It has been previously reported that compared to obese subjects with no liver impairment, obese subjects with steatosis or steatohepatitis had increased hs-CRP and other markers of inflammation.⁵ Also, serum C reactive protein was found associated with liver steatosis and visceral fat accumulation in type 2 diabetes mellitus patients but not in non-diabetic subjects.²⁶ Another study in 2,388 Brazilian subjects, showed that hepatic steatosis, obesity, central obesity (with a crude estimation of waist circumference) and metabolic syndrome are independently and additionally associated with the likelihood of having hs-CRP ≥ 3 mg/L.¹⁰ In a study of Mexican subjects in which AVF was not determined, those with FL also showed a significant increase in ultrasensitivity CRP, after adjustment for traditional cardiovascular risk factors.¹⁴ Our data confirm and extend the findings from those studies by demonstrating that FL is associated with hs-CRP even after controlling for visceral abdominal adiposity and insulin resistance. Although increases visceral abdominal adiposity and insulin resistance are closely related to hepatic steatosis,^{27,28} the observation that AVF and HOMA-IR are associated, independently from one another and together with systemic inflammation, and that they do not affect the association of FL with hs-CRP, could be interpreted as evidence that insulin resistance and AVF contribute to the risk of systemic inflammation by different but closely related pathophysiological mechanisms. Indeed previous evidence has shown that insulin-resistant adipose tissue plays a key role in the development of the metabolic and histological abnormalities present in obese subjects

with FL.^{29,30} In physiological conditions, the adipose tissue is the main source of free fatty acids for the liver synthesis of triglycerides. However, when the adipose tissue is insulin-resistant, releases excessive amounts of fatty acids that accumulate in the liver along with other compounds such as diacylglycerols and ceramides, which trigger proinflammatory cytokine production.^{31,32} Oxidative stress, produced by the lipotoxic effect related to the accumulation of saturated fatty acids in the liver, is another inductive factor of inflammation.^{33,34} Together, these abnormalities could explain the direct and independent link of FL with hs-CRP, and HOMA-IR observed in our study.

Women of Latin American descent have one of the highest average hs-CRP concentrations.³⁵⁻³⁷ Epidemiologic and clinical trials have consistently shown that compared to men, women have higher hs-CRP concentrations, which are unrelated to age, BMI, ethnicity and hormone replacement therapy. The present study confirms that plasma hs-CRP concentration is higher in women than in men.^{38,39} The reason for this gender-based difference is unknown but some plausible explanations have been suggested:

- Because hormone replacement therapy increases CRP concentrations,⁴⁰ it is possible that endogenous estrogen stimulates CRP synthesis.
- The larger total body fat reservoir in women could raise the synthesis of proinflammatory cytokines such as IL-6, that leads to increased production of CRP by the liver.⁴¹

The higher levels of subcutaneous abdominal fat and hs-CRP we observed in women support the latter hypothesis.

Study strengths

In most previous trials, FL and CRP association was studied jointly in men and women and the effect of gender has been controlled with statistical models. In the present study, the results from a population consisting of 47.4% male and 52.6% female subjects without personal or family history of coronary artery disease or diabetes mellitus, selected from all socioeconomic strata, we used analyses adjusted for potential confounders to demonstrate independent associations of cardiometabolic factors with hs-CRP by gender. The majority of reports have assessed abdominal obesity by the crude measure of waist circumference, while in this study we used CAT scans, the best

method for differential quantification of the subcutaneous and visceral central adiposity stores.^{18,19}

Study limitations

Due to the cross sectional nature of the study, it is not possible to establish causal or temporal relationships between cardiometabolic risk factors and increased hs-CRP. Nevertheless, the findings suggest that FL is not only a cardiovascular risk marker, but also a factor with atherogenic effect mediated by systemic inflammation. It has been suggested that at least two hs-CRP concentration measurements in metabolically stable subjects are needed to appropriately determine inflammatory status.⁴² In this study hs-CRP was determined only once; however a great care was taken to avoid sampling of subjects with acute or chronic inflammatory conditions. The diagnosis of FL was established by CAT scans, but was not confirmed by liver biopsies. However, a significant correlation has been demonstrated between the liver attenuation images on CAT and the histological grade of steatosis.⁴³ Although insulin resistance was not assessed by more sophisticated approaches, the HOMA-IR index has proven to be a reliable measure of insulin sensitivity in non-diabetic subjects.¹⁷

In conclusion, both steatosis and hs-CRP ≥ 3 mg/dL are present in 1 of 3 non-diabetic subjects free of clinical cardiovascular disease. FL, AVF $\geq P_{75}$ and HOMA-IR $\geq P_{75}$ are independently and additively associated in both genders with approximately twice the risk of increased hs-CRP values. The gender-based differences in the association of FL and the cardiometabolic risk factors with the systemic inflammatory status, suggest that hs-CRP related cardiovascular risk is greater in women than men.

ABBREVIATIONS

- **AVF:** abdominal visceral fat.
- **BMI:** body mass index.
- **CAD:** coronary artery disease.
- **CAT:** computed axial tomography.
- **FL:** fatty liver.
- **GEA:** Genetics of Atherosclerotic Disease Study.
- **HDL-C:** high density lipoprotein cholesterol.
- **HOMA-IR:** homeostasis model assessment of insulin resistance.
- **hs-CRP:** high sensitivity C reactive protein.
- **LDL-C:** low density lipoprotein cholesterol.
- **SAF:** subcutaneous abdominal fat.
- **TG:** triglycerides.

SUPPORTING

This study was supported by CONACYT grant No. SALUD 2010-2 150537.

ACKNOWLEDGEMENTS

We acknowledge to healthy volunteers and patients from the Instituto Nacional de Cardiología Ignacio Chávez, as well as the staff of the Endocrinology Department.

REFERENCES

1. Thakur ML, Sharma S, Kumar A, Bhatt SP, Luthra K, Guleria R, Pandey RM, et al. Nonalcoholic fatty liver disease is associated with subclinical atherosclerosis independent of obesity and metabolic syndrome in Asian Indians. *Atherosclerosis* 2012; 223: 507-11.
2. Brunt EM. Nonalcoholic steatohepatitis: definition and pathology. *Semin Liver Dis* 2001; 21: 3-16.
3. Bhatia LS, Curzen NP, Calder PC, Byrne CD. Non-alcoholic fatty liver disease: a new and important cardiovascular risk factor? *Eur Hearth J* 2012; 33: 1190-200.
4. Ampuero J and Romero-Gómez M. Influencia de la enfermedad por hígado graso no alcohólico en la enfermedad cardiovascular. *Gastroenterol Hepatol* 2012; 35: 585-93.
5. Assy N, Djibre A, Farah R, Grosovski M and Marmor A. Presence of coronary plaques in patients with nonalcoholic fatty liver disease. *Radiology* 2010; 254: 393-400.
6. Speliotes EK, Massaro JM, Udo Hoffmann U, Vasan RS, Meigs JB, Sahani DV, Hirschhorn JN, et al. Fatty Liver is Associated With Dyslipidemia and Dysglycemia Independent of Visceral Fat: The Framingham Heart Study. *Hepatology* 2010; 51: 1979-87.
7. Fabbrini E, Magkos F, Mohammed BS, Pietka T, Abumrad NA, Patterson BW, Okunade A, et al. Intrahepatic fat, not visceral fat, is linked with metabolic complications of obesity. *Proc Natl Acad Sci USA* 2009; 106: 15430-5.
8. Liu J, Musani SK, Bidulescu A, Carr JJ, Wilson JG, Taylor HA, Fox CS. Fatty liver, abdominal adipose tissue and atherosclerotic calcification in African Americans: the Jackson Heart Study. *Atherosclerosis* 2012; 224: 521-5.
9. Sung KC, Wild SH, Kwag HJ, Byrne CD. Fatty liver, insulin resistance, and features of metabolic syndrome: relationship with coronary artery calcium in 10,153 people. *Diabetes Care* 2012; 35: 2359-64.
10. Lavie CJ, Milani RV, Verma A, O'Keefe JH. C-reactive protein and cardiovascular diseases—is it ready for prime-time? *Am J Med Sci* 2009; 338: 486-92.
11. Haukeland JW, Damas JK, Konopski Z, Løberg EM, Haaland T, Goverud I, Torjesen PA, et al. Systemic inflammation in nonalcoholic fatty liver disease is characterized by elevated levels of CCL2. *J Hepatol* 2006; 44: 1167-74.
12. Ndumele CE, Nasir K, Conceição RD, Carvalho JA, Blumenthal RS, Santos RD. Hepatic steatosis, obesity, and the metabolic syndrome are independently and additively associated with increased systemic inflammation. *Atheroscler Thromb Vasc Biol* 2011; 31: 1927-32.
13. Targher G, Bertolini L, Rodella S, Lippi G, Franchini M, Zoppi G, Muggeoso M, et al. NASH Predicts plasma inflammatory biomarkers independently of visceral fat in men. *Obesity* 2008; 16: 1394-9.

14. Lizardi-Cervera J. Association among C-reactive protein, fatty liver disease, and cardiovascular risk. *Dig Dis Sci* 2007; 52: 2375-9.
15. Sugiuchi H, Uji Y, Okabe H, Irie T, Uekama K, Kayahara N, Miyauchi K. Direct measurement of high-density lipoprotein cholesterol in serum with polyethylene glycol-modified enzymes and sulfated alpha-cyclodextrin. *Clin Chem* 1995; 41: 717-23.
16. DeLong DM, DeLong ER, Wood PD, Lippel K, Rifkind BM. A comparison of methods for the estimation of plasma low- and very low-density lipoprotein cholesterol. The Lipid Research Clinics Prevalence Study. *JAMA* 1986; 256: 2372-7.
17. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985; 28: 412-9.
18. Sjöström L, Kvist H, Cederblad A, Tylén U. Determination of total adipose tissue and body fat in women by computed tomography, 40K, and tritium. *Am J Physiol* 1986; 250: E736-E745.
19. Kvist H, Chowdhury B, Grangard U, Tylén U, Sjöström L. Total and visceral adipose-tissue volumes derived from measurements with computed tomography in adult men and women: predictive equations. *Am J Clin Nutr* 1988; 48: 1351-61.
20. Ricci C, Longo R, Gioulis E, Bosco M, Pollesello P, Massutti F, Croce LS, et al. Non invasive in vivo quantitative assessment of fat content in human liver. *J Hepatol* 1997; 27: 108-13.
21. Piekarski J, Goldberg HI, Royal SA, Axel L, Moss AA. Difference between liver and spleen CT numbers in the normal adult: its usefulness in predicting the presence of diffuse liver disease. *Radiology* 1980; 137: 727-9.
22. Sánchez-Castillo CP, Velázquez-Monroy O, Berber A, Lara-Esqueda A., Tapia-Conyer R, James WP. Encuesta Nacional de Salud (ENSA) 2000 Working Group. Anthropometric cut-off points for predicting chronic diseases in the Mexican National Health Survey 2000. *Obes Res* 2003; 11: 442-51.
23. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, Fruchart JC, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009; 120: 1640-5.
24. Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO 3rd, Criqui M, Fadl YY, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: a statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 2003; 107: 499-511.
25. Baecke JA, Borema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982; 36: 936-42.
26. Iwasaki T, Nakajima A, Yoneda M, Terauchi Y. Relationship between the serum concentrations of C-reactive protein and parameters of adiposity and insulin resistance in patients with type 2 diabetes mellitus. *Endocrine J* 2006; 53: 345-56.
27. Lemieux I, Pascot A, Prud'homme D, Alméras N, Bogaty P, Nadeau A, Bergeron J, et al. Elevated C-reactive protein another component of the atherothrombotic profile of abdominal obesity. *Atheroscler Thromb Vasc Biol* 2001; 21: 961-7.
28. Sobhonslidsuk A, Jongjirasiri S, Thakkinstian A, Wisedopas N, Bunnag P, Puavilai G. Visceral fat and insulin resistance as predictor of non-alcoholic steatohepatitis. *World J Gastroenterol* 2007; 13: 3614-8.
29. Cusi K. Role of insulin resistance and lipotoxicity in nonalcoholic steatohepatitis. *Clin Liver Dis* 2009; 13: 545-63.
30. Larter CZ, Chitturi S, Heydet D, Farrell GC. A fresh look at NASH pathogenesis. Part 1: the metabolic movers. *J Gastroenterol Hepatol* 2010; 25: 672-90.
31. Kolak M, Westerbacka J, Velagapudi VR, Wagsäter D, Yetukuri L, Makkonen J, Rissanen A, et al. Adipose tissue inflammation and increased ceramide content characterize subjects with high liver fat content independent of obesity. *Diabetes* 2007; 56: 1960-8.
32. Pagadala M, Kasumov T, McCullough AJ, Zein NN, Kirwan JP. Role of ceramides in nonalcoholic fatty liver disease. *Trends Endocrinol Metab* 2012; 23: 365-71.
33. Fon Tacer K, Rozman D. Nonalcoholic Fatty liver disease: focus on lipoprotein and lipid deregulation. *J Lipids* 2011; 783976. doi: 10.1155/2011/783976.
34. Park S, Kim M, Paik JK, Jang YJ, Lee SH, Lee JH. Oxidative stress is associated with C-reactive protein in nondiabetic postmenopausal women, independent of obesity and insulin resistance. *Clin Endocrinol (Oxf)* 2013; 79: 65-70.
35. Lakoski SG. Gender and C-reactive protein: data from the multiethnic study of atherosclerosis (MESA) cohort. *Herat J* 2006; 152: 593-8.
36. Onat A, Ceyhan K, Başar O, Erer B, Toprak S, Sansoy V. Metabolic syndrome: major impact on coronary risk in a population with low cholesterol levels—a prospective and cross-sectional evaluation. *Atherosclerosis* 2002; 165: 285-92.
37. Han TS, Sattar N, Williams K, Gonzalez-Villalpando C, Lean ME, Haffner SM. Prospective study of C-reactive protein in relation to the development of diabetes and metabolic syndrome in the Mexico City Diabetes Study. *Diab Care* 2002; 25: 2016-21.
38. Thorand B, Baumert J, Döring A, Herder C, Kolb H, Rathmann W, Giani G, et al. Sex differences in the relation of body composition to markers of inflammation. *Atherosclerosis* 2006; 184: 216-24.
39. Clark CR, Ridker PM, Ommernborn MJ, Huisingh CE, Coull B, Buring JE, Berkman LF. Cardiovascular inflammation in healthy women: multilevel associations with state-level prosperity, productivity and income inequality. *BMC Public Health* 2012; 12: 211. doi: 10.1186/1471-2458-12-211.
40. Modena MG, Bursi F, Fantini G, Cagnacci A, Carbonieri A, Fortuna A, Rossi R. Effects of hormone replacement therapy on C-reactive protein levels in healthy postmenopausal women: comparison between oral and transdermal administration of estrogen. *Am J Med* 2002; 113: 331-4.
41. Rutter MK, Meigs JB, Sullivan LM, D'Agostino RB Sr, Wilson PW. C-reactive protein, the metabolic syndrome, and prediction of cardiovascular events in the Framingham Offspring Study. *Circulation* 2004; 110: 380-5.
42. Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO 3rd, Criqui M, Fadl YY, et al. Centers for Disease Control and Prevention; American Heart Association. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 2003; 107: 499-511.
43. Park SH, Kim KW, Lee SW, Yoon SE, Park SW, Ha HK, Lee MG, et al. Macrovesicular hepatic steatosis in living liver donors: use of CT for quantitative and qualitative assessment. *Radiology* 2006; 239: 105-12.