



Antifibrotic Mechanism of Pinocembrin: Impact on Oxidative Stress, Inflammation and TGF- β /Smad Inhibition in Rats

Marwa M. Said,* Samar S. Azab,** Noha M. Saeed,*** Ebtehal El-Demerdash**

* Pharmacist at Central Administration for Pharmaceutical Affairs, Ministry of Health, Cairo, Egypt.

** Pharmacology & Toxicology Department, Faculty of Pharmacy, Ain Shams University, Cairo, Egypt.

*** Pharmacology & Toxicology Department, Faculty of Pharmacy, Egyptian Russian University, Cairo, Egypt.

ABSTRACT

Introduction. The present study aimed to elucidate the potential antifibrotic effects of pinocembrin (PIN), a flavanone found abundantly in honey and propolis, by studying its effect on different oxidative stress, inflammatory and fibrosis markers in an experimental model of CCl_4 -induced liver fibrosis. **Material and methods.** PIN (20 mg/kg) was given orally 3 times/week for 6 consecutive weeks alternating with CCl_4 (0.5 mL/kg, 1:1 mixture with corn oil, i. p.) twice weekly. Different hepatotoxicity indices, oxidative stress, inflammatory and liver fibrosis markers were assessed. **Results.** PIN significantly restored liver transaminases and total cholesterol to normal levels. Also, PIN ameliorated oxidative stress injury evoked by CCl_4 as evidenced by inhibition of reduced glutathione depletion and lipid peroxidation as well as elevation of antioxidant enzyme superoxide dismutase (SOD). Further, PIN up-regulated the nuclear factor erythroid 2 (NF-E2)-related factor 2 (Nrf2), thereby inducing the expression and activity of the cytoprotective enzyme hemeoxygenase-1 (HO-1). Moreover, PIN alleviated pro-inflammatory cytokines such as TNF- α via inhibiting nuclear factor- κ B (NF- κ B) activation. As markers of fibrosis, collagen and α -SMA expression increased markedly in the CCl_4 group and PIN prevented these alterations. In addition, PIN down-regulated TGF β 1 and p-Smad2/3, thereby inhibiting TGF β 1/Smad signaling pathway. **Conclusion.** These results suggest that PIN possess potent antifibrotic effects that can be explained on its antioxidant properties. It ameliorates oxidative stress and inflammation during induction of fibrogenesis via its ability to augment cellular antioxidant defenses, activating Nrf2-mediated HO-1 expression and modulating NF- κ B and TGF- β 1/Smad signaling pathway.

Key words. Pinocembrin. CCl_4 . NF- κ B, Nrf2, HO-1.

INTRODUCTION

Liver fibrosis is a prevalent result of different ceaseless liver damages, which results from inordinate accumulation of extracellular matrix (ECM), because of inequality between ECM synthesis and deterioration.¹ Hepatic stellate cells (HSCs) undergo activation due to many factors,^{2,3} converting to vitamin A-losing and α -smooth muscle actin (α -SMA) expressing myoblasts.⁴ Also, activated HSCs produce fibrillar collagens leading to deposition of fibrotic matrix.⁵

Although growth factors, inflammatory factors and free radicals may participate in HSCs activation, data evoke a strong link with the transforming growth factor- β (TGF- β) signaling pathway.⁶ Indeed, Wynn reported that modu-

lation of TGF- β /Smad signaling by down-regulation of TGF- β expression may be efficient in inhibiting tissue fibrosis.¹ After binding of TGF- β to the type II receptor, kinases phosphorylate type I receptor, which further phosphorylates Smad2 and Smad3. Phosphorylated Smad2 and Smad3 then form a heteromeric complex with Smad4 that translocates to the nucleus and binds to the promoter of TGF- β -responsive genes to regulate their expression.⁷ TGF- β 1 also stimulates Smad3-dependent transcription of fibril collagens, which inhibit degradation of ECM.⁸ Therefore, TGF- β 1/Smad signaling pathway is considered a target to inhibit the activation of HSCs and ameliorate liver fibrosis.

In addition, oxidative stress promotes HSCs activation and collagen production and plays an important role in the

pathogenesis of liver fibrosis.⁹ Nuclear erythroid 2-related factor 2 (Nrf2) is an oxidative stress-mediated transcription factor with a variety of downstream targets aimed at cytoprotection. Activating Nrf2, promotes the expression of its target genes and increases the antioxidant enzymes activity.¹⁰ Furthermore, activation of Nrf2 may be a safe and effective strategy for anticipation of liver fibrosis and that Nrf2 activators may have great capability versus liver fibrosis.¹¹

Pinocembrin (5, 7-dihydroxyflavanone, PIN) is a flavanone found plentifully in honey and propolis.¹² Many studies have established that PIN possesses multiple activities, of which; neuroprotective,^{13,14} anti-inflammatory,¹⁵ anti-oxidant,^{16,17} hepatoprotective¹⁸ and anti-proliferative effects.¹⁹ PIN regulated the production of TNF- α via inhibiting NF- κ B, ERK1/2, JNK and p38MAPK in lipopolysaccharide-induced inflammatory responses¹⁵ and induced nuclear translocation of Nrf2 and expression of HO- β 1 in neuroblastoma SH-SY5Y cells.²⁰ Propolis also prevented the effects of TGF- β 1-induced Smad2 activation pathway in fibrotic lung diseases.²¹ Furthermore, Dodonaea viscosa ethanol extract, in which PIN was present, was examined *in vivo* against CCl₄-induced liver fibrosis in rats²² and managed to decrease liver fibrosis severity, hepatic enzymes and oxidative stress markers and normalized hepatic architecture. In addition, PIN inhibited thioacetamide-induced tissue Transglutaminase mRNA expression in Hep 3B carcinoma cell line, resulting in inhibition of hepatic fibrosis and cirrhosis.¹⁸ These findings indicated that PIN has protective effects on fibrosis but the mechanism by which PIN antagonizes liver fibrosis has not been fully examined. In the present study, we aimed at predicting the ability of PIN to attenuate the progression of chronic liver fibrosis induced by long-term administration of carbon tetrachloride (CCl₄) through reducing oxidative stress, cellular inflammation and down-regulating the expression of pro-fibrogenic markers by modulating TGF- β /Smad and Nrf-2 pathways.

MATERIALS AND METHODS

Drugs and chemicals

Pinocembrin (purity > 99.7%) was purchased from Sichuan Research Center of Traditional Chinese Medicine (Chengdu, China), 2-hydroxypropyl- β -cyclodextrin (HP β CD) from Roquette (France-Europe) and Carbon Tetrachloride (CCl₄), bovine serum albumin, reduced glutathione (GSH), Ellman's reagent [3,3'-dithiobis (6-nitrobenzoic acid)] and thiobarbituric acid (TBA) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). n-Butanol and formaldehyde 37% were purchased from ElNasr Chemical Co. (Egypt). All other chemicals and solvents were of highest grade commercially available.

Animals

Male Wistar rats (180-220 g) were obtained from Nile Co. for Pharmaceutical and Chemical Industries, Egypt. Rats were housed in an air-conditioned atmosphere; at a temperature of 25 °C. Animals were acclimated for 2 weeks before experimentation. They were kept on a standard diet and water ad libitum. Standard diet pellets (El-Nasr, Abu Zaabal, Egypt) contained not less than 20% protein, 5% fiber, 3.5% fat, 6.5% ash and 1% vitamin mixture. The study protocol was approved by the Ethical Committee, Faculty of Pharmacy, Ain Shams University, Egypt. The research was conducted in accordance with the internationally accepted principles for laboratory animal use and care as found in the National Institutes of Health guidelines, U.S.A.

Experimental design

Rats were randomly assigned into four groups (fifteen animals in each group) and treated for six weeks as follows; Group (A) served as control group and was given 1 mL/kg body weight from 20% HP β CD (the vehicle of PIN) through oral gavage three times per week and corn oil 0.5 mL/kg body weight i.p. twice per week alternating with HP β CD. Group (B) was given PIN 20mg/kg body weight dissolved in 20% HP CD through oral gavage three times per week. Group(C) was given CCl₄ (0.5 mL/kg body weight, 1:1 mixture with corn oil, i.p.), twice weekly to induce liver fibrosis and group (D) was given PIN 20 mg/kg body weight dissolved in 20% HP β CD through oral gavage three times per week alternating with CCl₄ (0.5 mL/kg body weight, 1:1 mixture with corn oil, i.p.), twice weekly.

Twenty-four hours after the last oral gavage, blood samples were collected from the retro-orbital plexus and allowed to clot. Serum was separated by centrifugation at 5,000 rpm for 10 min and used for the assessment of liver functions. Rats were sacrificed and liver tissues were dissected out and washed with ice-cold saline. Part of liver was homogenized in saline then the homogenate was used for assessment of oxidative stress and anti-inflammatory markers while other specimens from liver were fixed in 10% buffered formaldehyde for histopathological and immunohistochemical investigation.

Assessment of hepatotoxicity indices

Serum alanine aminotransferase (ALT), aspartate aminotransferase (AST) and total cholesterol (TC) were estimated using available commercial kits (Biodiagnostic,

Giza, Egypt). Liver index was calculated according to the formula:

$$\text{Liver index was} = (\text{liver weight}/\text{body weight}) \times 100$$

Assessment of oxidative stress markers

Reduced glutathione (GSH), lipid peroxides (malondi-aldehyde, MDA) and superoxide dismutase (SOD) were determined in liver homogenate using colorimetric kits (Biodiagnostics, Cairo, Egypt).

Quantitative ELISA immunoassay

TNF- α level in liver homogenate was assessed as an inflammatory marker. Determination of TNF- α was performed using commercial ELISA kit (R&D Systems, MN, USA) according to the manufacturer's instructions. The quantities of rat TNF- α were expressed as pg/mg protein and were determined according to method of Olsson.²³

Real-time RT-PCR analysis

NF- κ B as an inflammatory marker and collagen I as a liver fibrosis marker were determined using RT-PCR. Total RNA was extracted from liver tissue using SV Total RNA Isolation system as recommended by the manufacturer (Promega, Madison, WI, USA). Then, reverse-transcribed using reverse transcriptional polymerase chain reaction (RT-PCR) kit (Stratagene, USA) and quantitative real-time PCR was performed using SYBR Green. β -actin was used as standard housekeeping control. Levels of NF- κ B and collagen I were calculated based on the method of Pfaffl.²⁴ The primers were as follows: NF- κ B1; forward: 5'-TACTCTGGCGCAGAAATTAGGTC-3', reverse: 5'-CTGTCTCGGAGCTCGTCTATTG-3' and collagen I; Forward: 5'-TGGAGAGAGAGGTGAACAAGG-3 and Reverse: 5'-CATACCCTTAGCACCATCG-3.

Western blot analysis

Some markers for oxidative stress and liver fibrosis were assessed using western blot analysis including Nrf-2, HO-1 and p-Smad2/3. Total proteins were extracted from liver using RIPA buffer (Sigma). The protein concentration of lysate was quantified using a BCA Protein Assay kit according to Bradford technique.²⁵ Proteins were separated according to their molecular weight, transferred to nitrocellulose membrane [Protran BA 83 Cellulosenitrat (E), Whatman]. Primary antibodies of p-Smad2/3, Nrf-2 and HO-1 (Santa Cruz, CA, USA) were used. β -actin monoclonal antibody (Santa Cruz, CA, USA) then was added as an internal control.

Immunohistochemistry

As indices for liver fibrosis, TGF- β 1 and α -SMA were assessed using immunohistochemical detection. Paraffin embedded tissue sections of 3 μ m thickness were rehydrated first in xylene and then in graded ethanol solutions. The slides were then blocked with 5% bovine serum albumin (BSA) in tris buffered saline (TBS) for 2 h. The sections were then immunostained with one of the following primary antibodies; mouse monoclonal to rat α -SMA (SantaCruz Biotech, Cat No. sc-53142) and rabbit polyclonal to rat TGF- β 1 (SantaCruz Biotech, Cat No. sc-146) antibodies at a concentration of 1 μ g/mL containing 5% BSA in TBS and incubated overnight at 4 °C. After washing the slides with TBS, the sections were incubated with goat anti-rabbit secondary antibody. Sections were then washed with TBS and incubated for 5-10 min in a solution of 0.02% diaminobenzidine containing 0.01% H_2O_2 . Counter staining was performed using hematoxylin, and the slides were visualized under a light microscope.

Histopathological examination

For light microscopy, liver specimens were taken from the right lobe and fixed in 10% formalin and processed for paraffin sections of 4 μ m thickness. Sections were stained with hematoxylin and eosin for routine histopathological examination.

Statistical analysis

Data are presented as mean \pm SD. Multiple comparisons were performed using one-way ANOVA followed by Tukey-Kramer as a post hoc test. The 0.05 level of probability was used as the criterion for significance. All statistical analyses and graphs sketching were performed using GraphPad Prism (ISI1 software, USA) version 5 software.

RESULTS

Hepatotoxicity indices

Estimation of liver enzymes and TC revealed that CCl_4 induced a significant increase in serum ALT, AST and TC levels by 92, 67 and 114%, respectively as compared to the control group. Concurrent treatment of intoxicated group with PIN significantly reduced ALT and TC levels as well as relative liver weight similar to that of the control group and significantly reduced AST level but still significantly higher than the control group. PIN alone did not modify liver function (Table 1).

Histopathological examination

Histopathological examination (Figure 1) of liver tissue revealed that control group showed normal architecture of the central veins and surrounding hepatocytes in the hepatic parenchyma and no histopathological alterations were recorded (Figure 1A). Group received PIN alone showed normal histology of central vein and surrounding hepatocytes (Figure 1B). CCl_4 -intoxicated group showed focal fibrosis (f) with inflammatory cells infiltration (m) in the portal area with extension to the adjacent parenchyma and congestion in the hepatic sinusoids (s), along with fatty change (arrow) in some individual hepatocytes (Figure 1C). Concurrent treatment of intoxicated group with PIN illustrated minimal degeneration (d) in hepatocytes (Figure 1D).

Oxidative stress markers

Table 2 shows the effect of PIN treatment on CCl_4 -induced changes in protein thiols and lipid peroxides measured as GSH and MDA levels, respectively as well as liver antioxidant enzyme SOD. CCl_4 -intoxicated group showed

a significant depletion in hepatic GSH content by 56% and reduction in SOD by 86% in addition to a significant increase in MDA by 10 folds as compared to the control group. Nonetheless, pretreatment with 20 mg/kg PIN significantly restored the depleted GSH content and SOD activity and caused a significant decrease in lipid peroxides when compared to CCl_4 -intoxicated group. Rats treated with PIN only did not show any significant changes in all studied to the control group.

In addition, western blot analysis revealed significant decreases in Nrf-2 and HO-1 in CCl_4 -intoxicated group by 89 and 86%, respectively as compared to control group. Treatment with PIN 20 mg/kg significantly increased Nrf-2 and HO-1 levels by 72 and 78% respectively in comparison with CCl_4 -intoxicated group. No change was observed in Nrf-2 and HO-1 levels in animals treated with PIN only (Figures 2A and B).

Inflammatory markers

CCl_4 -intoxicated group showed proinflammatory response as evidenced by significant increase in both total TNF- α and NF- κ B in the liver tissue by 4 and 11 folds re-

Table 1. Liver indices for rats treated with CCl_4 and/or PIN.

	Liver index %	ALT (U/L)	AST (U/L)	TC (mg/dL)
Control	22.2 \pm 2.58 ^b	45.2 \pm 6.98 ^b	108.4 \pm 6.11 ^b	51.4 \pm 3.91 ^b
PIN (20 mg/kg)	22.0 \pm 1.58 ^b	39.6 \pm 4.16 ^b	109.4 \pm 8.11 ^b	48.6 \pm 3.36 ^b
CCl_4 (0.5 mL/kg)	49.8 \pm 1.92 ^a	87.0 \pm 7.02 ^a	180.8 \pm 7.39 ^a	110.4 \pm 9.15 ^a
CCl_4 +PIN (20 mg/kg)	29.2 \pm 3.34 ^{ab}	53.4 \pm 8.38 ^b	122.3 \pm 9.95 ^{ab}	58.6 \pm 2.97 ^b

Data are the mean \pm S.D. (n = 8). ^a or ^b: Significantly different from the control or CCl_4 group respectively at P < 0.05 using ANOVA followed by Tukey-Kramer as a post hoc test.




Figure 1. Photomicrographs of liver sections stained by H & E (x 40). **A.** Control group showing normal histological structure of the central veins and surrounding hepatocytes in the hepatic parenchyma. **B.** Group received PIN alone showed normal histology of central vein and surrounding hepatocytes. **C.** CCl_4 -intoxicated group showing focal fibrosis (f) with inflammatory cells infiltration (m) in the portal area with extension to the adjacent parenchyma and congestion in the hepatic sinusoids (s) along with fat deposition (arrow). **D.** CCl_4 -intoxicated group treated with PIN illustrated minimal degeneration (d) in hepatocytes.

spectively, as compared to the control group (Figures 3A & B). Treatment with 20 mg/kg PIN exhibited anti-inflammatory effects by significantly reducing liver TNF- α and NF- κ B contents by 2 and 7 folds respectively, as com-

pared to CCl_4 -intoxicated group. On the other hand, there was no significant change in liver TNF- α and NF- κ B content in animals treated with PIN alone.

Table 2. Oxidative stress markers of rats treated with PIN and/or CCl_4 .

	GSH (μ mol/g tissue)	MDA (nmol/g tissue)	SOD (U/mg protein)
Control	60.8 \pm 2.40 ^b	9.7 \pm 0.97 ^b	8.1 \pm 0.15 ^b
PIN (20 mg/kg)	59.1 \pm 7.22 ^b	9.4 \pm 0.76 ^b	8.5 \pm 0.37 ^b
CCl_4 (0.5 mL/kg)	26.5 \pm 4.11 ^a	104.9 \pm 11.22 ^a	1.1 \pm 0.17 ^a
CCl_4 +PIN (20 mg/kg)	46.7 \pm 4.47 ^{ab}	36.4 \pm 5.31 ^{ab}	6.7 \pm 0.61 ^{ab}

Data are the mean \pm S.D. ($n = 8$). ^a or ^b: Significantly different from the control or CCl_4 group respectively at $P < 0.05$ using ANOVA followed by Tukey-Kramer as a post hoc test.




Figure 2. Effect of treatment with 20 mg/kg PIN on Nrf-2, HO-1 and p-Smad2/3 released during CCl_4 -induced hepatotoxicity in rats assessed using Western Blot analysis. Values are given as mean \pm S.D. for groups of 15 rats for each. (a or b) significantly different from the control or CCl_4 group respectively at $P < 0.05$ using ANOVA followed by Tukey-Kramer as a post hoc test.




Figure 3. A. Effect of treatment with 20 mg/kg PIN on TNF- α released during CCl₄-induced hepatotoxicity in rats using ELISA. Values are given as mean + S.D. for groups of 15 rats for each. B. Effect of treatment with 20 mg/kg PIN on NF- κ B released during CCl₄-induced hepatotoxicity in rats using Real-time PCR. Values are given as mean + S.D. for groups of 15 rats for each (a or b) significantly different from the control or CCl₄ group respectively at $P < 0.05$ using ANOVA followed by Tukey-Kramer as a post hoc test.




Figure 4. Effect of treatment with 20 mg/kg PIN on Collagen I released during CCl₄-induced hepatotoxicity in rats assessed using Real-time PCR. Values are given as mean + S.D. for groups of 15 rats for each (a or b) significantly different from the control or CCl₄ group respectively at $P < 0.05$ using ANOVA followed by Tukey-Kramer as a post hoc test.

Liver fibrosis markers

Analysis of collagen I levels by RT-PCR showed an increase by 20 folds in CCl₄-intoxicated group as compared to control group. Treatment with PIN 20 mg/kg decreased collagen I level by 9 folds as compared to CCl₄-intoxicated group. PIN only treated group did not show any significant changes in collagen I levels from control group (Figure 4).

In addition, immunostaining of fibrogenic markers α -SMA and TGF- β (Figures 5 and 6, respectively) revealed that CCl₄ caused significant increase of α -SMA and TGF- β expression by 75 and 48% (Figures 5E, 5F and 6E, 6F), respectively which were evident from intense brown staining. Treatment of rats with 20 mg/kg PIN significantly inhibited the expression of α -SMA and TGF- β by 69 and 35% (Figures 5G, 5H and 6G, 6H), respectively when compared to CCl₄-intoxicated group. No change was observed in α -SMA and TGF- β expression in PIN only treated group when compared to control group (Figures 5C, 5D and 6C, 6D). Figures 5I and 6I represent OD quantification results of the stained regions for α -SMA and TGF- β , respectively. Further, western blot analysis detected significant increase in the p-Smad 2/3 levels by 10 folds in CCl₄-intoxicated group as compared to control group and treatment with PIN 20 mg/kg managed to decrease p-Smad 2/3 level by 8 folds as compared to CCl₄-intoxicated group. PIN only treated group did not show any significant changes in p-Smad 2/3 levels from control group (Figure 2C).

DISCUSSION

The present study was designed to examine the potential antifibrotic effect of PIN in an experimental model of liver fibrosis via studying its effects on different oxidative stress, inflammatory and fibrosis markers. The CCl₄-induced hepatotoxicity model is a commonly used model for the screening of the hepatoprotective activities of drugs. It is well known that prolonged administration of CCl₄ leads to liver fibrosis, cirrhosis, and hepatocellular carcinoma.

In the present study, chronic exposure of rats to CCl₄ caused a significant increase in serum ALT and AST, which resulted from the hepatocyte damage and inflammation.²⁶ Relative liver weight and serum TC were also significantly increased that confirming the induction of liver hypertrophy according to previous study.²⁷ Histopathological examination, the “gold” tool for the diagnosis of hepatic fibrosis, revealed the presence of fibrosis with several histopathological changes that are in accordance with previous study.²⁸ Treatment of animals with PIN (20 mg/kg) significantly decreased ALT, AST and TC




Figure 5. Expression of α -Smooth Muscle Actin (α -SMA) by immunohistochemical staining (**A, C, E, G** with magnification 40x and **B, D, F, H** with magnification 100x). Photomicrographs of histological sections of liver depicting (**A, B**) control group, (**C, D**) PIN only treated group, (**E, F**) CCl_4 (0.5 mL/kg) intoxicated group, (**G, H**) CCl_4 (0.5 mL/kg) + PIN (20 mg/kg) treated group, (**I**) quantitative image analysis for immunohistochemical staining expressed as optical densities (OD) across 10 different fields for each rat section. Values are given as mean \pm S.D. For groups of 15 rats for each. (a) $P < 0.05$ statistically significant from control group and (b) $P < 0.05$ statistically significant from CCl_4 group using one-way analysis of variance (ANOVA) followed by Tukey-Kramer as a post hoc test. For immunohistochemical analyses, brown color (positive) indicates specific immunostaining of α -SMA and light blue color (negative) indicates hematoxylin staining. CCl_4 intoxication significantly increased α -SMA expression as indicated by intense brown staining compared to the control group. PIN treatment significantly reduced α -SMA expression compared to CCl_4 group. However, there was no significant difference in α -SMA expression in PIN only treated group as compared to the control group.

and ameliorated the histopathologic changes induced by CCl_4 confirming that PIN effectively opposes the progression of CCl_4 -induced liver injury. These results are in agreement with Shalaby, Abd-Alla²² study which confirmed that an extract of *D. viscosa* extracts containing PIN as one of its prominent constituents managed to protect against hepatic fibrosis through down-regulation of liver enzymes and normalization of hepatic architecture.

Previous literature elucidated that CCl_4 -induced oxidative stress is the cornerstone underlying its hepatotoxicity and it is due to reductive dehalogenation of CCl_4 by the cytochrome P450 enzyme system to the highly reactive trichloromethyl radical in the presence of oxygen. These free radicals damage the hepatocellular membrane via lipid peroxidation, followed by release of inflammatory mediators from activated inflammatory cells which are

thought to potentiate CCl_4 -induced hepatic injury.²⁹ Oxidative stress occur when there is an inequality between production of reactive oxygen species (ROS) and cellular antioxidant activity.³⁰ Cellular antioxidant capacity depends on the presence of GSH and antioxidant enzymes such as SOD. A critical regulator of cellular defense against oxidative stress is Nrf2.³¹ It is responsible for regulating the antioxidant response element (ARE)-driven expression of genes encoding the majority of antioxidant and phase II detoxification enzymes such as HO-1.³² Several antioxidants have been reported to increase Nrf2 activation, mediated by HO-1 expression, thereby attenuating oxidative stress in vitro and in vivo.^{33,34} Thus, pharmacological activation of Nrf2 has been suggested to offer a novel therapeutic approach to the prevention and treatment of liver injury and fibrosis.³⁵ In the present study,




Figure 6. Expression of Transforming Growth Factor- β 1 (TGF- β 1) by immunohistochemical staining (**A, C, E, G** with magnification 40x and **B, D, F, H** with magnification 100x). Photomicrographs of histological sections of liver depicting (**A, B**) control group, (**C, D**) PIN only treated group, (**E, F**) CCl₄ (0.5 mL/kg) intoxicated group, (**G, H**) CCl₄ (0.5 mL/kg) + PIN (20 mg/kg) treated group, (**I**) quantitative image analysis for immunohistochemical staining expressed as optical densities (OD) across 10 different fields for each rat section. Values are given as mean \pm S.D. For groups of 15 rats for each. (a) $P < 0.05$ statistically significant from control group and (b) $P < 0.05$ statistically significant from CCl₄ group using one-way analysis of variance (ANOVA) followed by Tukey-Kramer as a post hoc test. For immunohistochemical analyses, brown color (positive) indicates specific immunostaining of TGF- β 1 and light blue color (negative) indicates hematoxylin staining. CCl₄ intoxication significantly increased TGF- β 1 expression as indicated by intense brown staining compared to the control group. PIN treatment significantly reduced TGF- β 1 expressions compared to CCl₄ group. However, there was no significant difference in TGF- β 1 expression in PIN only treated group as compared to the control group.

CCl₄ found to induce an increase in the oxidative stress profile as indicated by significant rise in lipid peroxidation, GSH depletion and significant reduction in antioxidant enzyme (SOD). The decline in SOD with concomitant MDA elevation, strongly indicate oxidative damage.³⁶ These results are in agreement with previous studies.^{22,37} Also, western blot analysis revealed that CCl₄ significantly down-regulated the expression of Nrf2 and HO-1 which is consistent with prior observation.³⁷ Pre-treatment with PIN significantly guarded against the oxidative stress effects by decreasing MDA and increasing GSH and SOD, which was previously proven by Shalaby, Abd-Alla,²² as well as restoring Nrf2 and HO-1 to normal level. Induction of HO-1 expression under regulation of Nrf2-ARE is consistent with previous studies^{38,39} who reported that up-regulation of HO-1 expression inhibited

oxidative stress in association with increased Nrf2-ARE-binding activity in hepatocytes and protected liver cells from apoptosis. Also, Jin, Liu²⁰ research had proved that PIN has cytoprotective role in dopaminergic cell culture systems by increasing Nrf2 and the subsequent HO-1 levels. These findings imply that PIN effectively attenuated the progression of CCl₄-induced oxidative stress by restoring liver antioxidant capacity.

Beside the oxidative stress, accumulating evidences have revealed that excessive ROS induced by CCl₄ can stimulate circulating monocytes and tissue macrophages, which lead to the synthesis and release of a variety of proinflammatory cytokines.^{40,41} In particular, TNF- α plays a key role in the development and maintenance of inflammation and that its elevation is associated with many liver diseases.^{41,42} Furthermore, NF- κ B is a nuclear transcrip-




Figure 7. A schematic diagram of the overall mechanism of amelioration of liver fibrosis by PIN.

tion factor that regulates the expression of a large number of genes that are critical for the regulation of apoptosis, viral replication, tumorigenesis, inflammation, and various autoimmune diseases.⁴³ In addition, HSCs activation is associated with elevation of NF-κB activity.⁴⁴ In the present study, CCl₄ significantly up-regulated the expression of TNF-α and NF-κB in rat liver and, treatment with PIN significantly reduced their expression. These results suggested that PIN could alleviate liver injury caused by CCl₄ through suppressing inflammatory response.

To further explore the mechanism underlying the anti-fibrotic effect of PIN, several fibrogenic markers were assessed. It is believed that HSCs are a major type of fibrogenic liver cells found during liver injury and are responsible for the progression of hepatic fibrosis.⁴⁵ Activation of HSCs towards myofibroblast-like cells is characterized by several key phenotypic features. Those features include proliferation, liver content of α-SMA; a microfilament protein that has been explored as a marker for activated HSCs,^{46,47} and finally accumulation of ECM mainly collagen, which is stimulated by the multifactorial growth factor TGF-β.⁴⁸ In this study, the levels of collagen I and α-SMA in liver tissue were significantly increased in CCl₄ intoxicated rats which is consistent with another study,⁴⁹ whereas PIN treatment markedly decreased the levels of these fibrogenic parameters. These data indicate that PIN can prevent col-

lagen and α-SMA accumulation caused by the chronic liver disease.

Furthermore, TGF-β1 is stored as an inactivated protein bound to a latency-associated peptide. Once activated, TGF-β1 binds its cognate receptors and functions in autocrine and paracrine manners to exert its biological and pathological activities via Smad-dependent and -independent signaling pathways. Smads are intracellular signal transductive molecules of the TGF-β super family. Smad signal transduction pathways are thought to mediate TGF-β 1-induced collagen synthesis and to play a crucial role in the process of liver damage and recovery, as well as in the development of liver fibrosis.^{50,51} Studies have demonstrated that TGF-β1/Smad signaling pathway might play a prominent role in the activation of HSCs and the regulation of the production, degradation, and accumulation of ECM proteins.⁵⁰ Therefore, the inhibition of the TGF-β/Smad signaling pathway is an attractive therapeutic target for the prevention of liver fibrosis.^{52,53} In the present study, PIN treatment attenuated the increase in level of the prominent profibrogenic cytokine TGF-β1 induced by CCl₄, a finding indicates the inhibitory effect of PIN to the proliferative activity of HSCs. Moreover, CCl₄-induced liver fibrosis was associated with a marked activation of Smad2/3 and PIN treatment was able to inhibit the phosphorylation of Smad2/3 and reversed the effect of CCl₄. Therefore, the decreased fibrosis level after PIN

treatment may be mediated via inhibition of TGF- β 1/Smad signaling pathway.

In summary, this study demonstrates that PIN has promising anti-fibrotic effects by inhibiting the TGF- β 1/Smad and activating Nrf2/HO-1 signaling pathways, as well as decreasing oxidative stress and proinflammatory cytokines production (Figure 7).

ABBREVIATIONS

- **ALT:** alanine aminotransferase.
- **AST:** aspartate aminotransferase.
- **CCl₄:** carbon tetrachloride.
- **GSH:** reduced glutathione.
- **HO-1:** heme oxygenase-1.
- **HP β CD:** 2-hydroxypropyl- β -cyclodextrin.
- **MDA:** malondialdehyde.
- **NF- κ B:** nuclear factor- κ B.
- **Nrf2:** nuclear factor erythroid 2 (NF-E2)-related factor 2.
- **PIN:** pinocembrin.
- **ROS:** reactive oxygen species.
- **SOD:** superoxide dismutase.
- **TC:** total cholesterol.
- **TGF- β :** transforming growth factor- β .
- **TNF- α :** tumor necrosis factor- α .
- **α -SMA:** α -smooth muscle actin.

FINANCIAL SUPPORT

The authors gratefully thank Roquette (France-Europe) for the free supplement of 2-hydroxypropyl- β -cyclodextrin (HP β CD).

REFERENCES

1. Wynn T. Cellular and molecular mechanisms of fibrosis. *J Pathol* 2008; 214: 199-210.
2. Chu Ps, Nakamoto N, Ebinauma H, Usui S, Saeki K, Matsumoto A, Mikami Y, et al. C?C motif chemokine receptor 9 positive macrophages activate hepatic stellate cells and promote liver fibrosis in mice. *Hepatology* 2013; 58: 337-50.
3. Ahmad A, Ahmad R. Understanding the mechanism of hepatic fibrosis and potential therapeutic approaches. *Saudi J Gastroenterol* 2012; 18: 155.
4. Gressner O, Weiskirchen R, Gressner A. Biomarkers of hepatic fibrosis, fibrogenesis and genetic pre-disposition pending between fiction and reality. *J Cel Mol Med* 2007; 11: 1031-51.
5. Henderson N, Iredale J. Liver fibrosis: cellular mechanisms of progression and resolution. *Clinical Science* 2007; 112: 265-80.
6. Dooley S, Streckert M, Delvoux B, Gressner AM. Expression of Smads during in vitro transdifferentiation of hepatic stellate cells to myofibroblasts. *Biochemical and Biophysical Research Communications* 2001; 283: 554-62.
7. Massague J. TGFbeta signalling in context. *Nat Rev Mol Cell Biol* 2012; 13: 616-30.
8. Ten Dijke P, Hill CS. New insights into TGF- β -Smad signalling. *Trends in Biochemical Sciences* 2004; 29: 265-73.
9. Ghatak S, Biswas A, Dhalil GK, Chowdhury A, Boyer JL, Santra A. Oxidative stress and hepatic stellate cell activation are key events in arsenic induced liver fibrosis in mice. *Toxicology and Applied Pharmacology* 2011; 251: 59-69.
10. Cong M, Liu T, Wang P, Fan X, Yang A, Bai Y, Peng Z, et al. Antifibrotic effects of a recombinant adeno-associated virus carrying small interfering RNA targeting TIMP-1 in rat liver fibrosis. *Am J Pathol* 2013; 182: 1607-16.
11. Yang J-J, Tao H, Huang C, Li J. Nuclear erythroid 2-related factor 2: A novel potential therapeutic target for liver fibrosis. *Food and Chemical Toxicology* 2013; 59: 421-7.
12. Jaganathan SK, Mandal M. Antiproliferative effects of honey and of its polyphenols: a review. *Journal of Biomedicine and Biotechnology* 2009; 2009.
13. Gao M, Liu R, Zhu S-Y, Du G-H. Acute neurovascular unit protective action of pinocembrin against permanent cerebral ischemia in rats. *Journal of Asian Natural Products Research* 2008; 10: 551-8.
14. Liu R, Gao M, Yang Z-H, Du G-H. Pinocembrin protects rat brain against oxidation and apoptosis induced by ischemia-reperfusion both in vivo and in vitro. *Brain Research* 2008; 1216: 104-15.
15. Soromou LW, Chu X, Jiang L, Wei M, Huo M, Chen N, Guan S, et al. In vitro and in vivo protection provided by pinocembrin against lipopolysaccharide-induced inflammatory responses. *International Immunopharmacology* 2012; 14: 66-74.
16. Kapoor S. Comment on isolation and identification of compounds from *Penthorum chinense* Pursh with antioxidant and antihepatocarcinoma properties: Pinocembrin and its rapidly emerging neuroprotective effects. *Journal of Agricultural and Food Chemistry* 2013; 61: 1416-16.
17. Yu Y-S, Hsu C-L, Yen G-C. Anti-inflammatory effects of the roots of *Alpinia pricei* Hayata and its phenolic compounds. *Journal of Agricultural and Food Chemistry* 2009; 57: 7673-80.
18. Chen C-S, Wu C-H, Lai Y-C, Lee W-S, Chen H-M, Chen R-J, Chen L-C, et al. NF- κ B-activated tissue transglutaminase is involved in ethanol-induced hepatic injury and the possible role of propolis in preventing fibrogenesis. *Toxicology* 2008; 246: 148-57.
19. Catchpole O, Mitchell K, Bloor S, Davis P, Suddes A. Antiproliferative activity of New Zealand propolis and phenolic compounds vs human colorectal adenocarcinoma cells. *Fitoterapia* 2015; 106: 167-74.
20. Jin X, Liu Q, Jia L, Li M, Wang X. Pinocembrin attenuates 6-OHDA-induced neuronal cell death through Nrf2/ARE pathway in SH-SY5Y cells. *Cellular and molecular neurobiology* 2015; 35: 323-33.
21. Kao H-F, Chang-Chien P-W, Chang W-T, Yeh T-M, Wang J-Y. Propolis inhibits TGF- β 1-induced epithelial-mesenchymal transition in human alveolar epithelial cells via PPAR γ activation. *International Immunopharmacology* 2013; 15: 565-74.
22. Shalaby N, Abd-Alla H, Hamed M, Al-Ghamdi S, Jambi S. Flavones composition and therapeutic potential of *Dodonaea viscosa* against liver fibrosis. *International Journal of Phyto-medicine* 2012; 4: 27-39.
23. Olsson I, Lantz M, Nilsson E, Peetre C, Thysell H, Grubb A, Adolf G. Isolation and characterization of a tumor necrosis factor binding protein from urine. *Eur J Haematol* 1989; 42: 270-5.
24. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research* 2001; 29: e45-e45.
25. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle

of protein-dye binding. *Analytical Biochemistry* 1976; 72: 248-54.

26. Bahcecioglu IH, Koca SS, Poyrazoglu OK, Yalniz M, Ozercan IH, Ustundag B, Sahin K, et al. Hepatoprotective effect of infliximab, an anti-TNF- α agent, on carbon tetrachloride-induced hepatic fibrosis. *Inflammation* 2008; 31: 215-21.
27. Yachi R, Igarashi O, Kiyose C. Protective effects of vitamin E analogs against carbon tetrachloride-induced fatty liver in rats. *J Clin Biochem Nutrit* 2010; 47: 148-54.
28. Fujii T, Fuchs BC, Yamada S, Lauwers GY, Kulu Y, Goodwin JM, Lanuti M, et al. Mouse model of carbon tetrachloride induced liver fibrosis: Histopathological changes and expression of CD133 and epidermal growth factor. *BMC Gastroenterology* 2010; 10: 79.
29. Recknagel RO, Glende EA, Dolak JA, Waller RL. Mechanisms of carbon tetrachloride toxicity. *Pharmacology & Therapeutics* 1989; 43: 139-54.
30. Ungvari Z, Sonntag WE, de Cabo R, Baur JA, Csiszar A. Mitochondrial protection by resveratrol. *Exercise and Sport Sciences Reviews* 2011; 39: 128-32.
31. He X, Wang L, Szklarz G, Bi Y, Ma Q. Resveratrol inhibits paraquat-induced oxidative stress and fibrogenic response by activating the nuclear factor erythroid 2-related factor 2 pathway. *Journal of Pharmacology and Experimental Therapeutics* 2012; 342: 81-90.
32. Na H-K, Surh Y-J. Modulation of Nrf2-mediated antioxidant and detoxifying enzyme induction by the green tea polyphenol EGCG. *Food and Chemical Toxicology* 2008; 46: 1271-8.
33. Yeh CT, Chiu HF, Yen GC. Protective effect of sulforaphane on indomethacin-induced cytotoxicity via heme oxygenase-1 expression in human intestinal Int 407 cells. *Molecular Nutrition & Food Research* 2009; 53: 1166-76.
34. Lee B-H, Hsu W-H, Chang Y-Y, Kuo H-F, Hsu Y-W, Pan T-M. Ankaflavin: a natural novel PPAR γ agonist upregulates Nrf2 to attenuate methylglyoxal-induced diabetes in vivo. *Free Radical Biology and Medicine* 2012; 53: 2008-16.
35. Osburn WO, Yates MS, Dolan PD, Chen S, Liby KT, Sporn MB, Taguchi K, et al. Genetic or pharmacologic amplification of nrf2 signaling inhibits acute inflammatory liver injury in mice. *Toxicological Sciences* 2008; 104: 218-27.
36. Ahmad A, Ahmad R. Resveratrol mitigate structural changes and hepatic stellate cell activation in N'-nitrosodimethylamine-induced liver fibrosis via restraining oxidative damage. *Chemico-Biological Interactions* 2014; 221: 1-12.
37. Mohamed MR, Emam MA, Hassan NS, Mogadem AI. Umbelliferone and daphnetin ameliorate carbon tetrachloride-induced hepatotoxicity in rats via nuclear factor erythroid 2-related factor 2-mediated heme oxygenase-1 expression. *Environmental Toxicology and Pharmacology* 2014; 38: 531-41.
38. Huang C-S, Lii C-K, Lin A-H, Yeh Y-W, Yao H-T, Li C-C, Wang T-S, et al. Protection by chrysins, apigenin, and luteolin against oxidative stress is mediated by the Nrf2-dependent up-regulation of heme oxygenase 1 and glutamate cysteine ligase in rat primary hepatocytes. *Archives of Toxicology* 2013; 87: 167-78.
39. Mani M, Golmohammadi T, Khaghani S, Zamani Z, Azadmanesh K, Meshkani R, Pasalar P. Homocysteine Induces Heme Oxygenase-1 Expression via Transcription Factor Nrf2 Activation in HepG2 Cells. *Iranian Biomedical Journal* 2013; 17: 93-100.
40. Wang J-Q, Chen X, Zhang C, Tao L, Zhang Z-H, Liu X-Q, Xu Y-B, et al. Phenylbutyric acid protects against carbon tetrachloride-induced hepatic fibrogenesis in mice. *Toxicology and Applied Pharmacology* 2013; 266: 307-16.
41. Shim J-Y, Kim M-H, Kim H-D, Ahn J-Y, Yun Y-S, Song J-Y. Protective action of the immunomodulator ginsan against carbon tetrachloride-induced liver injury via control of oxidative stress and the inflammatory response. *Toxicology and Applied Pharmacology* 2010; 242: 318-25.
42. Shin D-S, Kim KW, Chung HY, Yoon S, Moon J-O. Effect of sinapic acid against carbon tetrachloride-induced acute hepatic injury in rats. *Archives of Pharmacal Research* 2013; 36: 626-33.
43. Ma J-Q, Liu C-M, Qin Z-H, Jiang J-H, Sun Y-Z. Ganoderma applanatum terpenes protect mouse liver against benzo (α) pyren-induced oxidative stress and inflammation. *Environmental Toxicology and Pharmacology* 2011; 31: 460-8.
44. Kong D, Zhang F, Wei D, Zhu X, Zhang X, Chen L, Lu Y, et al. Paeonol inhibits hepatic fibrogenesis via disrupting nuclear factor- κ B pathway in activated stellate cells: In vivo and in vitro studies. *J Gastroenterol Hepatol* 2013; 28: 1223-33.
45. Park KC, Park JH, Jeon JY, Kim SY, Kim JM, Lim CY, Lee TH, et al. A new histone deacetylase inhibitor improves liver fibrosis in BDL rats through suppression of hepatic stellate cells. *Brit J Pharmacol* 2014; 171: 4820-30.
46. Friedman SL. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *Journal of Biological Chemistry* 2000; 275: 2247-50.
47. Ahmad R, Ahmed S, Khan NU, Hasnain A-U. Operculina turpethum attenuates N-nitrosodimethylamine induced toxic liver injury and clastogenicity in rats. *Chemico-Biological Interactions* 2009; 181: 145-53.
48. Liu Y, Wang Z, Kwong SQ, Lui EL, Friedman SL, Li FR, Lam RW, et al. Inhibition of PDGF, TGF- β , and Abl signaling and reduction of liver fibrosis by the small molecule Bcr-Abl tyrosine kinase antagonist Nilotinib. *J Hepatol* 2011; 55: 612-25.
49. Deng Y-R, Ma H-D, Tsuneyama K, Yang W, Wang Y-H, Lu F-T, Liu C-H, et al. STAT3-mediated attenuation of CCl4-induced mouse liver fibrosis by the protein kinase inhibitor sorafenib. *Journal of Autoimmunity* 2013; 46: 25-34.
50. Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF- β family signalling. *Nature* 2003; 425: 577-84.
51. Inagaki Y, Okazaki I. Emerging insights into transforming growth factor β Smad signal in hepatic fibrogenesis. *Gut* 2007; 56: 284-92.
52. Liu R-M, Pravia KG. Oxidative stress and glutathione in TGF- β -mediated fibrogenesis. *Free Radical Biology and Medicine* 2010; 48: 1-15.
53. Liu X, Hu H, Yin JQ. Therapeutic strategies against TGF- β signaling pathway in hepatic fibrosis. *Liver International* 2006; 26: 8-22.

Correspondence and reprint request:

Dr. Ebtehal El-Demerdash
 Prof. and Head of Pharmacology & Toxicology Department
 Faculty of Pharmacy, Ain Shams University, Abasia, Cairo,
 Egypt
 Mobile. 002-01001925375
 E-mail: ebtehal_dm@yahoo.com
 ebtehal_dm@pharma.asu.edu.eg