

LETTERS TO THE EDITOR

DOI: 10.15517/IJDS.2022.51544

Received:
11-IV-2022

Anatomical Variant in Middle Turbinate: Concha Bullosa, Finding on Cone Beam Computerized Tomography

Accepted:
21-VI-2022

Published Online:
27-VI-2022

Variante anatómica en cornete medio: concha bullosa. Hallazgo en tomografía computarizada de haz cónico

Katherine M. Aguirre Alanya DDS¹; Margarita A. Leandro Melgarejo DDS²;
Vilma E. Ruiz-De Chacón DDS, MSc³

1. Residente de segundo año, Segunda Especialidad de Radiología Bucal y Maxilofacial. Universidad Peruana Cayetano Heredia, Lima, Perú. <https://orcid.org/0000-0002-7937-2100>
2. Residente de segundo año, Segunda Especialidad de Radiología Bucal y Maxilofacial. Universidad Peruana Cayetano Heredia, Lima, Perú. <https://orcid.org/0000-0001-6043-6349>
3. Docente, Segunda Especialidad de Radiología Bucal y Maxilofacial. Universidad Peruana Cayetano Heredia, Lima, Perú. <https://orcid.org/0000-0002-4798-1710>

Correspondence to: Dr. Katherine Melissa Aguirre Alanya – katherine.aguirre@upch.pe

PhD. Jessie Reyes Carmona
Editor Jefe
ODOVTOS-International Journal of Dental Sciences
Universidad de Costa Rica

The nasal turbinates, also called nasal shells, are attached to the lateral and/or superior walls of the nostril and can present several anatomical variations. Thus, we can mention the concha bullosa, the paradoxical curvature of the middle turbinate (both common anatomical variants) and the secondary middle turbinate, the latter in turn, can be pneumatized (very rare) (1).

The term “concha bullosa” generally refers to pneumatization of the middle turbinate, however, pneumatization of the superior and inferior turbinates can also occur (2). Some authors mention it as a “variation in the pneumatization of the bone plate caused by an extension of the ethmoid cells” (3). The literature reports 3 types of concha bullosa: lamellar, bulbous and true; the latter refers to the complete pneumatization of the anterior third of the turbinate. The origin of this condition is undetermined (4), but it can be considered as predisposing causes to trauma, deviation of the nasal septum and mouth breathing (3). The frequency of pneumatization of the middle turbinate is in a range from 14% to 53%, while in the inferior turbinate it is 1% (2,4).

The anatomical variants of the nasal turbinates are important in the pathogenesis of the inflammatory processes of the paranasal sinuses, such as for the etiology of sinusitis associated with the middle turbinate (4,5). A significant correlation between the palatal dimensions, the presence of the concha bullosa and the deviation of the nasal septum that can contribute to the establishment of the skeletal malocclusion is also mentioned (6). In addition, the presence of the concha bullosa and deviation of the nasal septum would be related to compensatory hypertrophy of the contralateral turbinate (7). Most patients with concha bullosa of the middle turbinate are asymptomatic, but cases of blockade of the osteomeatal complex that could lead to chronic sinusitis have begun to be reported, in addition to reports of headaches in patients with pneumatization or the presence of concha bullosa in the superior turbinate (3,8).

Knowledge of anatomical structures and their spatial relationships is essential for diagnosis, as well as treatment planning. With the development and widespread use of new imaging technologies

such as cone beam computed tomography (CBCT) and the analysis tools that it includes, it is possible to observe anatomical structures outside the dentomaxillary complex; being the nasoethmoidal region (nose and ethmoid bone) represented in detail when the field of vision is wide enough (2). In addition, this technique offers multiple advantages such as costs and low radiation dose compared to other imaging techniques, high resolution and definition of details that could not be observed in two-dimensional radiographs. Likewise, it allows observing findings, from anatomical variants, benign lesions to pathologies of importance for the patient's health (9).

The analysis and interpretation of the images by the specialist in Oral and Maxillofacial Radiology must contemplate the description of the anatomical variants, which depending on the case could be the origin of some non-specific symptomatology of the patient. The identification of these will allow complete diagnostic evaluations and adequate treatment plans, leaving the decision to refer to the corresponding specialty to the clinician (9).




Figure 1. Images taken from the archive of the Oral and Maxillofacial Radiology Service of the Teaching Dental Center of the Universidad Peruana Cayetano Heredia.



Figure 2. Images taken from the archive of the Oral and Maxillofacial Radiology Service of the Teaching Dental Center of the Universidad Peruana Cayetano Heredia.




Figure 3. Images taken from the archive of the Oral and Maxillofacial Radiology Service of the Teaching Dental Center of the Universidad Peruana Cayetano Heredia.

REFERENCES

1. Rusu M.C., Stoenescu M.D. Rare Anatomic Variation: Pneumatized Secondary Middle Turbinate. *Curr Health Sci J.* 2020 Jul-Sep; 46 (3): 297-9.
2. von Arx T., Lozanoff S., Bornstein M.M. Extraoral anatomy in CBCT – a literature review. Part 1: Nasoethmoidal region. *Swiss Dent J.* 2019 Oct 14; 129 (10): 804-15.
3. Al-Rawi N.H., Uthman A.T., Abdulhameed E., Al Nuaimi A.S., Seraj Z. Concha bullosa, nasal septal deviation, and their impacts on maxillary sinus volume among Emirati people: A cone-beam computed tomography study. *Imaging Sci Dent.* 2019 Mar; 49 (1): 45-51.
4. Sava C.J., Rusu M.C., Săndulescu M., Dincă D. Vertical and sagittal combinations of concha bullosa media and paradoxical middle turbinate. *Surg Radiol Anat.* 2018 Jul; 40 (7): 847-53.
5. Rusu M.C., Măru N., Sava C.J., Motoc A., Săndulescu M., Dincă D. The sagittal grooves of the middle nasal turbinate determine paradoxical curvatures and bifidities. *Niger J Clin Pract.* 2020 Apr; 23 (4): 464-9.
6. Shetty S.R., Al Bayatti S.W., Al-Rawi N.H., Kamath V., Reddy S., Narasimhan S., Al Kawas S., Madi M., Achalli S., Bhat S. The effect of concha bullosa and nasal septal deviation on palatal dimensions: a cone beam computed tomography study. *BMC Oral Health.* 2021 Nov 23; 21 (1): 607. doi: 10.1186/s12903-021-01974-6. PMID: 34814910; PMCID: PMC8609805.
7. Shetty S., Al Bayatti S., Al-Rawi N., Marei H., Reddy S., Abdelatty H., et al. Analysis of inferior nasal turbinate width and concha bullosa in subjects with nasal septum deviation: a cone beam tomography study. *BMC Oral Health.* 2021; 21 (206). doi: 10.1186/s12903-021-01576-2. PMID: 33894757; PMCID: PMC8066481.
8. Rusu M.C., Măru N., Sava C.J., Săndulescu M., Dincă D. Rare anatomic variation: Giant unilateral concha bullosa superior. *Morphologie.* 2019 Mar; 103 (341): 54-9.
9. Avsever H., Gunduz K., Karakoç O., Akyol M., Orhan K. Incidental findings on cone-beam computed tomographic images: paranasal sinus findings and nasal septum variations. *Oral Radiol.* 2018 Jan; 34 (1): 40-8.

