

© 2020 Universidad Nacional Autónoma de México, Facultad de Estudios Superiores Zaragoza.

This is an Open Access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

TIP Revista Especializada en Ciencias Químico-Biológicas, 23: 1-10, 2020.

<https://doi.org/10.22201/fesz.23958723e.2020.0.270>

The Randle cycle, the precarious link between sugars and fats

Genaro Matus-Ortega^{1**}, Lucero Romero-Aguilar^{1***}, James González²,
Guadalupe Guerra Sánchez³, Maura Matus-Ortega⁴,
Víctor del Castillo-Falconi⁵ and Juan Pablo Pardo^{1*}

¹Universidad Nacional Autónoma de México, Facultad de Medicina, Depto. de Bioquímica; ²Facultad de Ciencias, Depto. de Biología Celular, Av. Universidad # 3000, Copilco, Ciudad Universitaria, Alcaldía de Coyoacán 04510, Ciudad de México, México. ³Instituto Politécnico Nacional, Escuela Nacional de Ciencias Biológicas, Depto. de Microbiología, Ciudad de México, México. ⁴Instituto Nacional de Psiquiatría “Ramón de la Fuente”, Ciudad de México, México. ⁵Universidad Nacional Autónoma de México, Instituto de Investigaciones Biomédicas, Unidad de Investigación en Cáncer, Ciudad Universitaria e Instituto Nacional de Cancerología, Alcaldía de Tlalpan, Ciudad de México, México. E-mails: *pardov@bq.unam.mx, **genaromatus@bq.unam.mx, ***lusromaguila@bq.unam.mx

ABSTRACT

Obesity is a growing global health concern, closely related to cardiovascular diseases. Understanding the correlation between excessive sugar consumption and the formation of fat deposits, described in the Randle cycle, will allow us to have a better grasp on metabolic processes that disrupt the balance between fat formation and degradation processes. The goal of this review is to expand and update the information about the Randle cycle and describe their different levels of regulation. In addition, the participation of mTORC1 and the AMP dependent Kinase (AMPK) during the postprandial and fasting states is described.

Keywords: metabolic syndrome, hepatotoxicity, fructose, lipogenesis, mTORC1, AMPK, dyslipidemia, Randle cycle.

El ciclo de Randle, el precario vínculo entre azúcares y grasas

RESUMEN

La obesidad es un problema de salud global, asociada con enfermedades cardiovasculares. El análisis de la relación existente entre el elevado consumo de glucosa y la concomitante formación de depósitos de grasa, descrita por el ciclo de Randle, permitirá desarrollar una comprensión de los procesos metabólicos involucrados en el balance entre la formación y la degradación de los lípidos. Esta revisión tiene como objetivo, proporcionar una actualización del ciclo y de sus diferentes niveles de regulación, así como la participación de mTORC1 y la cinasa dependiente de AMP (AMPK) durante el estado postprandial y de ayuno.

Palabras clave: síndrome metabólico, hepatotoxicidad, fructosa, lipogénesis, mTORC1, AMPK, dislipidemia, ciclo de Randle.

INTRODUCTION

In accordance with data from the World Health Organization (WHO), diseases associated with obesity have become one of the main health problems worldwide. The number of overweight people in almost every region of the world (except in certain sub-Saharan African regions and some Asian areas) has been increasing at a constant annual rate of 0.7% since 1975 to the end of the second decade of the 21st century (World Health Organization, 2018). Using the body mass index (BMI) scale, the WHO pointed out that in 2016, more than 39% of people older than 18 years old (more than 1,900 million) were overweight, while 13% of the world's population (more than 650 million people) was diagnosed with obesity. Among children and teenagers within the age interval of 5-19 years and children under 5 years old, 18% (over 340 million) and 6% (more than 113 million children) were overweight, respectively (Murray, 2019; Pearlman, Obert & Casey, 2017; Stanhope, 2016). This worldwide phenomenon in which there are more overweight than underweight people was recognized since the last third of the 20th century, indicating that two out of the three countries in North America (namely, México and the United States), and many countries of the European Union, had the most affected population by this health crisis (Hruby & Hu, 2015; Ogden, Yanovski, Carroll & Flegal, 2007; Pereira *et al.*, 2020; Smith & Smith, 2016).

It is generally stated that the main cause of obesity is related to an imbalance between the calories consumed and the calories expended. In accordance with WHO experts (World Health Organization, 2018), obesity problems can be explained considering that *“there is an increased intake of energy-dense foods that are high in fat, along with an increase in physical inactivity due to the increasingly sedentary nature of many forms of work, changing modes of transportation, and increasing urbanization”*.

However, to prevent and treat the obesity problem, experts need to clearly understand lipogenesis and lipolysis, as well as the processes that determine the formation of adipose tissue derived from both sugar-rich foods, whose main ingredient is fructose, and foods high in fat (Moran & Ladenheim, 2016; Priyadarshini & Anuradha, 2017). In other words, it is essential to understand the glucose-fatty acid cycle, also known as the Randle cycle, to recognize the causes of obesity and propose preventive and effective measures (Randle, Garland, Hales & Newsholme, 1963).

Likewise, the general population should be aware of the seventy different names given to sugar that are included in processed foods, in order to keep track of excessive carbohydrate consumption (Gómez Candela & Palma Milla,

2013; Rodríguez Delgado, 2017). It is estimated that sugar-sweetened beverages (soft drinks, juices, nectars, teas, energy drinks, yogurts, among others) are the main sources of sugar in the diet, accounting for more than 15% of the daily caloric intake. Besides, many people do not even realize that their consumption of sugar-sweetened beverages and low-nutrient density foods is much more frequent than they think (Jensen *et al.*, 2018; Rodríguez Delgado, 2017).

This increase in sugar consumption has been associated with pathologies such as liver steatosis, type 2 diabetes mellitus, simple and combined hyperlipidemias (hypertriglyceridemia and hypercholesterolemia), cardiovascular diseases (hypertension, and heart failure) and dental caries, the latter originally described as the only disorder due to sugar consumption. Therefore, in this review we updated the information regarding the Randle cycle, proposed in 1963 (Randle *et al.*, 1963), and the balance between the formation of acylglycerols and their breakdown (lipogenesis/lipolysis).

THE RANDLE CYCLE AND ITS ASSOCIATION WITH THE BALANCE BETWEEN LIPOGENESIS AND LIPOLYSIS

Postprandial state

Under hyperglycemic conditions, such as the postprandial state, insulin induces an increase in the expression of glycolytic regulatory enzymes (glucokinase; phosphofructokinase 1, PFK-1; and pyruvate kinase) and the glucose transporter GLUT 4 (Figure 1). Insulin also activates genes that code for enzymes involved in the Randle cycle (Table I), leading to an increase in the glycolytic and Krebs cycle fluxes and the stimulation of anabolic pathways, such as lipogenesis, β -reduction [synthesis of fatty acids in the cytosol catalyzed by the Fatty Acid Synthase (FAS)], phospholipogenesis and cholesterologenesis (Marcelino *et al.*, 2013; Nakamura, Yudell & Loo, 2014; Palomer, Salvado, Barroso & Vázquez-Carrera, 2013; Possik, Madiraju & Prentki, 2017).

In terms of metabolic pathways, it can be inferred that a sugar overload in glycolysis will drive some of the glucose carbons towards dihydroxyacetone phosphate (DHAP) (Figure 1), which is involved in the formation of acylglycerols (lipogenesis) and phospholipids (phospholipogenesis) (Song, Xiaoli & Yang, 2018) (Figure 1). Therefore, glycolytic flux and anaplerotic pathways are activated in the presence of insulin (Ameer, Scanduzzi, Hasnain, Kalbacher, & Zaidi, 2014; Bartelt *et al.*, 2013; Summermatter *et al.*, 2009).

Carbon overload in glycolysis is also associated with the transfer of citrate from mitochondria to the cytosol, where oxaloacetate (OAA) and acetyl-CoA are produced by the ATP citrate lyase (Figure 1). The first of these metabolites can be reduced or transaminated and returned to the mitochondrial matrix, forming part of the malate-aspartate shuttle.




Table I. Enzymes involved in the Randle cycle. Insulin increases the entry of glucose into the cells, the rate of glycolysis, the pentose phosphate pathway, as well as some anabolic pathways that are fed by the carbon skeletons derived from glucose. Some examples concerning these pathways are β -reduction and lipogenesis. Modified from Nelson & Cox, 2017.

Increased expression	Metabolic pathway
Hexokinase II	Glycolysis
Hexokinase IV	Glycolysis
Phosphofructokinase-1	Glycolysis
Pyruvate kinase	Glycolysis
Phosphofructokinase-2/ Fructose-2,6-bisphosphatase	Glycolysis/gluconeogenesis regulation
Glucose 6-phosphate dehydrogenase	Phosphopentose pathway
6-phosphogluconate dehydrogenase	Phosphopentose pathway
Pyruvate dehydrogenase complex	Krebs cycle entry
Acetyl-CoA carboxylase	β -reduction
Malic enzyme	β -reduction
Citrate lyase cytosolic	β -reduction
Fatty acid synthase	β -reduction
Acyl-CoA-glycerol transferase	Lipogenesis
Decreased expression	Metabolic pathway
Phosphoenolpyruvate carboxykinase	Gluconeogenesis
Glucose 6-phosphatase	Glycemic regulation

2012), such as apoprotein C-II on the smooth and skeletal muscles, and adipose tissue (Figure 1). Also, hydrolysis of TAG is more efficient when apoprotein C-V is active.

Fasting conditions

Under fasting or starvation conditions, lipolysis in white adipocytes is increased by hormones, such as glucagon (Pereira *et al.*, 2020) and norepinephrine, which activate the hormone-sensitive lipase (Figure 1), and decrease the activity of the enzymes that control lipid anabolism, such as HMG-CoA reductase, ACC, and LPL (Hilton, Karpe & Pinnick, 2015; Quiroga & Lehner, 2018; Rambold *et al.*, 2015). Due to their hydrophobic character, free fatty acids exported to the blood plasma are transported by albumin toward the muscle and liver tissues. Uptake of fatty acids into the liver or muscle cells is carried out by Fatty Acid Binding Protein (FABP), Fatty Acid Transporter Protein (FATP), and Fatty Acid Transporter (FAT-CD36) (Figure 1). Intracellular fatty acids are then activated in the form of acyl-CoA in hepatocytes

and muscle cells and subsequently translocated into the mitochondrial matrix by the fatty acid transporter CAT1 and degraded by β -oxidation (Figure 1).

Acetyl-CoA overproduction by β -oxidation of fatty acids causes the allosteric inhibition of the pyruvate dehydrogenase complex (Figure 1). This allows the production of OAA from pyruvate, and thus the beginning of hepatic gluconeogenesis (Figure 1) (Fuchs *et al.*, 2012; Sánchez-Gurmaches *et al.*, 2018). Glycerol obtained from TAG degradation is incorporated at the level of DHAP, feeding the gluconeogenesis in the liver (Figure 1). Glycerol is the most efficient gluconeogenic substrate, compared to alanine, lactate, and other carbon skeletons of some gluconeogenic amino acids (Figure 1). In energy terms, the synthesis of one molecule of glucose from glycerol requires two ATP molecules, instead of six ATP equivalents if gluconeogenesis begins from pyruvate (Fry & Carter, 2019; Pietrocola *et al.*, 2017).

HEPATIC METABOLISM OF FRUCTOSE

Fructose, obtained from fruits and honey, is an intense-flavor sweetener that is added to most processed foods (Bray, 2013; Feinman & Fine, 2013). Fructose presentations include free fructose, sucrose, polysaccharides (fructans) in syrups and nectars, among others (Choo *et al.*, 2018). The rise in fructose consumption has been associated with the increase in obesity and the onset of the metabolic syndrome (Elliott, Keim, Stern, Teff & Havel, 2002; Sievenpiper *et al.*, 2014) (Figure 2). This type of sugar is metabolized largely by hepatocytes, and its assimilation takes place in parallel with the catabolism of other hexoses in glycolysis (Ter Horst & Serlie, 2017). Glut 2 mediates the transport of fructose into the hepatocytes, and the monosaccharide is phosphorylated by fructokinase C, also known as ketohexokinase. Glyceraldehyde and DHAP are produced from fructose 1-phosphate by aldolase B, which allows the integration of fructose into the middle part of glycolysis (Figure 2).

Fructose is a highly lipogenic sugar in comparison with other monosaccharides (Loza-Medrano *et al.*, 2019; Mai & Yan, 2019), because it enters the glycolytic pathway without any allosteric or hormonal control of the fructokinase C. For instance, hexokinases and PFK-1 prevent an accelerated rate of ATP consumption and avoid the overproduction of ADP and trioses that feed lipogenesis (Abdelmalek *et al.*, 2012; Mock, Lateef, Benedito & Tou, 2017).

The increase in the formation of DHAP derived from fructose metabolism, augments the synthesis of fatty acids and the accumulation of triacylglycerol deposits that can progress to steatosis (Figure 2), along with an increase in VLDL and a decrease in High-Density Lipoproteins (HDL) (Ishimoto *et al.*, 2013; Roglans *et al.*, 2007).




Figure 2. Metabolic pathways involved in the assimilation of fructose. Abbreviations: TAG, Triacylglycerol; DAG, Diacylglycerol; MAG, Monoacylglycerol; IMP; Inosine monophosphate; AMP; Adenosine monophosphate. Modified from Nelson & Cox, 2017.

At the molecular level, frequent fructose intake increases the production of mRNAs for FAS and the stearoyl-CoA desaturase 1 (SCD1), which stimulates the synthesis of triacylglycerols and the introduction of the first double bond to the saturated fatty acids, respectively (Basaranoglu, Basaranoglu, Sabuncu & Senturk, 2013). In addition, fructose increases the mRNA of the Carbohydrate-Responsive Element-Binding Proteins (ChREBP) and the mRNA of proteins that participate in the STAT3 pathway involved in the release of leptin (Roglans *et al.*, 2007). It has been stated that ChREBP is a transcription factor that regulates the synthesis of enzymes participating in glycolysis, fructolysis and gluconeogenesis. Also, ChREBP is involved in the *de novo* synthesis of triacylglycerols and cholesterol, regardless of insulin activation (Ter Horst & Serlie, 2017).

Frequent fructose intake is associated with hypertension, insulin resistance, steatosis and hypertriglyceridemia, and causes non-alcoholic fatty liver disease in people with obesity, in which the nuclear receptor PPAR α and its target NF- κ B participate in the decrease of the rate of β -oxidation under gluconeogenesis conditions (Costa Gil & Spinedi, 2017; Laughlin *et al.*, 2014; Roglans *et al.*, 2002). High fructose intake is also related to the onset of gout disease (Figure 1). As a consequence of the increase in fructokinase C activity and the associated high rate of ATP consumption, there is a rise in the concentrations of ADP and AMP that causes a higher production of uric acid and inflammation of some joints (Mai & Yan, 2019). The link between fructose

intake and gout arthritis has been observed in various animal models within minutes after the ingestion of fructose (Jensen *et al.*, 2018).

In addition, the increase of uric acid levels results in the activation of cytosolic NADPH oxidase that translocates to the mitochondria, generating oxidative stress and the inhibition of the aconitase 2, and resulting in the accumulation of citrate in the mitochondrial matrix (Jamnik *et al.*, 2016; Jensen *et al.*, 2018). This causes the export of citrate to the cytoplasm and the stimulation of lipogenesis and cholesterologenesis (Figure 1). The oxidative stress in mitochondria spreads to the endoplasmic reticulum, activating the Sterol Regulatory Element-Binding transcription factor 1 (SREBP-1), which in turn increases the transcript levels of genes involved in lipogenesis and cholesterol synthesis (Jensen *et al.*, 2018; Lustig, 2010; Samuel, 2011) (Figure 1).

CONTROL OF RANDLE CYCLE BY mTORC1 AND AMPK

The mammalian target of rapamycin (mTOR) is a kinase that forms two complexes in mammals: mTORC1 and mTORC2. mTORC1 is activated by amino acids (Chen, Wei, Liu & Guan, 2014; Cheng & Saltiel, 2006), growth factors and hormones, such as insulin (Baena *et al.*, 2015; Verges, 2018). mTORC2 is also regulated by growth factors and is involved in cytoskeleton remodeling and sphingolipid synthesis (Figure 3). During the postprandial state, insulin stimulates phosphoinositide-dependent kinase 1 (PDK1),




Figure 3. Functional relationships between mTORC1 and the AMP activated protein kinase (AMPK) in the Randle cycle. Abbreviations: AMPK: AMP-Activated Protein Kinase; mTORC1: mammalian Target of Rapamycin Complex 1; PIP2: phosphatidylinositol (4,5)-bisphosphate; PI3K: phosphoinositide 3 kinase; PIP3: phosphatidylinositol (3,4,5)-trisphosphate; PDK1: 3-phosphoinositide-dependent kinase-1; PKB/Akt: protein kinase B/Akt; TSC1-TSC2: 1-2 tuberous sclerosis complex (or hamartin-tuberin complex); PKA: Protein Kinase A; LKB1: Liver Kinase B1. Modified from Yoon, 2017.

which leads to the activation of PKB/Akt signaling pathway, inhibition of the TSC1/TSC2 complex (tuberous sclerosis complex 1 and 2), and activation of mTORC1, which promotes lipogenesis, glycolysis, and glycogen synthesis (Asati, Mahapatra & Bharti, 2016; Jiang *et al.*, 2008; Kumar *et al.*, 2010; Naito, Kuma & Mizushima, 2013; Verges, 2018). On the contrary, the AMP-dependent Kinase (AMPK) is hormonally downregulated under the hyperglycemia status and activated during fasting or exercise conditions. Activation of the AMPK depends on the stimulation of both the AMPc-dependent protein kinase (PKA) and the human tumor suppressor liver kinase 1 (LKB1), and the increase in the concentration of AMP (Kim & He, 2013). Along with the stimulation of PKA and AMPK there is a decrease in the main lipogenic pathways, such as fatty acid synthesis, triacylglycerol accumulation and cholesterologenesis, and activation of gluconeogenesis (Hasenour *et al.*, 2017), glycogen degradation, lipolysis and mitochondrial β -oxidation, thereby increasing ketogenesis in the liver (Cardaci, Filomeni, & Ciriolo, 2012). AMPK, through the

phosphorylation of ACC and HMG-CoA reductase, inhibits the synthesis of fatty acids and cholesterol, respectively.

In short, triacylglycerol accumulation in white fat deposits, liver tissue, and between fiber bundles is caused by hypercaloric diets rich in fast-digesting carbohydrates, along with the sedentary lifestyle habits of Western societies (Perera & Turner, 2016). Hypertriglyceridemia and hypercholesterolemia are involved in the pathophysiology of health problems, such as high blood pressure, diabetes mellitus 2, atherosclerosis and obesity, among other diseases (Ke, Xu, Li, Luo & Huang, 2018; Nakamura *et al.*, 2014; Palomer *et al.*, 2013; Possik *et al.*, 2017).

CONCLUSIONS

There is a metabolic relationship between sugar consumption and fat accumulation. In the specific case of fructose, the excessive consumption of this sugar causes depletion of cellular ATP, steatosis, obesity, metabolic syndrome, and an increase in the production of uric acid. These adverse metabolic effects

are the consequence of the lack of regulatory mechanisms for the incorporation of fructose into the glycolytic pathway. A new addition to the Randle cycle is the incorporation of mTORC1 and the antagonistic effect of the AMPK to ensure an efficient regulation of lipogenesis and lipolysis, respectively. In terms of public policy, authorities of health institutes should advise against the abuse of carbohydrate consumption.

CONFLICTING INTERESTS

The authors declare that there is no conflict of interest.

ACKNOWLEDGMENTS

This work was supported by the Universidad Nacional Autónoma de México (UNAM), Programa de Apoyo a Proyectos de Investigación Tecnológica [PAPIIT IN222117]; Consejo Nacional de Ciencia y Tecnología, CONACYT [254904-JPP] and [256520-GGS]. Instituto Politécnico Nacional- Secretaría de Investigación y Posgrado, [IPN-SIP 20190200]. We are grateful to Oscar Iván Luqueño Bocardo for the design of Figure 1.

REFERENCES

- Abdelmalek, M. F., Lazo, M., Horska, A., Bonekamp, S., Lipkin, E. W., Balasubramanyam, A., Bantle, J. P., Johnson, R. J., Diehl, A. M. & Clark, J. M. Fatty Liver Subgroup of Look AHEAD Research Group. (2012). Higher dietary fructose is associated with impaired hepatic adenosine triphosphate homeostasis in obese individuals with type 2 diabetes. *Hepatology*, **56**(3), 952-960. DOI: 10.1002/hep.25741
- Aguiar, L. R. , Pardo, J. P., Lomelí, M. M., Bocardo, O. I. L., Juárez Oropeza, M. A. & Guerra Sánchez, G. (2017). Lipid droplets accumulation and other biochemical changes induced in the fungal pathogen *Ustilago maydis* under nitrogen-starvation. *Arch. Microbiol.*, **199**(8):1195-1209. DOI: 10.1007/s00203-017-1388-8
- Ameer, F., Scanduzzi, L., Hasnain, S., Kalbacher, H. & Zaidi, N. (2014). De novo lipogenesis in health and disease. *Metabolism*, **63** (7), 895-902. DOI: 10.1016/j.metabol.2014.04.003
- Asati, V., Mahapatra, D. K. & Bharti, S. K. (2016). PI3K/Akt/mTOR and Ras/Raf/MEK/ERK signaling pathways inhibitors as anticancer agents: Structural and pharmacological perspectives. *Eur. J. Med. Chem.*, **109**, 314-341. DOI: 10.1016/j.ejmech.2016.01.012
- Baena, M., Sanguesa, G., Hutter, N., Sánchez, R. M., Roglans, N., Laguna, J. C. & Alegret, M. (2015). Fructose supplementation impairs rat liver autophagy through mTORC activation without inducing endoplasmic reticulum stress. *Biochim. Biophys. Acta*, **1851**(2), 107-116. DOI: 10.1016/j.bbali.2014.11.003
- Barbosa, A. D. & Siniosoglou, S. (2017). Function of lipid droplet-organelle interactions in lipid homeostasis. *Biochim. Biophys. Acta Mol. Cell Res.*, **1864**(9), 1459-1468. DOI: 10.1016/j.bbamcr.2017.04.001
- Bartelt, A., Weigelt, C., Cherradi, M. L., Niemeier, A., Todter, K., Heeren, J. & Scheja, L. (2013). Effects of adipocyte lipoprotein lipase on de novo lipogenesis and white adipose tissue browning. *Biochim Biophys Acta*, **1831**(5), 934-942. DOI: 10.1016/j.bbali.2012.11.011
- Basaranoglu, M., Basaranoglu, G., Sabuncu, T. & Senturk, H. (2013). Fructose as a key player in the development of fatty liver disease. *World J. Gastroenterol.*, **19**(8), 1166-1172. DOI: 10.3748/wjg.v19.i8.1166
- Bray, G. A. (2013). Energy and fructose from beverages sweetened with sugar or high-fructose corn syrup pose a health risk for some people. *Adv. Nutr.*, **4**(2), 220-225. DOI: 10.3945/an.112.002816
- Cardaci, S., Filomeni, G. & Ciriolo, M. R. (2012). Redox implications of AMPK-mediated signal transduction beyond energetic clues. *J. Cell Sci.*, **125**(Pt 9), 2115-2125. DOI: 10.1242/jcs.095216
- Chen, Y., Wei, H., Liu, F. & Guan, J. L. (2014). Hyperactivation of mammalian target of rapamycin complex 1 (mTORC1) promotes breast cancer progression through enhancing glucose starvation-induced autophagy and Akt signaling. *J. Biol. Chem.*, **289**(2), 1164-1173. DOI: 10.1074/jbc.M113.526335
- Cheng, A. & Saltiel, A. R. (2006). More TORC for the gluconeogenic engine. *Bioessays*, **28**(3), 231-234. DOI: 10.1002/bies.20375
- Choo, V. L., Vigiouliouk, E., Blanco Mejia, S., Cozma, A. I., Khan, T.A., Ha, V., Wolever, T. M. S., Leiter, L. A., Vuksan, V., Kendall, C. W. C., de Souza, R. J., Jenkins, D. J. A. & Sievenpiper, J. L. (2018). Food sources of fructose-containing sugars and glycaemic control: systematic review and meta-analysis of controlled intervention studies. *BMJ*, **363**, k4644. DOI: 10.1136/bmj.k4644
- Costa Gil, J. E. & Spinedi, E. (2017). La tormentosa relación entre las grasas y el desarrollo de la diabetes mellitus de tipo 2: actualizado. Parte I. *Revista Argentina de Endocrinología y Metabolismo*, **54**, 109-123. DOI: 10.1016/j.raem.2017.06.001
- Elliott, S. S., Keim, N. L., Stern, J. S., Teff, K. & Havel, P. J. (2002). Fructose, weight gain, and the insulin resistance syndrome. *Am. J. Clin. Nutr.*, **76**(5), 911-922. DOI: 10.1093/ajcn/76.5.911
- Feinman, R. D. & Fine, E. J. (2013). Fructose in perspective. *Nutr. Metab. (Lond.)*, **10**(1), 45. DOI: 10.1186/1743-7075-10-45
- Fry, B. & Carter, J. F. (2019). Stable carbon isotope diagnostics of mammalian metabolism, a high-resolution isotomics approach using amino acid carboxyl groups. *PLoS One*, **14**(10), e0224297. DOI: 10.1371/journal.pone.0224297
- Fuchs, C. D., Claudel, T., Kumari, P., Haemmerle, G., Pollheimer, M. J., Stojakovic, T., Schrnagl, H.,

- Halilbasic, E., Gumhold, J., Silbert, D., Koefeler, H. & Trauner, M. (2012). Absence of adipose triglyceride lipase protects from hepatic endoplasmic reticulum stress in mice. *Hepatology*, **56**(1), 270-280. DOI: 10.1002/hep.25601
- Gómez Candela, C. & Palma Milla, S. (2013). Una visión global, actualizada y crítica del papel del azúcar en nuestra alimentación. *Nutrición Hospitalaria*, **28**, 1-4.
- Hasenour, C. M., Ridley, D. E., James, F. D., Hughey, C. C., Donahue, E. P., Viollet, B., Foretz, M., Young, J. D. & Wasserman, D. H. (2017). Liver AMP-Activated Protein Kinase Is Unnecessary for Gluconeogenesis but Protects Energy State during Nutrient Deprivation. *PLoS One*, **12**(1), e0170382. DOI: 10.1371/journal.pone.0170382
- Hilton, C., Karpe, F. & Pinnick, K. E. (2015). Role of developmental transcription factors in white, brown and beige adipose tissues. *Biochim. Biophys. Acta*, **1851**(5), 686-696. DOI: 10.1016/j.bbali.2015.02.003
- Hruby, A. & Hu, F. B. (2015). The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*, **33**(7), 673-689. DOI: 10.1007/s40273-014-0243-x
- Ishimoto, T., Lanaspá, M. A., Rivard, C. J., Roncal-Jimenez C. A., Orlicky, D. J., Cicerchi, C., McMahan, R. H., Abdelmalek, M. F., Rosen, H. R., Jackman, M. R., MacLean, P. S., Diggle, C. P., Asipu, A., Inaba, S., Kosugi, T., Sato, W., Maruyama, S., Sánchez-Lozada, L. G., Sautin, Y.Y., Hill, J. O., Bonthron, D. T. & Johnson, R. J. (2013). High-fat and high-sucrose (western) diet induces steatohepatitis that is dependent on fructokinase. *Hepatology*, **58**(5), 1632-1643. DOI: 10.1002/hep.26594
- Jamnik, J., Rehman, S., Blanco Mejia, S., de Souza, R. J., Khan, T.A., Leiter, L.A., Wolever, T. M., Kendall, C. W., Jenkins, D. J. & Sievenpiper, J. L. (2016). Fructose intake and risk of gout and hyperuricemia: a systematic review and meta-analysis of prospective cohort studies. *BMJ Open*, **6**(10), e013191. DOI: 10.1136/bmjopen-2016-013191
- Jensen, T., Abdelmalek, M. F., Sullivan, S., Nadeau, K. J., Green, M., Roncal, C., Nakagawa, T., Kuwabara, M., Sato, Y., Kang, D. H., Tolan, D. R., Sanchez-Lozada, L. G., Rosen, H. R., Lanaspá, M. A., Diehl, A. M. & Johnson, R. J. (2018). Fructose and sugar: A major mediator of non-alcoholic fatty liver disease. *J. Hepatol.*, **68**(5), 1063-1075. DOI: 10.1016/j.jhep.2018.01.019
- Jiang, X., Kenerson, H., Aicher, L., Miyaoka, R., Eary, J., Bissler, J. & Yeung, R. S. (2008). The tuberous sclerosis complex regulates trafficking of glucose transporters and glucose uptake. *Am. J. Pathol.*, **172**(6), 1748-1756. DOI: 10.2353/ajpath.2008.070958
- Ke, R., Xu, Q., Li, C., Luo, L. & Huang, D. (2018). Mechanisms of AMPK in the maintenance of ATP balance during energy metabolism. *Cell Biol. Int.*, **42**(4), 384-392. DOI: 10.1002/cbin.10915
- Kim, I. & He, Y. Y. (2013). Targeting the AMP-Activated Protein Kinase for Cancer Prevention and Therapy. *Front. Oncol.*, **3**, 175. DOI: 10.3389/fonc.2013.00175
- Kory, N., Farese, R. V., Jr. & Walther, T. C. (2016). Targeting Fat: Mechanisms of Protein Localization to Lipid Droplets. *Trends Cell Biol.*, **26**(7), 535-546. DOI: 10.1016/j.tcb.2016.02.007
- Kumar, A., Lawrence, J. C. Jr., Jung, D.Y., Ko, H. J., Keller, S. R., Kim, J. K., Magnuson, M. A. & Harris, T. E. (2010). Fat cell-specific ablation of rictor in mice impairs insulin-regulated fat cell and whole-body glucose and lipid metabolism. *Diabetes*, **59**(6), 1397-1406. DOI: 10.2337/db09-1061
- Laughlin, M. R., Bantle, J. P., Havel, P. J., Parks, E., Klurfeld, D. M., Teff, K. & Maruvada, P. (2014). Clinical research strategies for fructose metabolism. *Adv. Nutr.*, **5**(3), 248-259. DOI: 10.3945/an.113.005249
- Loza-Medrano, S. S., Baiza-Gutman, L. A., Manuel-Apolinar, L., García-Macedo, R., Damasio-Santana, L., Martínez-Mar, O. A., Sánchez-Becerra, M. C., Cruz-López, M., Ibáñez-Hernández, M. A. & Díaz-Flores, M. (2019). High fructose-containing drinking water-induced steatohepatitis in rats is prevented by the nicotinamide-mediated modulation of redox homeostasis and NADPH-producing enzymes. *Mol. Biol. Rep.*, **47**(1), 337-351. DOI: 10.1007/s11033-019-05136-4
- Lustig, R. H. (2010). Fructose: metabolic, hedonic, and societal parallels with ethanol. *J. Am. Diet. Assoc.*, **110**(9), 1307-1321. DOI: 10.1016/j.jada.2010.06.008
- Mai, B. H. & Yan, L. J. (2019). The negative and detrimental effects of high fructose on the liver, with special reference to metabolic disorders. *Diabetes Metab. Syndr. Obes.*, **12**, 821-826. DOI: 10.2147/DMSO.S198968
- Marcelino, H., Veyrat-Durebex, C., Summermatter, S., Sarafian, D., Miles-Chan, J., Arsenijevic, D., Zani, F., Montani, J. P., Seydoux, J., Solinas, G., Rohner-Jeanrenaud, F. & Dulloo, A. G. (2013). A role for adipose tissue de novo lipogenesis in glucose homeostasis during catch-up growth: a Randle cycle favoring fat storage. *Diabetes*, **62**(2), 362-372. DOI: 10.2337/db12-0255
- Mock, K., Lateef, S., Benedito, V. A. & Tou, J. C. (2017). High-fructose corn syrup-55 consumption alters hepatic lipid metabolism and promotes triglyceride accumulation. *J. Nutr. Biochem.*, **39**, 32-39. DOI: 10.1016/j.jnutbio.2016.09.010
- Moran, T. H. & Ladenheim, E. E. (2016). Physiologic and Neural Controls of Eating. *Gastroenterol. Clin. North Am.*, **45**(4), 581-599. DOI: 10.1016/j.gtc.2016.07.009
- Mottillo, E. P., Balasubramanian, P., Lee, Y. H., Weng, C., Kershaw, E. E. & Granneman, J. G. (2014). Coupling of lipolysis and de novo lipogenesis in brown, beige, and white adipose tissues during chronic beta3-adrenergic receptor activation. *J. Lipid. Res.*, **55**(11), 2276-2286. DOI: 10.1194/jlr.M050005
- Murray, R. D. (2019). 100% Fruit Juice in Child and Adolescent Dietary Patterns. *J. Am. Coll. Nutr.*, **39**(2),

- 122-127. DOI: 10.1080/07315724.2019.1615013
- Naito, T., Kuma, A. & Mizushima, N. (2013). Differential contribution of insulin and amino acids to the mTORC1-autophagy pathway in the liver and muscle. *J. Biol. Chem.*, **288**(29), 21074-21081. DOI: 10.1074/jbc.M113.456228
- Nakamura, M. T., Yudell, B. E. & Loor, J. J. (2014). Regulation of energy metabolism by long-chain fatty acids. *Prog. Lipid. Res.*, **53**, 124-144. DOI: 10.1016/j.plipres.2013.12.001
- Nelson, D. L. & Cox, M. (2017). *Lehninger principles of biochemistry*. W.H. Freeman . New York.
- Ogden, C. L., Yanovski, S. Z., Carroll, M. D. & Flegal, K. M. (2007). The epidemiology of obesity. *Gastroenterology*, **132**(6), 2087-2102. DOI: 10.1053/j.gastro.2007.03.052
- Palomer, X., Salvado, L., Barroso, E. & Vazquez-Carrera, M. (2013). An overview of the crosstalk between inflammatory processes and metabolic dysregulation during diabetic cardiomyopathy. *Int. J. Cardiol.*, **168**(4), 3160-3172. DOI: 10.1016/j.ijcard.2013.07.150
- Pearlman, M., Obert, J. & Casey, L. (2017). The Association Between Artificial Sweeteners and Obesity. *Curr. Gastroenterol. Rep.*, **19**(12), 64. DOI: 10.1007/s11894-017-0602-9
- Pereira, M. J., Thombare, K., Sarsenbayeva, A., Kamble, P. G., Almby, K., Lundqvist, M. & Eriksson, J. W. (2020). Direct effects of glucagon on glucose uptake and lipolysis in human adipocytes. *Mol. Cell Endocrinol.*, **503**, 110696. DOI: 10.1016/j.mce.2019.110696
- Perera, N. D. & Turner, B. J. (2016). AMPK Signalling and Defective Energy Metabolism in Amyotrophic Lateral Sclerosis. *Neurochemical Research*, **41**(3), 544-553. DOI: 10.1007/s11064-015-1665-3
- Pietrocola, F., Demont, Y., Castoldi, F., Enot, D., Durand, S., Semeraro, M., Baracco, E. E., Pol, J., Bravo-San Pedro, J. M., Bordenave, C., Levesque, S., Humeau, J., Chery, A., Métivier, D., Madeo, F., Maiuri, M. C. & Kroemer, G. (2017). Metabolic effects of fasting on human and mouse blood *in vivo*. *Autophagy*, **13**(3), 567-578. DOI: 10.1080/15548627.2016.1271513
- Possik, E., Madiraju, S. R. M. & Prentki, M. (2017). Glycerol-3-phosphate phosphatase/PGP: Role in intermediary metabolism and target for cardiometabolic diseases. *Biochimie*, **143**, 18-28. DOI: 10.1016/j.biochi.2017.08.001
- Priyadarshini, E. & Anuradha, C. V. (2017). Glucocorticoid Antagonism Reduces Insulin Resistance and Associated Lipid Abnormalities in High-Fructose-Fed Mice. *Can. J. Diabetes*, **41**(1), 41-51. DOI: 10.1016/j.cjcd.2016.06.003
- Quiroga, A. D. & Lehner, R. (2012). Liver triacylglycerol lipases. *Biochim. Biophys. Acta*, **1821**(5), 762-769. DOI: 10.1016/j.bbalip.2011.09.007
- Quiroga, A. D. & Lehner, R. (2018). Pharmacological intervention of liver triacylglycerol lipolysis: The good, the bad and the ugly. *Biochem. Pharmacol.*, 155, 233-241. DOI: 10.1016/j.bcp.2018.07.005
- Rambold, A. S., Cohen, S. & Lippincott-Schwartz, J. (2015). Fatty acid trafficking in starved cells: regulation by lipid droplet lipolysis, autophagy, and mitochondrial fusion dynamics. *Dev. Cell*, **32**(6), 678-692. DOI: 10.1016/j.devcel.2015.01.029
- Randle, P. J., Garland, P. B., Hales, C. N. & Newsholme, E. A. (1963). The glucose fatty-acid cycle. Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet*, **1**(7285), 785-789. DOI: 10.1016/s0140-6736(63)91500-9
- Rodríguez Delgado, J. (2017). Azúcares... ¿los malos de la dieta? *Pediatría Atención Primaria*, **19**, 69-75.
- Roglans, N., Sanguino, E., Peris, C., Alegret, M., Vázquez, M., Adzet, T., Díaz, C., Hernández, G., Laguna, J. C. & Sánchez, R. M. (2002). Atorvastatin treatment induced peroxisome proliferator-activated receptor alpha expression and decreased plasma nonesterified fatty acids and liver triglyceride in fructose-fed rats. *J. Pharmacol. Exp. Ther.*, **302**(1), 232-239. DOI: 10.1124/jpet.302.1.232
- Roglans, N., Vila, L., Farre, M., Alegret, M., Sánchez, R. M., Vázquez-Carrera, M. & Laguna, J. C. (2007). Impairment of hepatic Stat-3 activation and reduction of PPARalpha activity in fructose-fed rats. *Hepatology*, **45**(3), 778-788. DOI: 10.1002/hep.21499
- Samuel, V. T. (2011). Fructose induced lipogenesis: from sugar to fat to insulin resistance. *Trends Endocrinol. Metab.*, **22**(2), 60-65. DOI: 10.1016/j.tem.2010.10.003
- Sánchez-Gurmaches, J., Tang, Y., Jespersen, N. Z., Wallace, M., Martinez Calejman, C., Gujja, S., Li, H., Edwards, Y. J. K., Wolfrum, C., Metallo, C. M., Nielsen, S., Scheele, C. & Guertin, D. A. (2018). Brown Fat AKT2 Is a Cold-Induced Kinase that Stimulates ChREBP-Mediated De Novo Lipogenesis to Optimize Fuel Storage and Thermogenesis. *Cell Metab.*, **27**(1), 195-209 e196. DOI: 10.1016/j.cmet.2017.10.008
- Sievenpiper, J. L., de Souza, R. J., Cozma, A. I., Chiavaroli, L., Ha, V. & Mirrahimi, A. (2014). Fructose vs. glucose and metabolism: do the metabolic differences matter? *Curr. Opin. Lipidol.*, **25**(1), 8-19. DOI: 10.1097/MOL.0000000000000042
- Smith, K. B. & Smith, M. S. (2016). Obesity Statistics. *Primare*, **43**(1), 121-135, ix. DOI: 10.1016/j.pop.2015.10.001
- Song, Z., Xiaoli, A. M. & Yang, F. (2018). Regulation and Metabolic Significance of De Novo Lipogenesis in Adipose Tissues. *Nutrients*, **10**(10). DOI: 10.3390/nu10101383
- Stanhope, K. L. (2016). Sugar consumption, metabolic disease and obesity: The state of the controversy. *Crit. Rev. Clin. Lab. Sci.*, **53**(1), 52-67. DOI: 10.3109/10408363.2015.1084990
- Summermatter, S., Marcelino, H., Arsenijevic, D., Buchala, A., Aprikian, O., Assimacopoulos-Jeannet, F., Seydoux, J., Montani, J. P., Solinas, G. & Dulloo, A. G. (2009).

- Diabetes*, **58(10)**, 2228-2237. DOI: 10.2337/db08-1793
- Ter Horst, K. W. & Serlie, M. J. (2017). Fructose Consumption, Lipogenesis, and Non-Alcoholic Fatty Liver Disease. *Nutrients*, **9(9)**. DOI: 10.3390/nu9090981
- Verges, B. (2018). mTOR and Cardiovascular Diseases: Diabetes Mellitus. *Transplantation*, **102(2S Suppl 1)**, S47-S49. DOI: 10.1097/TP.0000000000001722
- Yoon, M-S. (2017). The Role of Mammalian Target of Rapamycin (mTOR) in Insulin Signaling. *Nutrients* 2017, **9**, 1176. DOI:10.3390/nu9111176
- World Health Organization. World Health Statistics (2018): Monitoring Health for the SDGs., 2018.