

Postconditioning to protect the human heart

Dr. Pastor Luna-Ortiz,* Dr. Gustavo Pastelin,* Dr. Juan Carlos Torres,* Dr. Martín Martínez-Rosas**

* Department of Pharmacology.

** Department of Physiology.

Instituto Nacional de Cardiología
«Ignacio Chávez», México, D.F.

Reprints requests:

Dr. Martín Martínez Rosas
Department of Physiology
Instituto Nacional de Cardiología
«Ignacio Chávez», Juan Badiano Núm. 1
Col. Sección 16
14080 México, D.F.
E-mail: martin5163@hotmail.com

Recibido para publicación: 28-09-10.

Aceptado para publicación: 21-10-10.

Este artículo puede ser consultado en versión completa en
<http://www.medigraphic.com/rma>

SUMMARY

Recently, it has been shown that the heart can be protected against the ischemia reperfusion injury if brief coronary occlusions are performed just at the beginning of the reperfusion. This procedure has been called postconditioning (PostC). It can also be elicited by pharmacological interventions, which are named pharmacological PostC. In general, PostC reduces the reperfusion-induced injury, blunts oxidant-mediated damages and attenuates the local inflammatory response to reperfusion, decrease infarct size, diminishes apoptosis, neutrophil activation and endothelial dysfunction. The mechanisms that participate in PostC still are not completely understood. In this regard, adenosine, glycine, bradykinin, ciclosporin A, are involved in post C triggering. Similar to preconditioning, PostC triggers several signaling pathways and molecular components including nitric oxide (NO), Protein Kinase C (PKC), adenosine triphosphate-sensitive potassium channels (IKATP), the pathway: Reperfusion Injury Salvage Kinases (RISK) which compresses to: phosphatidylinositol-3-OH kinase (PI3K) and Extracellular signal-Regulated Kinase (ERK 1/2), and finally, the Survivor Activating Factor Enhancement (SAFE) pathway. In this review we describe the mechanisms of reperfusion-induced injury as well as the proposed protective pathways activated by PostC, which appear to converge in inhibition of mitochondrial permeability transition pores (mPTP) opening. On the other hand, experimental evidence indicated that volatile anesthetics and opioids are capable of exerting cardioprotective effects under certain conditions, constituting a very useful pharmacological Post C. In conclusion, the first minutes of reperfusion represent a window of opportunity for triggering the aforementioned mediators, which acting in concert lead to protection against reperfusion injury (PostC). Pharmacological, especially anesthetic, PostC may have a promising future in the clinical scenarios.

Key words: Postconditioning, cardioprotection and ischemia-reperfusion injury.

RESUMEN

Recientemente, se ha demostrado que el corazón puede protegerse contra el daño por isquemia reperusión si se aplican breves oclusiones coronarias justo al inicio de la reperusión. Este procedimiento ha sido llamado postcondicionamiento (PostC) y puede ser producido por intervenciones farmacológicas, las cuales constituyen el PostC farmacológico. En general, el PosC induce el daño inducido por la reperusión, disminuyendo el daño oxidativo y atenuando la respuesta inflamatoria local durante la reperusión, así también disminuye el tamaño del infarto, disminuye el proceso de apoptosis, la activación neurotrófica y la disfunción endotelial. Los mecanismos que participan en el PostC aún no son bien entendidos. En este sentido, se sabe que moléculas como la adenosina, la glicina, la bradicinina y la ciclosporina A están involucradas en la activación de PostC. De manera similar al preconditionamiento, el PsotC dispara diversas rutas de señalización en las cuales participan diversos componentes moleculares como el óxido nítrico (NO), la proteína cinasa de fosfatidilinositol-3-OH y a la cinasa

reguladora por señales extracelulares (ERK $\frac{1}{2}$), y finalmente la ruta de aumento del factor activación de sobrevivencia (SAFE). En esta revisión describimos los mecanismos de daño inducido por la reperfusión así como las vías protectoras propuestas activadas por el PostC, las cuales parecen converger en una inhibición de la apertura de los poros de transición de la permeabilidad mitocondrial (mPTP). Por otro lado, la evidencia experimental indica que los anestésicos volátiles y los opiáceos son capaces de ejercer efectos cardioprotectores bajo ciertas condiciones, constituyendo un PostC farmacológico muy útil, de esta manera los primeros minutos de la reperfusión representan una ventana de oportunidad para activar los mediadores antes mencionados, los cuales actúan en concierto para llevar a la protección del miocardio contra el daño por reperfusión. El PostC farmacológico especialmente el anestésico puede tener un futuro promisorio en los escenarios clínicos de las salas de operaciones.

Palabras clave: Postcondicionamiento, cardioprotección, daño por isquemia, reperfusión.

INTRODUCTION

Despite advances in prevention and treatment, cardiovascular disease remains as the number one cause of death in men and women in the United States⁽¹⁾. Acute myocardial infarction (AMI) is still a frequent and disabling disease being the infarct size a major determinant of myocardial functional recovery and mortality after an AMI⁽²⁾. Thus, limitation of infarct size represents an appropriate strategy with which to prevent postinfarction heart failure and to improve survival. In the last 15 years there has been a significant improvement in the outcome of AMI patients. This improvement is mainly based on the introduction of efficient reperfusion strategies, such as primary percutaneous coronary intervention (PCI) and thrombolysis⁽³⁾. However, coronary heart disease still presents a mortality rate of around 10% at one year for all AMI patients⁽¹⁾. This remaining high mortality rate is partly a consequence of the phenomenon called «myocardial reperfusion injury»⁽⁴⁾, originally described by Jennings and Sommers⁽⁵⁾ in the sixties on experimental model of myocardial infarction. In the myocardial reperfusion injury, the reperfusion causes additional functional and structural damage to myocardium after a period of ischemia when is acutely reperfused⁽⁵⁾. Studies of AMI in animal models suggest that lethal reperfusion injury, which starts immediately after the opening of the culprit coronary artery, accounts for up to 50% of the final size of a myocardial infarct⁽⁴⁾. In human clinical studies, this phenomenon is associated with detrimental myocardial remodeling, ventricular arrhythmias and no-reflow, and predicts a negative prognosis in myocardial infarction patients⁽⁶⁾. Thus, lethal reperfusion injury has become a major therapeutic target in the search for improvement in AMI patient's recovery and numerous strategies have been tested in experimental settings to reduce reperfusion injury. Recently, it has been shown that the heart can be effectively protected from reperfusion injury not only with ischemic preconditioning (IPC), which is a classic strategy of cardioprotection, but also with brief episodes of

ischemia during the early reperfusion period after a prolonged period of ischemia. This process is called: ischemic postC.

In this review we describe the mechanisms of reperfusion injury, the possible protective mechanisms of the postC and pharmacological PostC, specially using anesthetics to induce the heart protection.

MECHANISMS OF REPERFUSION INJURY

The heart is a pump that converts chemical energy into mechanical work. The power produced by cardiac muscle is generated almost entirely by the oxidation of carbon fuels with oxygen. The fuels are provided to a great extent by coronary blood flow. Oxidative metabolism is the function of mitochondria within the cell, and thus the bulk of cardiac energy is supplied by oxidative phosphorylation within cardiac mitochondria. Like many cells, when deprived of oxygen (anoxia), cardiac cells can maintain ATP levels by anaerobic glycolysis, and can then revert smoothly to oxidative metabolism on reperfusion⁷. However, if blood flow is restricted, as in myocardial infarct, the cells accumulate glycolytic by-products (lactate, H⁺) in addition to suffering from oxygen deprivation⁽⁸⁾. This is a condition known as ischemia and can damage cardiac cells irreversibly. Paradoxically, however, the major damage to ischemic cells comes on the re-introduction of oxygen (reperfusion). During reperfusion, the cells typically undergo further contraction (hypercontracture) and membrane damage, followed by cell death^(9,10).

The reperfusion injury of the myocardium is a complex phenomenon that encompasses a list of events including: reperfusion arrhythmias, microvascular damage, reversible myocardial mechanical dysfunction («stunned» myocardium), and cell death (due to apoptosis or necrosis). These events may occur together or separately⁽²⁷⁾. Most of the detrimental effects of reperfusion are triggered within the first minutes following the re-opening of the occluded coronary artery⁽¹⁰⁾. However, most of the cellular disturbances that occur at the time of re-

perfusion are determined by the abnormalities induced during the ischemic period (Figure 1). During ischemia, the increase in anaerobic glycolysis results in the progressive accumulation of protons and lactic acid, with the acidosis increased eventually inhibiting glycolytic flux and the synthesis of ATP. As the cardiomyocyte attempts to correct acidosis through the Na^+/H^+ exchanger (this transporter moves H^+ out cell and Na^+ into the cell), the Na^+ is accumulated and this event activates the Na^+/K^+ ATPase to pump Na^+ out of the cell. However, taking in account the diminished ATP production, the Na^+/K^+ ATPase, progressively fails without sufficient




Figure 1. Schematic representation of the main events contributing to rapid lethal cardiomyocyte injury during ischemia and hypercontracture-cell death.

ATP. Subsequent activation of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger, in its reverse mode, helps to move Na^+ out of the cell, but favors cytosolic accumulation of Ca^{2+} . Prolonged ischemia induces a progressive failure of ionic homeostasis, which ultimately causes accumulation of intracellular Na^+ and Ca^{2+} and a decline in ATP levels. If ischemia continues for a long time, a hypercontracture is developed and cell death occurs.

During early reperfusion (in the first minutes), occurs a rapid correction of acidosis (low intracellular pH) through the Na^+/H^+ exchanger, and through the entry of ion HCO_3^- by Na/HCO_3 symporter (Figure 2). Additionally the lactate previously accumulated, is washout with the restored blood flow diminishing the acidosis. However, all these events cause accumulation of Na^+ . This additional augment in intracellular sodium induces a secondary activation of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger in the reverse mode to move out ions Na^+ , however this aggravates cytosolic Ca^{2+} accumulation. On the other hand, the abrupt re-exposure of the ischemia-inhibited mitochondrial respiratory chain to oxygen generates a membrane potential to drive ATP synthesis, which leads to a rapid overload of Ca^{2+} in the matrix and massive production of reactive oxygen species (ROS) which by themselves are capable to damage cellular membranes and induce oxidative stress. These two factors (cytosolic Ca^{2+} accumulation and oxidative stress) trigger the mPTP to open, and this pore is a key element in cell death⁽¹¹⁾. The mPTP is a nonspecific channel located on the inner mitochondrial membrane. The mPTP was originally proposed to consist of 3 major components: a voltage-dependent anion channel, the adenine nucleotide translocase, and cyclophilin D, contained within the mitochondrial matrix, however, its structure is not completely defined. Under normal physiological conditions, the mitochondrial inner membrane is impermeable to almost all metabolites and ions, and the mPTP is in a closed conformation. Under some stress conditions, the mPTP may open and triggers several deleterious events. Opening of the mPTP abolished the mitochondrial membrane potential ($\Delta\psi_m$) inhibiting oxidative phosphorylation. The mPTP opening allows the equilibration of molecules that are smaller than approximately 1,500 daltons. Osmotic force generated by matrix proteins results in matrix swelling (mitochondrial swelling), leading to further rupture of the outer membrane and the release of proapoptotic factors, such as cytochrome c, into the cytosol. These actions rapidly produce cell death. In addition, disruption of the mitochondrial membrane potential also causes the ATP synthase to behave as an ATPase and accelerate energy depletion secondary to the ischemic insult⁽¹²⁾.

There are experimental evidences that support that mPTP opening is a postischemic event. In the isolated rat heart model, it has been demonstrated that the cytosolic release of NAD^+ (this molecule was used as a surrogate marker of mPTP opening), occurs at the time of reperfusion following

a prolonged ischemic insult⁽¹³⁾. Griffiths and Halestrap used the [³H]2-deoxyglucose entrapment technique to investigate the kinetics of *in situ* mPTP opening, and demonstrated that mPTP opening does not happen during ischemia, but occurs within the first 5 minutes of reflow following a 30-minute period of ischemia in the isolated rat heart⁽¹⁴⁾. Importantly, the time course of mPTP opening appeared to match the rapid correction of pH that occurs at reperfusion. Recent *in vivo* studies support his concept by showing that PostC may mediate its cardioprotective effects through prolonged transient acidosis during the early reperfusion phase⁽¹⁵⁾.

Inflammatory changes and endothelial function during reperfusion

During reperfusion there are several altered responses in other tissues or systems besides the cardiac muscle, forming a set of events that make up the reperfusion damage. Alterations of endothelial function are pivotal in the development of reperfusion damage and the no-reflow phenomenon. The inflammatory process characterizing early and late periods of reperfusion is an important aspect of changes leading to tissue damage by its effects on endothelial cells. Neutrophils feature prominently in the inflammatory component of post-ischemic injury. This occurs because ischemia-reperfusion prompts a release of oxygen free radicals, cytokines and other pro-inflammatory mediators that activate both the neutrophils and the coronary vascular endothelium⁽¹⁶⁾. Activation of these

cells promotes the expression of adhesion molecules on both neutrophils and the endothelium, which recruit neutrophils on the endothelial surface and initiate a specific cascade of cell-cell interaction. This lead first to the adhesion of neutrophils to the vascular endothelium and subsequently to their transendothelial migration and their direct interactions with interstitial matrix and myocytes^(17,18). This specific series of events is a prerequisite for the full expression of reperfusion injury, including endothelial dysfunction, microvascular collapse, and impairment of blood flow (the «no reflow» phenomenon), myocardial infarction and apoptosis.

POSTCONDITIONING TO PROTECT THE HEART

In the procedure called myocardial PostC, the heart can be protected against the ischemia reperfusion injury with brief coronary occlusions performed just at the beginning of the reperfusion. The postC it was first described by Zhao and colleagues in dogs⁽¹⁹⁾, which reduced the myocardial injury to an extent comparable to IPC. Beneficial outcomes observed with PostC included reduction in infarct size^(20,21), in endothelial dysfunction, in neutrophil adherence^(19,21), and in apoptosis⁽²²⁾. Recently, it has been shown that a range of pharmacological agents given at the moment of reperfusion after an ischemic insult can significantly protect the myocardium, and this effect has been shown to involve protection against necrosis and apoptosis. This maneuver it has been called: pharmacological postC.




Figure 2. ATP dissipation during ischemia leads to rises in resting cytosolic free [Ca²⁺]. Reperfusion leads to excessive mitochondrial Ca²⁺ uptake. Mitochondrial Ca²⁺ overload together with oxidative stress and the prevailing low ATP provoke mPTP opening. These events initiate a «vicious cycle», i. e. inner-membrane depolarization, ATP hydrolysis by the mitochondrial ATP synthase, further increases in cytosolic Ca²⁺, and so on, leading finally, to cell death.

PROPOSED MECHANISMS OF POSTC

Studies have documented that several pathways and molecular components are involved in the cardioprotective effects of postC. These including: NO⁽²³⁾, phosphatidylinositol 3-kinase (PI3K)⁽²⁴⁾, Extracellular signal-Regulated Kinase (ERK)⁽²⁵⁾, PKC⁽²⁶⁾, and mitochondrial adenosine triphosphate-sensitive potassium channels (mitoKATP)⁽²⁷⁾, Reperfusion Injury Salvage Kinases (RISK), and Survivor Activating Factor Enhancement (SAFE) pathways.

mPTP as potential end-effector of postC

As was described, the mitochondria and the mPTP opening have been proposed to play an essential role in reperfusion injury⁽²⁸⁾, and in this way, inhibition of mPTP opening has been reported to be an important mechanism underlying postC protection⁽²⁹⁾. A number of studies have discovered a common finding with regard to the timing of postC. The protection induced can only be taken advantage of if postC is initiated at the onset of reperfusion and is lost if is delayed by 1 minute in the rat⁽³⁰⁾, and rabbit models⁽³¹⁾. Therefore, this would suggest that the end-effector of protection must exert its actions during the initial stages of reperfusion. In this regard, pharmacologically inhibiting the mPTP during the first few minutes of reperfusion⁽³²⁾ has been shown to be cardioprotective and delaying this inhibition by 15 minutes abolishes this protection. Similarly, delaying the administration of insulin, which is known to activate the RISK pathway, until after the first 15 minutes of reperfusion, abrogates its infarct-limiting effect⁽³³⁾. Therefore, one could speculate that the mPTP, which is known to regulate cell death during the first few minutes of reperfusion⁽³⁴⁾, is potentially the main candidate as the end-effector. Others studies have assessed the interaction between the mPTP opening, and the postconditioning interventions. In a rabbit model of ischemia-reperfusion, Argaud et al. found that both, mechanical and pharmacological PostC with cyclosporin A (CsA) or NIM811, a specific inhibitor of the mPTP that was given 1 minute before reperfusion, limit infarct size by at least 45% compared with control animals⁽³⁵⁾. Thus, they demonstrated that the suppression of mPTP opening at reperfusion provided powerful antinecrotic and anti-apoptotic protection to the ischemic myocardium. They also showed that both mechanical and pharmacological PostC, increase the Ca²⁺ load that is required to open the mPTP⁽³⁶⁾. Additional evidence for a major role of the mPTP in lethal reperfusion injury recently came from the use of transgenic mice lacking cyclophilin D (CypD)^(37,38). CypD, which is recognized as a key molecular component of the mPTP, is a mitochondrial member of the family of peptidyl-prolyl cis-trans isomerases (PPIases). Although still debated, it was reported that, in the presence of a high matrix Ca²⁺ concentration, CypD modifies

the conformation of the inner membrane proteins to form a mega-channel. The molecular structure of the mPTP remains poorly known and, besides CypD, might involve various others proteins including voltage-dependent anion channel (VDAC) or adenine nucleotide translocator (ANT), as was mentioned above. Unfortunately, their precise role is still elusive and no pharmacological agent targeting these proteins is currently available for clinical trials. *In vivo*, CypD-deficient mice develop smaller infarcts after a prolonged coronary artery occlusion followed by reperfusion^(37,38). Recently, Lim et al. reported that CypD-deficient mice cannot be postconditioned, further suggesting that lethal reperfusion injury is mediated by mPTP opening⁽³⁹⁾. These results strongly support the proposal that mPTP opening, triggered by mitochondrial Ca²⁺ overload and overproduction of reactive oxygen species (ROS), plays a central role in lethal reperfusion injury and, specific inhibition of mPTP opening at the time of reperfusion is central to the cardioprotective effect induced by PostC.

ADDITIONAL MECHANISMS OF POSTC

RISK Pathway

During the early stages of reperfusion, there is an up-regulation of pro-survival kinases termed: the Reperfusion Injury Salvage Kinase (RISK) pathway, which recently, has promoted a renewed interest in cardioprotective reperfusion strategies⁽⁴⁰⁾ (Figure 3). While the mechanisms involved in reperfusion injury are known to involve apoptosis and necrosis, among other, it is also realized that cells have an inherent program for survival after ischemia-reperfusion insults, via the recruitment of innate prosurvival kinase cascades. This pathway compresses to PI3-kinases and p42/p44 extra-cellular signal-regulated kinases (ERK 1/2) which could activated by ischemia-reperfusion injury or by PostC or Pharmacological PostC. The PI3-kinases (PI3Ks) are a family of enzymes involved in several functions such as cell growth, proliferation, differentiation, motility, survival and intracellular trafficking. These enzymes are capable of phosphorylating the 3-position hydroxyl group of the inositol ring of phosphatidylinositol. Particularly, the PI3K-Akt (Akt = Protein kinase B) and MEK kinases (MEK kinase = MAPK/ERK kinase; Mitogen-Activated Protein kinase/Extracellular signal-Regulated Kinase) have shown to be important components of the cell survival pathway and have antiapoptotic effects. There are several potential mechanisms, none of them mutually exclusive, through which this pathway mediates inhibition of mPTP opening. These are: 1) Phosphorylating and inhibiting GSK3 β ⁽⁴¹⁾, 2) Phosphorylating eNOS, producing nitric oxide, which has been demonstrated to inhibit mPTP opening⁽⁴²⁾, and 3) phosphorylating BAD, either directly or indirectly via p70S6 Kinase^(43,44), thereby negating its pro-

apoptotic effect. Recently, Davidsson et al.⁽⁴⁵⁾ found that insulin-mediated phosphorylation of PI3K-Akt protects the myocyte against oxidative stress by inhibiting mPTP. This suggests that the pharmacological activation of the RISK pathway by insulin at the onset of reperfusion protects against ischemia-reperfusion injury by reducing the probability that the mPTP will open. This mechanism could participate in the cardioprotective effect of «GIK» solution, which is composed by insulin in combination with glucose and potassium. However, clinical studies have suggested that the GIK solution may be cardioprotective following myocardial ischemia, via a direct, non-metabolic cardioprotective effect^(46,47). Albeit, whether the RISK pathway kinases and the mPTP is involved in the clinical setting remains to be examined. At this moment is necessary to ascertain the potential mechanism where by the RISK pathway delays the mPTP opening and protects the cells from injury, and this issue is part of the setting of

preconditioning or protection against reperfusion-induced injury in experimental models.

POSTCONDITIONING THE HUMAN HEART

The ultimate validation and utility of a cardioprotective therapy is its application to humans presenting with coronary artery disease and concomitant risk factors. However, at date there are very few studies conducted in humans, this situation is understandable because of high risks that his approach represents. Two studies have recently reported that conventional PostC is an effective treatment in a select patient population with coronary artery disease. In a study by Laskey⁽⁴⁸⁾, 17 patients undergoing PCI were enrolled to receive standard of care angioplasty involving 90 s of uninterrupted balloon inflation without further treatment (n = 7) or repeated balloon inflation («conditioning», n = 10) of 90 s duration applied 3-5 min after the angioplasty inflation. «Conditioning» after angioplasty reduced the magnitude of ST-segment elevation compared to controls, and accelerated the rate at which ST elevation normalized after reperfusion. Furthermore, blood flow velocity reserve was significantly improved in «conditioned» hearts. Staat et al.⁽⁴⁹⁾, reported a multi-center randomized clinical trial of 37 patients with total coronary artery occlusion undergoing angioplasty/stenting. Patients that achieved a TIMI flow grade of 2-3 at completion of the angioplasty/stent procedure were randomized to receive either standard of care treatment thereafter or postC with 4 cycles of 1 min re-inflation followed by 1 min deflation of the angioplasty balloon. Infarct size was significantly less, and the coronary blood flow achieved was greater in the postconditioned patients. There were no adverse events in the postconditioned patients. Together, these two studies suggest that postC represents a safe and efficient cardioprotective intervention for treatment of reperfusion injury in patients with ischemic heart disease. These data must be reproduced by other clinical trials, and in broader patient populations, particularly those presenting with severe coronary artery disease involving high risk factors (hypertension, hypercholesterolemia, obesity, diabetes).




Figure 3. Scheme showing the proposed mechanism of protection induced by ischemic postconditioning (that could be triggered by ischemia reperfusion, too). The gradual reperfusion may have an effect via activation of phosphatidylinositol 3-kinase (PI3K)-Akt or ERK 1/2, phosphorylates downstream targets such as glycogen synthase kinase-3 β (GSK- β), BAD/Bax, and endothelial (NO) synthase (eNOS), producing NO, which inhibits mitochondrial permeability transition pore (mPTP) opening. Phosphorylation of p70s6K confers protection by inactivating Bcl-2-Associated Death promoter (BAD) or through protein translation.

PHARMACOLOGICAL POSTC

In the clinical situations, ischemic PostC may be difficult conceptually to introduce, i.e., reintroduction of ischemia at the time of reperfusion may lead to potential complications. These complications are the cause of the initial refusal to use this approach in clinical setting. For example, during primary angioplasty, repetitive inflations and deflations of the balloon may result in coronary plaque rupture with consequences for restenosis or embolic events. During coronary bypass

surgery, interruptions to reperfusion via the newly grafted conduit may only lead to regional myocardial protection in the area supplied by that bypass. An alternative strategy during bypass surgery would be the repetitive clamping and unclamping of the ascending aorta to achieve ischemic PostC, a concept, however, that many cardiac surgeons would be unwilling to perform due to the high risk of disrupting atheromatous plaque debris and subsequent risk of stroke. The use of ischemic PostC would not be possible in an acute myocardial infarction patient referred for thrombolysis, in case of unstable angina, in patients presenting with non-ST segment elevation myocardial infarction or in the event of a cardiac arrest. Such a protocol might be possible to apply in a patient with myocardial infarction referred for Percutaneous Coronary Intervention (PCI), elective PCI or at the time of cardiac surgery. However, the concept of «pharmacological PostC» by the administration of agents that activate the RISK pathway and mediate the protective effects of postC is a more practical solution. Agents such as insulin⁽⁴⁴⁾, atorvastatin⁽⁵⁰⁾, bradikinin⁽⁵¹⁾, tumor growth factor (TGF)-B⁽⁵²⁾, and glucagon-like peptide 1 (GLP-1)⁽⁵³⁾, anesthetic⁽⁵⁴⁻⁵⁹⁾, opioids⁽⁶⁰⁻⁶³⁾ could individually be used as adjuvant to current reperfusion strategies such as thrombolytics and primary PCI to limit lethal reperfusion injury and could form the basis of much needed and important reperfusion strategies.

Anesthetic postconditioning (APC)

Experimental evidence indicated that volatile anesthetics are capable of exerting cardioprotective effects under these conditions. For example, halothane prevented reoxygenation-induced hypercontracture of cardiac myocytes *in vitro*, a potential cause of myocyte necrosis during early reperfusion⁽⁶⁴⁾. Halothane also reduced reperfusion injury after regional myocardial ischemia in rabbit hearts⁽⁶⁵⁾. Desflurane and sevoflurane reduced infarct size when administered during the first 15 minutes of reperfusion in rabbits⁽⁶⁶⁾. Isoflurane enhanced the functional recovery of isolated rat hearts when administered solely during reperfusion⁽⁶⁷⁾. The administration of sevoflurane after ischemia also improved contractile and metabolic function concomitant with reduced myoplasmic calcium loading in isolated guinea pig hearts⁽⁶⁸⁾. These and other finding indicated that volatile anesthetics may prevent intracellular calcium overloading during early reperfusion, presumably by virtue of their actions as voltage-dependent calcium channel antagonist isoflurane and sevoflurane also reduced postischemic adhesion of neutrophils⁽⁶⁹⁾, an important source of oxygen-derived free radicals during reperfusion that are known to be critical mediators of reperfusion injury⁽⁷⁰⁾. Several protective effects of inhalation anaesthetics have been proposed (Figure 4).

Opioid and postC

Activation of G protein-coupled receptors such as adenosine and opioid receptors has been demonstrated to initiate preconditioning. In the case of PostC, the G protein-coupled receptor activation may serve as essential mechanisms that trigger the protection too. Indeed, recent reports have addressed that adenosine A2a and A2b receptors are responsible for the protection of postC⁽⁷¹⁾. Recently, Gross et al, demonstrated that opioids can reduce infarct size when administered just before reperfusion, an effect that was similar to that observed when opioids were given before ischemia⁽⁷²⁾. Furthermore, Weihrauch et al.⁽⁷³⁾, reported that morphine enhanced the isoflurane-induced postC in rabbit hearts. Therefore, it is possible that postC protects the heart by activating opioid receptors during reperfusion. On other hand, activation of delta opioid receptors plays an important role in postC and may protect the heart at reperfusion by modulating the mPTP opening via a signaling cascade involving delta and kappa opioid receptors, NO, PKC and other pathways (Figure 5). Gross et al.⁽⁷²⁾ and Fryer et al.⁽⁷⁴⁾ showed that opioid-induced postC was mediated by activation of G protein-coupled delta 1 opioid receptors, PI3K-mediated signaling, and mitoKATP channels. Two selective delta 1 opioid receptor agonist (TAN-67 and BW373U86) and morphine were initially shown to augment isoflurane preconditioning in rats⁽⁷⁵⁾, and the nonselective opioid antagonist naloxone inhibited these salutatory actions. Interestingly, naloxone pretreatment also abolished reductions in infarct size produced by administration of isoflurane alone before ischemia and reperfusion, showing for the first time that opioid receptors mediate anesthetic preconditioning. Our group, at the pharmacology department at Instituto Nacional




Figure 4. Protective effects of potent inhalation anaesthetics. APD = Action Potential Duration.

de Cardiología «Ignacio Chávez» showed that remifentanyl, an agonist of opioid receptors, elicited antiarrhythmic and cardioprotective effects in experimental ventricular arrhythmias induced by digoxin and in a model of two atrial arrhythmias induced by aconitine and by electrical stimulation⁽⁷⁶⁾.

Glycine cardioprotection

The amino acid glycine (gly) has been shown to exert a protective effect against cell death after diverse insults including ischemia-reperfusion injury in different cell types other than cardiomyocytes⁷⁷ (Figure 6). Gly is an inhibitory neurotransmitter in the central nervous system, which has been demonstrated by a wide range of studies in the past few years. Gly prevents or mitigates a variety of pathological processes in experimental models, such as inflammation, shock, endotoxaemic shock, and those associated with transplantation. The classic inhibitory gly receptors (GlyR) in the central nervous system is a ligand-gated membrane-spanning ion channel^(78,79). GlyR consists of three types of subunits: (alpha), a ligand binding subunit; (beta), a structural subunit, and gephyrin, a cytoplasmic anchoring protein⁽⁸⁰⁾. Gly in the central nervous system exerts its inhibitory actions by binding GlyR that is located mainly on the postsynaptic neuronal membranes. Activation of GlyR leads to increase in chlo-

ride conductance⁽⁸¹⁾. An influx of chloride hyperpolarizes postsynaptic membranes, thus counteracting the depolarizing action of excitatory neurotransmitters. It is proposed that hyperpolarization decreases the opening of calcium




Figure 6. Proposed role of Gly during reperfusion injury.




Figure 5. Scheme of the activation on the RISK pathway induced by δ - and κ -opioid receptor. ROS = Reactive Oxygen Species; PKC = Protein Kinase C.

channels, thereby blocking the movement of calcium across the plasma membrane and the subsequent events in the inflammatory process. Recent studies have provided pharmacological and molecular evidence for the existence of GlyR in endothelial cells⁽⁸²⁾, renal proximal tubular cells⁽⁸³⁾, and most leukocytes⁽⁸⁴⁾. Several investigators reported the benefits of N-(2-mercaptopropionyl)-glycine (MPG), a hydroxyl radical scavenger, for ischemic-reperfused hearts; treatment of anesthetized animals with this agent reduced ischemia-reperfusion-induced formation of the infarct size⁽⁸⁵⁾, and improved ischemia-reperfusion induced myocardial stunning⁽⁸⁶⁾. Furthermore, treatment of perfused hearts with MPG enhanced the recovery of contractile function and preserved the ultrastructural integrity of the myocardium⁽⁸⁷⁾. Ruiz-Meana study shows that the amino acid Gly has a powerful and previously unrecognized inhibitory effect on mPTP, similar of that of low pH, in rat heart mitochondria. They also demonstrate for the first time that addition of Gly during reoxygenation prevents acute necrotic cell death associated with pH normalization in cultured cardiac myocytes and LDH release in isolated rat hearts. Moreover, exposure to Gly during reoxygenation not only reverses cellular Gly depletion observed in cells submitted to ischemia-reoxygenation, but markedly raises intracellular Gly content up to the range proven to inhibit mPTP in isolated mitochondria⁽⁸⁸⁾.

Postconditioning the human heart with adenosine

Adenosine is a primary mediator of postconditioning effects. Studies performed in animal models indicate that adenosine receptor activation is involved in the cardioprotection conferred by postC. The underlying mechanism may involve delaying the washout of endogenously released adenosine during the early minutes of reperfusion⁽⁸⁹⁾. The protective effects could be blocked by the nonselective adenosine receptor antagonist, 8SPT⁽⁹⁰⁾ and by A2a and A3 selective antagonists given before postC. Adenosine has also been implicated in the cardioprotection exerted by remote postC⁽⁹¹⁾. In one study of sixty patients with rheumatic heart valve disease undergoing heart valve replacement operations were randomized to an adenosine (1.5 mg/kg) or saline (control) bolus injection through an arterial catheter immediately after the aorta cross-clamp was removed. The results of this study were, the inotrope scores in the intensive care unit (ICU) were much lower, and the ICU time was significantly shorter in adenosine group. More important, cardiac troponin I release was less especially at 12 and 24 hours after reperfusion. The present study demonstrated that pharmacologic postC with adenosine produces protective effects against myocardial reperfusion injury in cardiac operations⁽⁹²⁾.

ADVANCES IN POSTC

Activation of the protective survivor activating factor enhancement (SAFE) pathway against reperfusion injury

Novel strategies to protect the heart during myocardial reperfusion have emerged as very promising experimental therapies against lethal reperfusion injury. One of them, is the activation of the protective survivor activating factor enhancement (SAFE) pathway⁽⁹³⁾. This path includes the participation of the Tumor Necrosis Factor-alpha, TNF α , and the STAT-3 (STAT = Signal Transducer and Activator of Transcription). TNF α levels (blood and tissue) are increased in acute myocardial infarction and generally, TNF α is implicated as a mediator of adverse remodeling in heart failure⁽⁹⁴⁾. TNF α exerts its action after binding onto its cell surface receptors. TNF α receptor 1 (TNFR1 or p55) and TNF α receptor 2 (TNFR2 or p75). Both receptors subtypes are found in human and rat cardiac myocytes⁽⁹⁵⁾. There are controversial findings that have been described with beneficial and detrimental effects for TNF α ^(96,97). These differences about its effects can be explained on which TNF α receptor is activated. Thus, TNF α 1 may be cardiotoxic while TNF α receptor 2 could be cardioprotective. Lecour et al.⁽⁹⁸⁾ have demonstrated that exogenous TNF α , in a dose and time dependent manner, could mimic ischemic preconditioning in the rat or mouse model.

The mechanisms proposed for the protective effect of TNF α is the following: After activation of the membrane receptors by TNF α (and others receptors could be participating like interleukin 6 or growth factors receptors), two adjacent juxtaposed JAKs (Janus kinase) are transphosphorylated and subsequently activate STAT pathway. JAKs are a family of tyrosine kinases that are associated with membranes receptors and play a major role in the transduction of signals from the cytosol to the nucleus and they are activated during ischemia and ischemia reperfusion in the heart. The STAT family of transcription factor proteins consists of seven identified members: STAT-1, STAT-2, STAT-3, STAT-4, STAT-5A, STAT-5B and STAT6. After activation with JAKs, STAT form homo- or heterodimers that are translocated to the nucleus, resulting in gene transcription (Figure 7). The activation of the JAK/STAT pathway plays a crucial role in the expression of stress-responsive genes that could be participating in the protective effects⁽⁹⁹⁾.

Acting together SAFE and RISK pathways

These distinct pathways are activated at the time of reperfusion following an ischemic preconditioning insult, but whether these two pathways may interact together or be totally independent remains unclear. The Lecour group has found that




Figure 7. The JAK/STAT signaling pathway. JAK-2 is activated in response to a ligand (like $\text{TNF}\alpha$). Activated JAK-2 recruits STAT-3 (signal transducer and activator of transcription) from the cytoplasm and phosphorylates STAT-3 on a tyrosine residue, which allows STAT-3 to form homodimers and translocate to the nucleus where it up regulates transcription. STAT-3 is further phosphorylated on a serine residue to improve DNA interaction.

$\text{TNF}\alpha$ -mediated preconditioning acts independently of the activation of the RISK pathway, therefore suggesting that activation of both pathways at the time of reperfusion is not always mandatory to protect the heart against reperfusion injury⁽⁹⁸⁾. However, the inhibition of either the RISK or the SAFE pathway in ischemic preconditioning totally abolished the protection in the ischemic preconditioning, suggesting the possibility of a cross-talk between the two pathways at level of PI3K/Akt and JAK/STAT-3⁽⁹⁸⁻¹⁰⁰⁾ (Figure 8).

The precise molecular mechanisms and pathways of signaling protective cascades are far from being fully elucidated.




Figure 8. Activation of the RISK and SAFE pathways following ischemic pre- or post conditioning. Extracellular signal-regulated protein kinase (ERK).

However, there is now strong evidence for the existence of at least two pathways with $\text{TNF}\alpha$, which play a key role in the immune system. Therefore it represents a novel and alternative protective SAFE path during both ischemic pre- and postconditioning.

REFERENCES

- Lloyd-Jones D, Adams R, Carnethon M, De Simone G, Ferguson TB, Flegal K, et al. Heart disease and stroke statistics-2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2009;119:e182.
- Gibbons RJ, Valeti US, Araoz PA, Jaffe AS. The quantification of the infarct size. *J Am Coll Cardiol* 2004;44:1533-1542.
- McGovern PG, Pankow JS, Shahar E, Doliszny KM, Folsom AR, Blackburn H, and Luepker RV. Recent trends in acute coronary heart disease-mortality, morbidity, medical care and risk factors. The Minnesota Heart Survey Investigators. *N Engl J Med* 1996;334:884-890.
- Yellon DM, Hausenloy DJ. Myocardial reperfusion injury. *N Engl J Med* 2007;357:1121-1135.
- Jennings RB, Sommers HM, Smyth GA, Flack HA, Linn H. Myocardial necrosis induced by temporary occlusion of a coronary artery in the dog. *Arch Pathol* 1960;70:68-78.
- Wu KC, Zerhouni EA, Judd RM, Lugo-Olivieri CH, Barouch LA, Schulman SP, Blumenthal RS, Lima JA. Prognostic significance of microvascular obstruction by magnetic resonance imaging in patients with acute myocardial infarction. *Circulation* 1998;97:765-772.
- Das AM, Harris DA. Regulation of the mitochondrial ATP synthase in intact rat cardiomyocytes. *Biochem J* 1990;266:355-361.

8. Dennis S C, Gevers W, Opie LH. Protons in ischemia: where do they come from; where do they go? *J Mol Cell Cardiol* 1991;23:1077-1086.
9. Schlüter KD, Jakob G, Ruiz-Meana M, Garcia-Dorado D, Piper HM. Protection of reoxygenated cardiomyocytes against osmotic fragility by NO donors. *Am J Physiol* 1996;271:H428-H434.
10. Piper HM, Abdallah Y, Schafer C. The first minutes of reperfusion: a window of opportunity for cardioprotection. *Cardiovasc Res* 2004;61:365-371.
11. Crompton M. The mitochondrial permeability transition pore and its role in cell death. *Biochem J* 1999;341:233-249.
12. Halestrap AP, Clarke SJ, Javadov SA. Mitochondrial permeability transition pore opening during myocardial reperfusion –a target for cardioprotection. *Cardiovasc Res* 2004;61:372-385.
13. Di Lisa et al. 2001 = Di Lisa F, Menabo R, Canton M, Barile M, Bernardi P. Opening of the mitochondrial permeability transition pore causes depletion of mitochondrial and cytosolic NAD⁺ and is a causative event in the death of myocytes in postischemic reperfusion of the heart. *J Biol Chem* 2001;276:2571-2575.
14. Griffiths EJ, Halestrap AP. Mitochondrial non-specific pores remain closed during cardiac ischaemia, but open upon reperfusion. *Biochem J* 1995;307: 93-98.
15. Cohen MV, Yang XM, Downey JM. Acidosis, oxygen, and interference with mitochondrial permeability transition pore formation in the early minutes of reperfusion are critical to postconditioning's success. *Basic Res Cardiol* 2008;103:464-471.
16. Jordan JE, Zhao ZQ, Vinten-Johansen J. The role of neutrophils in myocardial ischemia-reperfusion injury. *Cardiovasc Res* 1999;43:860-878.
17. Engler RL, Schmid-Schnbein GW, Pavelec RS. Leukocyte capillary plugging in myocardial ischemia and reperfusion in the dog. *Am J Pathol* 1983;111:98-111.
18. Moore KL, Patel KD, Bruel RE. P-selectin glycoprotein ligand-1 mediates rolling of human neutrophils on P-selectin. *J Cell Biol* 1995;128:661-671.
19. Zhao ZQ, Corvera JS, Halkos ME. Inhibition of myocardial injury by ischemic postconditioning during reperfusion: comparison with ischemic preconditioning. *Am J Physiol Heart Circ Physiol* 2003;285:H579.
20. Kin H, Zatta AJ, Lofye MT. Postconditioning reduces infarct size via adenosine receptor activation by endogenous adenosine. *Cardiovasc Res* 2005;67:124.
21. Halkos ME, Kerendi F, Corvera JS. Myocardial protection with postconditioning is not enhanced by ischemic preconditioning. *Ann Thorac Surg* 2004;78:961.
22. Sun HY, Wang NP, Halkos M. Postconditioning attenuates cardiomyocyte apoptosis via inhibition of JNK and p38 mitogen activated protein kinase signaling pathways. *Apoptosis* 2006;11:1583.
23. Yang X-M, Proctor JB, Cui L, Krieg T, Downey JM, Cohen MV. Multiple, brief coronary occlusions during early reperfusion protect rabbit hearts by targeting cell signaling pathways. *J Am Coll Cardiol* 2004;44:1103-1110.
24. Yang X-M, Philipp S, Downey JM, Cohen MV. Postconditioning's protection is not dependent on circulating blood factors or cells but involves adenosine receptors and requires PI3Kinase and guanylyl cyclase activation. *Basic Res Cardiol* 2005;100:57-63.
25. Tsang A, Hausenloy DJ, Mocanu MM, Yellon DM. Postconditioning a form of "modified reperfusion" protects the myocardium by activating the phosphatidylinositol 3-kinase-akt pathway. *Circ Res* 2004;95:230-232.
26. Zatta AJ, Kin H, Lee G, Wang N, Jiang R, Lust R, Reeves JG, Mykitenko J, Guyton RA, Zhao ZQ, Vinten-Johansen J. Infarct-sparing effect of myocardial postconditioning is dependent on protein kinase C signaling. *Cardiovasc Res* 2006;70:315-324.
27. Jang Y, Xi J, Wang H, Mueller RA, Norfleet EA, Xu Z. Postconditioning prevents reperfusion injury by activating μ -opioid receptors. *Anaesthesiology* 2006;108:243-250.
28. Griffiths EJ, Halestrap AP. Protection by cyclosporin A of ischemia/reperfusion-induced damage in isolated rat hearts. *J Mol Cell Cardiol* 1993;25:1461-1469.
29. Argaud I, Gateau-Roesch O, Risky O, Loufoat J, Robert D, Ovize M. Postconditioning inhibits mitochondrial permeability transition. *Circulation* 2005;111:194-197.
30. Philipp S, Downey JM, Cohen MV. Postconditioning must be initiated in less than 1 minute following reperfusion and is dependent on adenosine receptors and PI3-kinase. (Abstract) *Circulation* 2004;110:suppl 111:804.
31. Hausenloy DJ, Duchon MR, Yellon DM. Inhibiting mitochondrial permeability transition pore opening at reperfusion protects against ischaemia-reperfusion injury. *Cardiovasc Res* 2003;60:617-625.
32. Hausenloy DJ, Maddock HL, Baxter GF, Yellon DM. Inhibiting mitochondrial permeability transition pore opening: a new paradigm for myocardial preconditioning? *Cardiovasc Res* 2002;55:534-543.
33. Jonassen AK, Sack MN, Mjos OD, Yellon DM. Myocardial protection by insulin at reperfusion requires early administration and is mediated via Akt and p70s6 kinase cell-survival signaling. *Circ Res* 2001;89:1191-1198.
34. Sun K, Liu ZS, Sun Q. Role of mitochondria in cell apoptosis during hepatic ischemia-reperfusion injury and protective effect of ischemic postconditioning. *World J Gastroenterol* 2004;10:1934-1938.
35. Argaud L, Gateau-Roesch O, Risky O, Loufoat J, Robert D, Ovize M. Postconditioning inhibits mitochondrial permeability transition. *Circulation* 2005;111:194-197.
36. Argaud L, Gomez L, Gateau-Roesch O, Couture-Lepetit E, Loufoat J, Robert D, Ovize M. Trimetazidine inhibits mitochondrial permeability transition pore opening and prevents lethal ischemia-reperfusion injury. *J Mol Cell Cardiol* 2005;39:893-899.
37. Baines CP, Kaise RA, Purcell NH, Blair NS, Osinska H, Hambleton MA, Brunskill EW, Sayen MR, Gottlieb RA, Dorn GW, et al. Loss of cyclophilin D reveals a critical role for mitochondrial permeability transition in cell death. *Nature* 2005;434:658-662.
38. Nakagawa T, Shimizu S, Watanabe T, Yamaguchi O, Otsu K, Yamagata H, Inohara H, Kubo T, Tsukimoto Y. Cyclophilin D-dependent mitochondrial permeability transition regulates some necrotic but not apoptotic cell death. *Nature* 2005;434:652-658.
39. Lim SY, Davidson SM, Hausenloy DJ, Yellon DM. Preconditioning and postconditioning: the essential role of the mitochondrial permeability transition pore. *Cardiovasc Res* 2007;75:530-535.
40. Hausenloy DJ, Yellon DM. New directions for protecting the heart against ischaemia-reperfusion injury: targeting the reperfusion injury salvage kinase (RISK)-pathway. *Cardiovasc Res* 2004;61:448-460.
41. Juhaszova M, Zorov DB, Kim SH, Pepe S, Fu Q, Fishbein KW, et al. Glycogen synthase kinase-3 beta mediates convergence of protection signaling to inhibit the mitochondrial permeability transition pore. *J Clin Invest* 2004;113:1535-1549.
42. Kim JS, Ohshima S, Padiaditakis P, Lemasters JJ. Nitric oxide protects rat hepatocytes against reperfusion injury mediated by the mitochondrial permeability transition. *Hepatology* 2004;39:1533-1543.
43. Jonassen AK, Mjos OD, Sack MN. p70s6 kinase is a functional target of insulin activated Akt cell-survival signaling. *Biochem Biophys Res Commun* 2004;315:160-165.
44. Jonassen AK, Sack MN, Mjos OD, Yellon DM. Myocardial protection by insulin at reperfusion requires early administration and is mediated via Akt and p70s6 kinase cell-survival signaling. *Circ Res* 2001;89:1191-1198.
45. Davidson SM, Hausenloy D, Duchon MR, Yellon DM. Signalling via the reperfusion injury signalling kinase (RISK) pathway links closure of the mitochondrial permeability transition pore to cardioprotection. *The International Journal of Biochemistry & Cell Biology* 2006;38:414-419.
46. Doenst T, Bothe W, Beyersdorf F. Therapy with insulin in cardiac surgery: Controversies and possible solutions. *Ann Thorac Surg* 2003;75:S721-S728.

47. Sack MN, Yellon DM. Insulin therapy as an adjunct to reperfusion after acute coronary ischemia: A proposed direct myocardial cell survival effect independent of metabolic modulation. *J Am Coll Cardiol* 2003;41:1404-1407.
48. Laskey WK. Brief repetitive balloon occlusions enhance reperfusion during percutaneous coronary intervention for acute myocardial infarction: a pilot study. *Catheter Cardiovasc Interv* 2005;65:361-367.
49. Staat P, Rioufol G, Piot C, Cottin Y, Cung TT. Postconditioning the human heart. *Circulation* 2005;112:2143-2148.
50. Bell RM, Yellon DM. Atorvastatin, administered at the onset of reperfusion and independent of lipid lowering, protects the myocardium by up-regulating a pro-survival pathway. *J Am Coll Cardiol* 2003;41:508-515.
51. Bell RM, Yellon DM. Bradykinin limits infarction when administered as an adjunct to reperfusion in mouse heart: the role of PI3k, Akt and eNOS. *J Mol Cell Cardiol* 2003;35:185-193.
52. Baxter GF, Mocanu MM, Brar BK, Latchman DS, Yellon DM. Cardioprotective effects of transforming growth factor-beta 1 during early reoxygenation of reperfusion are mediated by p42/p44 MAPK. *J Cardiovasc Pharmacol* 2001;38:930-939.
53. Nikolaidis LA, Mankad S, Sokos GG, Miske G, Shan A, Elani D, Shannon RP. Effects of glucagon-like peptide-1 in patients with acute myocardial infarction and left ventricular dysfunction after successful reperfusion. *Circulation* 2004;109: 962-965.
54. Schlack W, Preckel B, Barthel H. Halothane reduces reperfusion injury after regional ischaemia in the rabbit heart *in vivo*. *Br J Anaesth* 1997;79:88-96.
55. Preckel B, Schlack W, Comfere T. Effects of enflurane, isoflurane, sevoflurane and desflurane on reperfusion injury after regional myocardial ischaemia in the rabbit heart *in vivo*. *Br J Anaesth* 1998;81:905-912.
56. Schlack W, Preckel B, Stunneke D. Effects of halothane, enflurane, isoflurane, sevoflurane and desflurane on myocardial reperfusion injury in the isolated rat heart. *Br J Anaesth* 1998;81:913-919.
57. Varadarajan SG, An J, Novalija E. Sevoflurane before or after ischemia improves contractile and metabolic function while reducing myoplasmic Ca²⁺ loading in intact hearts. *Anesthesiology* 2002;96:125-133.
58. Heindl B, Reichle FM, Zahler S. Sevoflurane and isoflurane protect the reperfused guinea pig heart by reducing postischemic adhesion of polymorphonuclear neutrophils. *Anesthesiology* 1999;91:521-530.
59. Vinten-Johansen J. Involvement of neutrophils in the pathogenesis of lethal myocardial reperfusion injury. *Cardiovasc Res* 2004;61:481-497.
60. Jang Y, Xi J, Wang H, Mueller RA, Norfleet EA, Xu Z. Postconditioning prevents reperfusion injury by activating δ -opioid receptors. *Anesthesiology* 2006;108:243-250.
61. Gross ER, Hsu AK, Gross GJ. Opioid-induced cardioprotection occurs via glycogen synthase kinase β inhibition during reperfusion in intact rat hearts. *Circ Res* 2004;94:960-966.
62. Weihrauch D, Riotkowski JG, Bienengracher M, Hersten JR, Wartier DC, Pagel PS. Morphine enhances isoflurane-induced postconditioning against myocardial infarction: The role of phosphatidylinositol-3-kinase and opioid receptors in rabbits. *Anesth Analg* 2005;101:942-949.
63. Gross ER, Hsu AK, Gross GJ. GSK3 β inhibition and KATP channel opening mediate acute opioid-induced cardioprotection at reperfusion. *Basic Res Cardiol* 2007;102:341-349.
64. Siegmund B, Schlack W, Ladilov YV. Halothane protects cardiomyocytes against reoxygenation-induced hypercontracture. *Circulation* 1997;96:4372-4379.
65. Schlack W, Preckel B, Barthel H. Halothane reduces reperfusion injury after regional ischaemia in the rabbit heart *in vivo*. *Br J Anaesth* 1997;79:88-96.
66. Preckel B, Schlack W, Comfere T. Effects of enflurane, isoflurane, sevoflurane and desflurane on reperfusion injury after regional myocardial ischaemia in the rabbit heart *in vivo*. *Br J Anaesth* 1998;81:905-912.
67. Schlack W, Preckel B, Stunneke D. Effects of halothane, enflurane, isoflurane, sevoflurane and desflurane on myocardial reperfusion injury in the isolated rat heart. *Br J Anaesth* 1998;81:913-919.
68. Varadarajan SG, An J, Novalija E. Sevoflurane before or after ischemia improves contractile and metabolic function while reducing myoplasmic Ca²⁺ loading in intact hearts. *Anesthesiology* 2002;96:125-133.
69. Heindl B, Reichle FM, Zahler S. Sevoflurane and isoflurane protect the reperfused guinea pig heart by reducing postischemic adhesion of polymorphonuclear neutrophils. *Anesthesiology* 1999;91:521-530.
70. Vinten-Johansen J. Involvement of neutrophils in the pathogenesis of lethal myocardial reperfusion injury. *Cardiovasc Res* 2004;61:481-497.
71. Kin H, Zatta AJ, Lofye MT, Amerson BS, Halkos ME, Kerendi F, Zhao ZQ, Guyton RA, Headrick JP, Vinten-Johansen J. Postconditioning reduces infarct size via adenosine receptor activation by endogenous adenosine. *Cardiovasc Res* 2005;67:124-133.
72. Gross ER, Hsu AK, Gross GJ. Opioid-induced cardioprotection occurs via glycogen synthase kinase β inhibition during reperfusion in intact rat hearts. *Circ Res* 2004;94:960-966.
73. Weihrauch D, Riotkowski JG, Bienengracher M, Hersten JR, Wartier DC, Pagel PS. Morphine enhances isoflurane-induced postconditioning against myocardial infarction: The role of phosphatidylinositol-3-kinase and opioid receptors in rabbits. *Anesth Analg* 2005;101:942-949.
74. Fryer RM, Wang Y, Hsu AK. Essential activation of PKC-delta in opioid-initiated cardioprotection. *Am J Physiol Heart Circ Physiol* 2001;280:H1346-H1353.
75. Ludwig LM, Patel HH, Gross GJ. Morphine enhances pharmacological preconditioning by isoflurane: Role of mitochondrial K (ATP) channels and opioid receptors. *Anesthesiology* 2003;98:705-711.
76. Luna-Ortiz P, Zarco-Olvera G, Ramírez-Ortega M, Tenorio-López FA, Gutierrez A, Martínez-Rosas M, Del-Valle-Mondragón L, Pastelin G. Antiarrhythmic and cardioprotective of remifentanyl in anesthetized dogs. *Arch Cardiol Mex* 2009;79:182-188.
77. Nishimura Y, Lemasters JJ. Glycine blocks opening of a death channel in cultured hepatic sinusoidal endothelial cells during chemical hypoxia. *Cell Death Differ* 2001;8:850-858.
78. Rajendra S, Lynch JW, Schofield PR. The glycine receptor. *Pharmacol Ther* 1997;73:121-146.
79. Laube B, Maksay G, Schemm R, Betz H. Modulation of glycine receptor function: a novel therapeutic approach for therapeutic intervention at inhibitory synapses? *Trends Pharmacol Sci* 2002;23:519-527.
80. Pfeiffer F, Betz H. Solubilization of the glycine receptor from rat spinal cord. *Brain Res* 1981;226:273-279.
81. Werman R, Davidoff RA, Aprison MH. Is glycine a neurotransmitter? *Nature* 1967;214:681-683.
82. Yamashima S, Konno A, Wheeler MD, Rusyn IV, Rusyn EL, Cox AD, Thurman RG. Endothelial cells contain a glycine-gated chloride channel. *Nutr Cancer* 2001;40:197-204.
83. Reeves WB. Effects of chloride channel blockers on hypoxic injury in rat proximal tubules. *Kidney Int* 1997;51:1529-1534.
84. Ikejima K, Limuro Y, Forman DT, Thurman RG. A diet containing glycine improves survival in endotoxic shock in the rat. *Am J Physiol* 1996;271:G-97-103.
85. Koerner JE, Anderson BA, Dage RC. Protection against postischemic myocardial dysfunction in anesthetized rabbits with scavengers of oxygen-derived free radicals. *J Cardiovasc Pharmacol* 1991;17:185-191.
86. Myers ML, Bolli R, Lekich RF, Hartley CJ, Roberts R. N-2mercaptopropionylglycine improves recovery of myocardial function after reversible regional ischemia. *J Am Coll Cardiol* 1986;8:1161-1168.
87. Tanaka M, Fujiwara H, Yamasaki K, Sasayama S. Superoxide dismutase and N-2 mercaptopropionylglycine attenuate infarct size limitation effect of ischemic preconditioning in the rabbit. *Cardiovasc Res* 1994;28:980-986.
88. Ruiz-Meana M, Pina P, Garcia Dorado D, Rodríguez-Sinovas A. Glycine protects cardiomyocytes against lethal reoxygenation injury by inhibiting mitochondrial permeability transition. *J Physiol* 2004;558:873-882.
89. Kin H, Zatta AJ, Lofye MT, et al. Postconditioning reduces infarct size via adenosine receptor activation by endogenous adenosine. *Cardiovasc Res* 2005;67:124-133.

90. Yang XM, Philipp S, Downey JM, Cohen MV. Postconditioning protection is not dependent on circulating blood factors. *Basic Res Cardiol* 2005;100:57-63.
91. Kerendi F, Kin H, Halkos ME, et al. Remote postconditioning: brief renal ischemia and reperfusion applied before coronary artery reperfusion reduces myocardial infarct size via endogenous activation of adenosine receptors. *Bas Res Cardiol* 2005;100:404-412.
92. Jin ZX, Zhou JJ, Xing M. Postconditioning the human heart with adenosine in heart valve replacement surgery. *Ann thorac Surg* 2007;83:2066-2073.
93. Lecour S. Multiple protective pathways against reperfusion injury: a SAFE path without Aktion? *J Mol Cell Cardiol* 2009;46:607-609.
94. Mann DL. Stress-activated cytokines and the heart: from adaptation to maladaptation. *Annu Rev Physiol* 2003;65:81-101.
95. Torre-Amione G, Kapadia S, Lee J, Bies RD, Lebovitz R, Mann DL. Expression and functional significance of tumor necrosis factor receptors in human myocardium. *Circulation* 1995;92:1487-1493.
96. Kurreimeyer KM, Michael LH, Baumgarten G, Taffet GE, Peschon JJ, Sivasubramanian N, et al. Endogenous tumor necrosis factor protects the adult cardiac myocyte against ischemic-induced apoptosis in a murine model of acute myocardial infarction. *Proc Natl Acad Sci USA* 2000;97:5456-5461.
97. Flaherty MP, Guo Y, Tiwari S, Rezazadeh A, Hunt G, Sangsaimath SK, et al. The role of TNF-alpha receptors p55 and p75 in acute myocardial ischemia/reperfusion injury and late preconditioning. *J Mol Cell Cardiol* 2008;45:735-741.
98. Lecour S, Suleman N, Deuchar GA, Somers S, Lacerda L, Huisamen B, et al. Pharmacological preconditioning with tumor necrosis factor-alpha activates signal transducer and activator of transcription-3 at reperfusion without involving classic prosurvival kinases (Akt and extracellular signal-regulated kinase). *Circulation* 2005;112:3911-3918.
99. Boengler K, Hilfiker-Kleiner D, Drexler H, Heusch G, Schulz R. The myocardial JAK/STAT pathway: from protection to failure. *Pharmacol Ther* 2008;120:172-185.
100. Gross ER, Hsu AK, Gross GJ. The JAK/STAT pathway is essential for opioid-induced cardioprotection: JAK2 as a mediator of STAT3, Akt, and GSK-3 beta. *Am J Physiol Heart Circ Physiol* 2006;291:H827-834.