

Social connections and depressive symptoms among older adults during the initial lockdown period for COVID-19

Sara Estefania Solis-López,¹ Virgilio Hernández-Ruiz,² Hélène Amieva,² Jose Alberto Avila-Funes^{1,2}

¹ Departamento de Geriátrica, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Ciudad de México, México.

² Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR 1219, F-33000 Bordeaux, France.

Correspondence:

José Alberto Avila-Funes
Departamento de Geriátrica, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán.
Vasco de Quiroga 15,
C.P. 14080, Tlalpan,
Ciudad de México, México.
Phone: +52 (55) 5487 0900,
ext. 5703
Email: avilafunes@live.com.mx

Received: 20 May 2022
Accepted: 21 October 2022

Citation:

Solis-López, S. E., Hernández-Ruiz, V., Amieva, H., & Avila-Funes, J. A. (2023). Social connections and depressive symptoms among older adults during the initial lockdown period for COVID-19. *Salud Mental*, 46(3), 131-136.

DOI: 10.17711/SM.0185-3325.2023.017



ABSTRACT

Introduction. Little information from developing countries during the first lockdown for COVID-19 is available. We hypothesized that the use of communication tools, and not living alone would provide a protective effect against DS. **Objective.** To determine the association between social connections and depressive symptoms (DS) during the first lockdown period among Mexican community-dwelling older adults. **Method.** Cross-sectional phone survey including 269 participants aged 65 years or older. Participants were asked about their social connections and the presence of DS during the first lockdown for COVID-19. **Results.** Mean age was 83.2 (SD = 6.7). Compared with those without DS, those with DS reported a greater number of phone calls or videocalls although this was not statistically significant. However, when stratifying by housing situation, only the participants who lived alone and that received less calls from friends had more DS ($p = .04$). **Discussion and conclusion.** Living alone allowed the participants not to have much contact with family and this caused friends to represent the most important social relationship outside the home. We hypothesize that the means to stay socially active for older adults in Latin America are different and have a different impact. Because, DS only were present among the participants who reported living alone and having fewer calls from friends during the confinement period.

Keywords: Depressive symptoms, COVID-19, social connections, older adults.pp

RESUMEN

Introducción. Hay poca información disponible sobre los países en desarrollo durante el primer período de confinamiento por COVID-19. Planteamos la hipótesis de que el uso de herramientas de telecomunicación y vivir acompañado proporciona un efecto protector frente a la presencia de síntomas depresivos (SD). **Objetivo.** Determinar la asociación entre las conexiones sociales y los SD durante el primer período de confinamiento en adultos mayores mexicanos que viven en la comunidad. **Método.** A través de un estudio transversal, 269 participantes de 65 años o más completaron una encuesta telefónica sobre sus conexiones sociales y la presencia de SD durante el primer período de confinamiento por COVID-19. **Resultados.** La edad media fue de 83.2 (DE = 6.7). En comparación con los que no tenían SD, los que tenían SD reportaron un mayor número de llamadas telefónicas o videollamadas, pero esto no fue estadísticamente significativo. Sin embargo, al estratificar por situación de vivienda, los participantes que vivían solos y que recibían menos llamadas de amigos tenían más SD ($p = .04$). **Discusión y conclusión.** Vivir solo permitió a los participantes no tener mucho contacto con la familia y esto provocó que los amigos representaran la relación social más importante fuera del hogar. Creemos que los medios para mantenerse socialmente activos de los adultos mayores en América Latina son diferentes y tienen un impacto diferente. Debido a que los SD solo estuvieron presentes entre los participantes que reportaron vivir solos y tener menos llamadas de amigos durante período de confinamiento.

Palabras clave: Síntomas depresivos, COVID-19, conexiones sociales, adultos mayores.

INTRODUCTION

Social isolation is a known risk factor for diverse negative health-related and mental health-related outcomes among older adults. These outcomes include the development of depression and depressive symptoms (DS). This risk was recognized well before the COVID-19 pandemic (Cipolleta & Gris, 2021; Santini et al., 2020). Therefore, as the World Health Organization warned, mental health-related complications associated with social isolation could increase in this already vulnerable population due to the adoption of the lockdown periods to limit the spread of SARS-CoV-2 at the beginning of the pandemic (World Health Organization [WHO], 2020). In this sense, older adults are not only at higher risk for increased mortality from COVID-19, but are also vulnerable for its indirect adverse mental health effects, such as the development of DS (Fraser et al., 2020).

Consequently, during the COVID-19 pandemic, multiple projects have been conducted in order to study the impact that social isolation may have had on older adults' mental health during diverse lockdown periods. In the pre-pandemic era, several studies have already shown an association between loneliness and social isolation and various adverse health-related outcomes among older adults including all-cause mortality, poor physical health (Noone et al., 2020), and increased risk for dementia (Cipolleta & Gris, 2021; Santini et al., 2020).

However, results coming from the early months of the pandemic and the first lockdown periods have been very heterogeneous, either because they did not observe changes in mental health during that period (Röhr, Reininghaus, & Riedel-Heller, 2020), or because they reported a worsening of DS (Robb et al., 2020). As the future of the pandemic remains uncertain, it is pertinent to continue studying the possible associations between isolation and its negative mental health-related consequences in older adults, such as the development of DS. Equally important is the need to study variables that could mitigate these consequences, such as maintaining social interactions (i.e., social connections). Social connections are the relationships that we have with the people around us and are shaped by verbal and non-verbal performances that co-construct the meaning of such interactions, either in person or remotely by using technology (Birt et al., 2020). For instance, studies conducted in high-income countries before the pandemic have reported that living with other people and maintaining remote contact through communication tools, such as telephone calls or videocalls, had a positive impact on the reduction of DS (Noone et al., 2020; Stevic, Schmuck, Matthes, & Karsay, 2021). However, little information from developing countries is available concerning the potential effect that the first lockdown period may have had on older adults' mental health (Buenaventura, Ho, & Lapid, 2020). In Mexico, in order to limit the number of infections by SARS-CoV-2,

between March 23 and May 30, 2020, strict social distancing (domiciliary lockdown), as well as the suspension of all non-essential activities were recommended by the Government (Ibarra-Nava, Cardenas-de la Garza, Ruiz-Lozano, & Salazar-Montalvo, 2020). Relying on the PAMCOVID telephone survey aimed at communitarian older adults, and administered during the first lockdown period in Mexico (Hernández-Ruiz, 2021), the objective of the present study was to determine the association between social connections – assessed by the housing situation (if they lived alone or in the company of someone else) and the use of communication tools (phone-calls or video-calls) to maintain contact with family, friends, and neighbors during the first lockdown period in Mexico City – and the presence of DS among community-dwelling older adults.

METHOD

Participants

Cross-sectional study including the 269 participants from the PAMCOVID telephone survey. Briefly, the PAMCOVID survey is the Mexican counterpart of the French PACOVID (“Personnes Agées face au COVID”) longitudinal telephone survey. These surveys were set up in the region of Bordeaux (France), and Mexico City a few days after the first global lockdown. Both were aimed at communitarian older adults and administered by trained health staff. Both surveys shared the initial main objective of addressing the issue; what are the attitudes, psychological, and social experiences of the older persons regarding the COVID-19 crisis and the lockdown measures? The complete methodology and initial results for both surveys have been described elsewhere (Hernández-Ruiz, 2021).

The collected data included information on the housing situation during the lockdown (i.e., if participants were living alone or with someone else), mental-health (three items from the CES-D questionnaire: how often have they felt “sad”, “depressed”, and “lonely” over the last week; Radloff, 1977), health status (including self-reported comorbidities), and social variables (non-professional phone or video contact over the last week).

For the present study, the included population consisted in older persons (≥ 65 years-old) from a Geriatrics outpatient clinic at a third-level university-affiliated hospital in Mexico City whose consultation was canceled during the first national mandatory lockdown. Thus, eligible participants were contacted, and asked if they wished to participate in a 30-minute phone interview. Data was collected in May-July 2020. At the beginning of the interview the objectives of the study were presented, and verbal informed consent was obtained. If the persons themselves could not respond to the questionnaire but wished to participate, a proxy was invited

to answer in their place. Exclusion criteria were refusal to participate in the study, inability to answer the interview due to cognitive impairment or severe hearing problems as well as the absence of a proxy informant who could answer for them if needed.

Measurement

Depressive symptoms

DS were established by three items from the Center for Epidemiological Studies depression (CES-D) scale (*how often in the past week have you felt sad / how often in the past week have you felt depressed, and how often in the past week have you felt lonely*). Each item was scored on a four-point scale, and then it was dichotomized in “no” (not at all) or “yes” (rarely or less than a day, some of the time 1-2 days, occasionally 3-4 days or most of the time 5-7 days), being “yes” if in the past week they had felt these feelings at least one day. If the participants answered “yes” for at least one question, they were considered as having DS. If the participants reported DS they were oriented to receive the correspondent attention and also were referred to the hotline for older adults that was implemented by the Geriatrics clinic (Navarrete-Reyes & Avila-Funes, 2020).

Social connections

Several measures of social connections were included. Participants were asked about their housing condition during the lockdown period: if they lived alone or in the company of a partner/another member of the family. Likewise, they were asked if they have had remote contact via communication tools (phone calls or videocalls) with other persons (family members, friends and/or neighbors). Finally, participants were asked the number of phone calls or videocalls received on the seven days preceding the interview, and if that number represented a lower / same / higher quantity of calls respect to *the usual* (previous to the lockdown period).

Other variables

We obtained demographic data (age, sex, and level of education) from the medical records. Health conditions included self-reported diagnoses of hypertension, diabetes, or chronic lung disease.

Statistical analysis

Variables were described using arithmetic means and standard deviations (SD) or frequencies and proportions when appropriate. The following statistical procedures were used according to the characteristic of each variable: chi square test for qualitative data or Student t test for continuous data. In order to test the association between every measure of social connection and the presence of DS, univariate logistic regression models were conducted. All statistical tests were

evaluated using 95% confidence intervals (CI) and p value < .05. Statistical tests were performed using the SPSS software for Windows® (SPSS Inc., Chicago, IL, version 25).

Ethical considerations

The patients/participants provided their informed consent to participate in this study. PAMCOVID ethics approval number 3361 (CONBIOÉTICA-09-CEI-011-2016027).

RESULTS

From 269 participants, the 71.7% ($n = 193$) of the responses were directly obtained from participants themselves and the 28.2% ($n = 76$) from a proxy. Mean age of participants was 83.2 (SD = 6.5; range 67-100), and 72.5% were women. The most prevalent chronic diseases were hypertension 66.5% ($n = 179$) and diabetes ($n = 96$; 35.7%; Table 1).

Regarding the housing situation during the lockdown period, most of the participants lived with a family member ($n = 195$; 72.5%). The most used means to keep social interactions were phone calls ($n = 243$; 90.3%) with an average of 29 (SD = 30.6) calls received from family members, friends and/or neighbors, in the seven days prior to the interview. And most of the participants perceived having received a greater number of calls ($n = 231$; 85.9%), being the calls from family members the ones who reported a greater perceived increase ($n = 223$; 94.9%), followed by the calls from friends ($n = 95$; 87.2%) and neighbors ($n = 28$; 78.5%). On the other hand, the use of videocalls was the less frequently used communication tool ($n = 117$; 43.5%). Among the participants that reported videocall use, an average of 4.2 (SD = 6.0) videocalls were received in the week before the interview from family members, friends and/or neighbors.

Half of the participants ($n = 146$; 54.3%) reported the presence of at least one depressive symptom in the week before the interview and “*feeling sad*” was the most frequently reported answer ($n = 130$; 48.3%). Compared with those who did not have DS, participants who reported DS were younger ($p = .04$), were predominantly women ($p = .02$), and more frequently reported living alone ($p < .001$). Despite that a higher number of received phone calls from their families, friends and/or neighbors was reported by participants with DS respect to their counterparts without DS, the difference was not statistically significant ($p = .23$). In the same vein, participants who had DS more frequently reported that the number of phone calls received from family, friends and/or neighbors during the lockdown was higher if compared to the pre-pandemic period; however, this difference was not statistically significant either ($p = .39$). In the same vein, the univariate logistic regression analysis showed that the number of phone-calls from friends (OR = .90, 95% CI = [.54,

Table 1
Sociodemographic and mental health characteristics

	All study sample (n = 269)	Without DS (n = 123)	With DS (n = 146)	p
Age, mean (SD)	83.2 (6.5)	83.4 (6.8)	82.8 (6.3)	.04
Female, n (%)	195 (72.5)	81 (65.9)	114 (78.1)	.02
Health conditions:				
Hypertension, n (%)	179 (66.5)	79 (44.1)	100 (55.9)	.52
Diabetes mellitus, n (%)	96 (35.7)	50 (52.1)	46 (47.9)	.56
Pneumopathy, n (%)	52 (19.3)	20 (38.5)	32 (61.5)	.25
Cancer, n (%)	30 (11.2)	14 (46.7)	16 (53.3)	.89
Housing condition:				
Living with family, n (%)	195 (72.5)	94 (76.4)	101 (70.1)	.25
Living alone, n (%)	20 (7.4)	1 (0.8)	19 (13.2)	< .001
Number of phone contacts received, mean (SD)	29 (30.6)	28 (30.2)	30.7 (31.3)	.23
Number of videocalls contacts received, mean (SD)	4.2 (6.0)	4.2 (6.9)	4.2 (5.0)	.86
More calls perceived from family, n (%)	223 (94.9)	108 (96.4)	115 (93.5)	.31
More calls perceived from friends, n (%)	95 (87.2)	45 (86.5)	50 (87.7)	.85
More calls perceived from neighbors, n (%)	28 (75.7)	14 (73.7)	14 (77.8)	.77

Note: DS = depressive symptoms.

1.4], $p = .68$), family members (OR = .51, 95% CI = [.26, 1.00], $p = .52$) and neighbors (OR = .83, 95% CI [.37, 1.8], $p = .63$) was not statistically significant associated with the presence of DS. However, stratifying by housing situation (living alone vs. those who lived in the company of another person), only those who were living alone and had a lower number of phone-calls from friends presented more DS ($p = .04$; Table 2).

DISCUSSION AND CONCLUSION

This study did not find an association between keeping social connections and the presence of DS in community-dwelling older adults. And only when stratifying by housing situation, we found that those who were living alone and had lower number of phone-calls from friends presented more DS.

During the first waves of the COVID-19 pandemic, restrictive measures such as lockdowns and physical distancing were implemented in order to protect the population, particularly the more vulnerable groups as the older adults. However, it was warned that these restrictions could become a risk factor for the development of psychological problems such as DS (Cipolleta & Gris, 2021). That is why we hypothesized that the use of communication tools, like phone calls or video-calls, and not living alone (i.e., social connections) would provide a protective effect against DS (Käll et al., 2020). Nevertheless, this study does not prove it.

Even if our results are in the same line with existing literature reporting that populations with higher risks of developing DS during the first lockdown period were younger and living alone, we failed to show a statistical and significant association between the number of calls, as a proxy for social connection, and the presence of DS. For instance, Robb and collaborators showed that in 7,127 English community-dwelling older adults, women, younger persons, those single/widowed/divorced and those who were living alone more frequently reported a worsening of DS during the first lockdown (Robb et al., 2020). However, the evidence showing that the initial lockdown did not have the impact that we might have thought is increasing. For instance, in the cohort of Röhr and collaborators consisting of 1,005 community dwelling German older adults (aged 65-94) who were being followed for depression and anxiety prior to the pandemic, the authors report that mental health-related measures remained largely unchanged during the first lockdown period. This being assessed by a computer-assisted standardized telephone interview (Röhr et al., 2020). In the same vein, van Tilburg et al. found that despite an increase in the feeling of loneliness in older people, depression and anxiety feelings remained roughly stable during the first lockdown. This was evaluated in 1,679 Dutch community-dwelling persons aged 65-102 who completed a digital survey of loneliness and mental health (van Tilburg, Steinmetz, Stolte, van der Roest, & de Vries, 2021). It is possible that this can be explained by older adults' ability to apply coping strategies, like engaging in leisure activities, maintaining their daily routine, or by the

Table 2
Comparative analysis of depressive symptoms according calls number and housing condition

<i>Living with family</i>	<i>Calls from friends</i>	<i>With DS</i>	<i>Without DS</i>	<i>p</i>
No	No, <i>n</i> (%)	0 (0)	5 (19.2)	.04
	Yes, <i>n</i> (%)	19 (100)	21 (80.8)	
Yes	No, <i>n</i> (%)	7 (21.2)	2 (6.5)	.09
	Yes, <i>n</i> (%)	26 (78.8)	29 (93.5)	
<i>Calls from family</i>				
No	No, <i>n</i> (%)	0 (0)	4 (10)	.09
	Yes, <i>n</i> (%)	27 (100)	36 (90)	
Yes	No, <i>n</i> (%)	4 (4.7)	4 (4.9)	.94
	Yes, <i>n</i> (%)	81 (95.3)	77 (95.1)	
<i>Calls from neighbors</i>				
No	No, <i>n</i> (%)	0 (0)	2 (22.2)	.30
	Yes, <i>n</i> (%)	4 (100)	7 (77.8)	
Yes	No, <i>n</i> (%)	5 (33.3)	2 (22.2)	.56

Note: DS = depressive symptoms.

acceptance of the lockdown situation, as a way of facing these new circumstances. Likewise, they may display lower stress reactivity and more emotional resources to adapt in better ways to face adverse situations, as suggested by the French PACOVID study (Hernández-Ruiz et al., 2021).

It is important to recall that most of the evidence recommending the use of communication tools to maintain remote social interactions is based on studies that only found a positive impact of these tools in younger populations. However, a multi-national cross-sectional online survey by Bonsaksen and collaborators conducted in community-dwelling older adults during the first months of the pandemic, reported that only the participants between 60-69 years endorsed a reduction in the feeling of loneliness and an increase in quality-of-life estimators with the use of video-based communication (Bonsaksen et al., 2021). The mean age of our sample was higher in comparison to studies that recommend the use of newer communication tools, which may influence why in our sample only 43% of participants used video-calls. Nevertheless, this proportion is larger than the reported prevalence of use in other populations, which may reflect the characteristics of the Latin-American housing, where intergenerational cohabitation is more frequent (Kinsella, Velkoff, & U.S. Census Bureau, 2002). Another factor to account is that most of the studies evaluating older adults' mental health during the different lockdown periods have not been performed in Latin-American countries, where older adults probably experienced different impacts of the pandemic respect to their counterparts in high-income countries. Likewise, the means to keep socially active or the social connections adopted

by older adults in Latin America could have been different, as suggested by the study of Tyler and collaborators. The work examined the relationships amongst demographics, COVID-19 life impacts, depression and anxiety in adults aged 60 or older from 33 countries (Tyler et al., 2021). This study found that in the first months of the pandemic being separated from, and having conflicts with loved ones predicted depression, as did residing in a country with higher income. In our study, most of the participants kept their family and friends close, and most of the participants lived in the company of someone else, who most likely acted like their social connections physically and helped them to maintain contact out of the household. This may be supported by the finding that in our study we only found DS among those participants who lived alone and did receive fewer calls from friends. Which may be a representation of other social constraints in these individuals. We hypothesize that living alone constrained them to not have much contact with their family members, and consequently their friends represented their most important social relationship outside of the household. Indeed, in Geriatrics it is well recognized that living alone and being socially isolated are two factors that can have a significant impact on older adults' mental health (Tanskanen & Anttila, 2016).

Some of our study's limitations are the lack of a previous mental health status assessment, its cross-sectional design, its sample size and that we didn't include participants who did not have access to phone-calls. However, several strengths can be underlined. This study is the first-one that has sought to determine whether maintaining social connections during the first lockdown either by household status,

phone or video-calls had an association with the presence of DS in a middle-income country. Moreover, by relying on the telephone to do our survey, it was possible to achieve the participation of people in more vulnerable states of health, or who do not have the electronic means to answer an online survey. Which also contributes to reduce the selection bias, as described in other publications (Bethlehem, 2010). However, we are not exempt from other selection biases.

In conclusion, the present study showed that relatively younger participants, women, and people that were living alone and receive fewer calls from friends, more frequently reported depressive symptoms during the initial COVID-19 lockdown period in Mexico City. Social connections did not had associations with the presences of DS. This may reflect the impact of in person social interactions in the people who lived in the company of someone else. The results of this study offer an insight at some of the conditions that the older adults from Mexico City experienced during the first lockdown period of the COVID-19 pandemic, and thus, may promote the creation of tools to address the situation of those individuals who may have fewer social connections.

Funding

None.

Conflict of interest

The authors declare they have no conflicts of interest.

REFERENCES

- Bethlehem, J. (2010). Selection bias in web surveys. *International Statistical Review*, 78(2), 161-188. doi: 10.1111/j.1751-5823.2010.00112.x
- Birt, L., Griffiths, R., Charlesworth, G., Higgs, P., Orrell, M., Leung, P., & Poland, F. (2020). Maintaining Social Connections in Dementia: A Qualitative Synthesis. *Qualitative Health Research*, 30(1), 23-42. doi: 10.1177/1049732319874782
- Bonsaksen, T., Thygesen, H., Leung, J., Ruffolo, M., Schoultz, M., Price, D., & Østertun Geirdal, A. (2021). Video-Based Communication and Its Association with Loneliness, Mental Health and Quality of Life among Older People during the COVID-19 Outbreak. *International Journal of Environmental Research and Public Health*, 18(12), 6284. doi: 10.3390/ijerph18126284
- Buenaventura, R. D., Ho, J. B., & Lapid, M. I. (2020). COVID-19 and mental health of older adults in the Philippines: a perspective from a developing country. *International Psychogeriatrics*, 32(10), 1129-1133. doi: 10.1017/S1041610220000757
- Cipolletta, S., & Gris, F. (2021). Older People's Lived Perspectives of Social Isolation during the First Wave of the COVID-19 Pandemic in Italy. *International Journal of Environmental Research and Public Health*, 18(22), 11832. doi: 10.3390/ijerph182211832
- Fraser, S., Lagacé, M., Bongué, B., Ndeye, N., Guyot, J., Bechard, L., ... Tougas, F. (2020). Ageism and COVID-19: what does our society's response say about us?. *Age Ageing*, 49(5), 692-695. doi: 10.1093/ageing/afaa097
- Hernández-Ruiz, V., Meillon, C., Avila-Funes, J.-A., Bergua, V., Dartigues, J.-F., Koleck, M., ... Amieva, H. (2021). Older Adults and the COVID-19 Pandemic, What About the Oldest Old? The PACOVID Population-Based Survey. *Frontiers in Psychiatry*, 12, 711583. doi: 10.3389/fpsy.2021.711583
- Ibarra-Nava, I., Cardenas-de la Garza, J. A., Ruiz-Lozano, R. E., & Salazar-Montalvo, R. G. (2020). Mexico and the COVID-19 Response. *Disaster Medicine and Public Health Preparedness*, 14(4), e17-e18. doi: 10.1017/dmp.2020.260
- Kinsella, K., Velkoff, V. A., & U.S. Census Bureau, (2002). Living arrangements. *Aging Clinical and Experimental Research*, 14(6), 431-438. doi: 10.1007/bf03327344
- Käll, A., Jägholm, S., Hesser, H., Andersson, F., Mathaldi, A., Norkvist, B. T., ... Andersson, G. (2020). Internet-Based Cognitive Behavior Therapy for Loneliness: A Pilot Randomized Controlled Trial. *Behavior Therapy*, 51(1), 54-68. doi: 10.1016/j.beth.2019.05.001
- Navarrete-Reyes, A. P., & Avila-Funes, J. A. (2020). Staying in a Burning House: Perks and Perils of a Hotline in the Times of COVID-19. *Journal of the American Geriatrics Society*, 68(5), E10-E11. doi: 10.1111/jgs.16479
- Noone, C., McSharry, J., Smalle, M., Burns, A., Dwan, K., Devane, D., & Morrissey, E. C. (2020). Video calls for reducing social isolation and loneliness in older people: a rapid review. *Cochrane Database of Systematic Reviews*, 5, (5), CD013632. doi: 10.1002/14651858.CD013632
- Radloff, L. S. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1(3), 385-401. doi: 10.1177/014662167700100306
- Robb, C. E., Jager, C. A., Ahmadi-Abhari, S., Giannakopoulou, P., Udeh-Momoh, C., McKean, J., ... Middleton, L. (2020). Associations of Social Isolation with Anxiety and Depression During the Early COVID-19 Pandemic: A Survey of Older Adults in London, UK. *Frontiers in Psychiatry*, 11, 591120. doi: 10.3389/fpsy.2020.591120
- Röhr, S., Reininghaus, U., & Riedel-Heller, S. G. (2020). Mental wellbeing in the German old age population largely unaltered during COVID-19 lockdown: results of a representative survey. *BMC Geriatrics*, 20(1),489. doi: 10.1186/s12877-020-01889-x
- Santini, Z. I., Jose, P. E., Cornwell, E. Y., Koyanagi, A., Nielsen, L., Hinrichsen, C., Meilstrup, C., ... Koushede, V. (2020). Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal mediation analysis. *Lancet Public Health*, 5(1), e62-e70. doi: 10.1016/S2468-2667(19)30230-0
- Stevic, A., Schmuck, D., Matthes, J., & Karsay, K. (2021). 'Age Matters': A panel study investigating the influence of communicative and passive smartphone use on well-being. *Behaviour & Information Technology*, 40(2), 176-190. doi: 10.1080/0144929X.2019.1680732
- Tanskanen, J., & Anttila, T. (2016). A Prospective Study of Social Isolation, Loneliness, and Mortality in Finland. *American Journal of Public Health*, 106(11), 2042-2048. doi: 10.2105/AJPH.2016.303431
- Tyler, C. M., McKee, G. B., Alzueta, E., Perrin, P. B., Kingsley, K., Baker, F. C., & Arango-Lasprilla, J. C. (2021). A Study of Older Adults' Mental Health across 33 Countries during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 18(10), 5090. doi: 10.3390/ijerph18105090
- van Tilburg, T. G., Steinmetz, S., Stolte, E., van der Roest, H., & de Vries, D. H. (2021). Loneliness and mental health during the COVID-19 pandemic: A study among Dutch older adults. *The Journals of Gerontology*, 76(7), e249-e255. doi: 10.1093/geronb/gbaa111
- World Health Organization. (2020). *Mental health and psychosocial considerations during the COVID-19 outbreak*. World Health Organization.