



Facultad de Estudios Superiores  
**IZTACALA**

# Artificial neural networks model: Neuropsychological variables and their relationship with body fat percentage in adults

## *Modelo de redes neuronales artificiales: Variables neuropsicológicas y su relación con el porcentaje de grasa corporal en adultos*

Bárbara Itzel Méndez-Peña<sup>1</sup>, María Magdalena Murillo-Tovar<sup>1</sup>, Gerardo Leija-Alva<sup>1</sup>,  
Itzihuari Iratzi Montufar Burgos<sup>1</sup>, Angélica Serena-Alvarado<sup>1</sup>, Roxana Sarai Durán-Arciniega<sup>1</sup>,  
Nadia Mabel Pérez-Vielma<sup>1</sup>, Víctor Ricardo Aguilera-Sosa<sup>1,2</sup>

<sup>1</sup>Centro Interdisciplinario de Ciencias de la Salud Unidad Santo Tomás, Instituto Politécnico Nacional. 11340. CDMX.

<sup>2</sup>Departamento de Investigación y Posgrado

Recibido: 26-11-2021

Revisado: 28-12-2021

Aceptado: 14-01-2022

**Autor de correspondencia:** [vaguilera@ipn.mx](mailto:vaguilera@ipn.mx) (V. R. Aguilera-Sosa)

**Funding.:** This work received no funding.

**Conflict of Interest:** Authors declare no conflicts of interest.

### Abstract

There is a growing interest to understand the neural functions and substrates of complex cognitive processes related to Obesity (OB). Artificial Intelligence (AI) is being applied, specifically the perceptron model of Artificial Neural Networks (ANN) in non-communicable chronic diseases, to identify with greater certainty the connective factors (synaptic networks) between the input variables and the output variables associated. **Objective:** Identify the synaptic weights of the ANN whose input variables are the executive functions (EF) and healthy lifestyles as predictors of Body Fat Percentage (BFP) in a group of adult subjects with different levels of BFP. **Methods:** It was an exploratory, quantitative, cross-sectional, comparative, convenience, and explanatory research. The Neuropsychological Battery (BANFE-2) and the Overeating Questionnaire (OQ) were administered

to 40 participants aged between 18-38 years. BFP was measured using a RENPHO ES-24M Smart Body Composition Scale. The perceptron ANN model with ten trials was applied with a multilayer-perceptron. **Results:** The ANN showed that the sensory variables with greater synaptic weight for BFP were Stroop A and B Errors and Successes of BANFE-2, and OQ scales Rationalizations and Healthy Habits. **Conclusions:** ANN proved to be important in the simultaneous analysis of neuropsychological and healthy lifestyle data for the analysis of OB prevention and treatment by identifying the variables that are closely related. These findings open the door for the use of non-linear analysis models, which allow the identification of relationships of different weights, between input and output variables, to more effectively direct interventions to modify obesity habits.

Keywords: healthy habits, neuropsychological variables, body fat, artificial neural networks.

## Resumen

Existe un interés creciente por comprender las funciones neuronales y sustratos cognitivos complejos relacionados con la obesidad. Se está aplicando Inteligencia Artificial, en concreto el modelo perceptrón de Redes Neuronales Artificiales en enfermedades crónicas no transmisibles, para identificar con mayor certeza los factores de conexión (redes sinápticas) entre las variables de entrada y las variables de salida. **Objetivo:** Identificar pesos sinápticos de la RNA cuyas variables de entrada fueron las funciones ejecutivas y los estilos de vida saludable, como predictores del Porcentaje de Grasa Corporal en un grupo de sujetos adultos con diferentes niveles del Porcentaje de Grasa. **Métodos:** se trató de una investigación exploratoria, cuantitativa, transversal, comparativa, de conveniencia y explicativa. Se administró la Batería Neuropsicológica (BANFE-2) y el Cuestionario de Sobreingesta (OQ), a 40 participantes con edades comprendidas entre los 18-38 años. El porcentaje de grasa se midió con una báscula de composición corporal (RENPHO ES-24M). El modelo redes neuronales de perceptrón, se ejecutó con diez ensayos. **Resultados:** El modelo de Red Neuronal mostró que las variables sensoriales con mayor peso sináptico para el porcentaje de grasa, fueron Errores Stroop A y B y Aciertos de BANFE-2, y Racionalizaciones de las escalas OQ y Hábitos Saludables. **Conclusiones:** las redes neuronales artificiales demostró ser importante en el análisis simultáneo de datos neuropsicológicos y de estilo de vida saludable para el análisis de prevención y tratamiento de la obesidad, al identificar las variables que están estrechamente relacionadas. Estos hallazgos abren la puerta al uso de modelos de análisis no lineales, que permiten identificar relaciones de diferente peso, entre variables de entrada y salida, más eficientes que los modelos lineales.

Palabras clave: hábitos saludables, variables neuropsicológicas, grasa corporal, redes neuronales artificiales.

## Introduction

Understanding OB requires the development of models that facilitate the analysis, prognosis, and recognition of its behavior, all which enables us to address it (WHO, 2005). One proposal for model generation used in Artificial Intelligence (AI) is the Perceptron Model of Artificial Neural Networks (ANN). It has proved

to be useful for a broad understanding of the factors that influence the relationship between the input, the hidden layers (synapsis) and the output or dependent variables. Multilayer perceptron (MLP) is a nonlinear interaction model of the relationship between variables with a high level of connectivity. Backpropagation allows for networks to learn based on least squares error correction, and its main objective is that the networks

learn to identify patterns and generate categories of information (García & Espinosa, 2013). This process is as follows: among  $n$  cases, the ANN algorithm randomly chooses some and excludes others until it finds, depending on the number of trials, the model that best fits the input and output variables. In this way, the use of the results of ANN models enables health professionals to make more specific and effective decisions for interventions; influencing well-being and life quality of patients (Bajo & Ballesteros, 2002; Arias et al. 2019; Sarmiento-Ramos, 2020).

From a neuropsychological perspective, there is a growing interest to understand the neural functions and substrates of complex cognitive processes related to OB, particularly executive functions (EF), which are supported in planning, self-monitoring, inhibition, and decision making performed daily (Allan et al., 2016; Tirapu et al., 2012).

It has been suggested that obese individuals with a sedentary lifestyle show low results in inhibitory control and working memory tasks, as well as a lower neurocognitive performance (Tsai, 2019) correlative to BFP, which has a negative impact on performance of cognitive control tasks, (Huang et al., 2019; Yang et al., 2018) while in individuals with a healthy weight executive functioning is not impaired (Calvo et al., 2014; Narimani et al., 2019).

OB entails a greater accumulation of body fat and its redistribution into adipose tissue for storage (Carlson, 2014; Brandan et al., 2014). This accumulation results in an energy imbalance which is reflected mainly in weight gain and waist and hip circumference (Carlson, 2014; Cherbuin & Walsh, 2019). This is the reason why BFP is a more effective measure at predicting health risks than the simple measurement of gross weight or Body Mass Index (BMI) (Calvo et al., 2014; Huang et al., 2019; Narimani et al., 2019).

The use of ANNs has been helpful in finding the relation between environment and the prevalence of OB. In a study conducted in the USA, AI was used to identify more than 150, 000 satellite images and establish links with OB prevalence and health indicators, such as the level of physical activity of people. The final algorithm found, indeed, a relation between high population density, a high number of expressways, fewer parks and green spaces and high prevalence of OB (Cardozo

et al., 2016). Heydari, Ayatollahi & Zare compared ANN and binary logistic regression to identify degree of obesity in a group of adults with different degrees of fat percentage. They found that both models were efficient in identifying the degree of obesity (2012). In another study, Ergün (2009) based on the need to have an automated system for the recognition and monitoring of obesity, also compared a multilayer neural network model against a binary logistic regression model, concluding that the neural network is better than the logistic one.

Therefore, the objective of this study was to identify the synaptic weights of the ANN whose input variables are the executive functions (EF) and health lifestyles as predictors of Body Fat Percentage (BFP) in a group of adult subjects with low, normal, high, and very high body fat levels.

## Material and Methods

This was a non-experimental, cross-sectional, explanatory, quantitative with a non-probabilistic design, convenience sampling,  $n = 40$  adults aged between 18-38 years old. All the participants were residents of Azcapotzalco, Miguel Hidalgo and Cuauhtémoc, mayors of Mexico City, and were evaluated from November 2019 to February 2020.

### *Instruments and Measurements*

#### *OQ, Overeating Questionnaire.*

It is a self-administered instrument with a five-point Likert-type scale (Not at all, A little, Moderately, A lot and very much) used to evaluate habits, thoughts and attitudes related to overweight and OB. It contains 80 items that measure cognitions, behavioral cognitions, behaviors-habits, emotions, and attitudes related to obesity. For its construction, it was applied to an  $n=1,788$  North American individuals between the ages of 9 and 98 years old. Two areas were found: one for Inconsistency (INC) and one for Defensiveness (DEF). Eight scales for Habits and Attitudes: Overeating (SOB), Undereating (SUB), Cravings (CRAV) and (SUB), Food cravings (ANT), Expectations with eating

(EXP), Rationalizations (RAC), and Motivation to lose weight (MOT). The latter scales assess variables related to general habits and psychosocial functioning: Habits (Hábitos), Eating Habits (HAB), Eating Expectations (EXP) psychosocial functioning: Health Habits (SAL), Body Image (COR), Social Isolation (AIS), and (AIS) and Affective Disturbances (AFE). It has an internal consistency of 0.94 to 0.88, and correlates with BMI scores and with other scales that measure health habits, social functioning and social functioning. The authors report a reliability of 0.82 (Cronbach's alpha) and construct validity on subscales of Food Cravings and Motivation of 0.26, for Food Cravings and Motivation; 0.68 between Food and Motivation; 0.68 between Social Isolation and Affective Disturbance; Health Habits and Food Intake 0.53 (William and Warren, 2007). In 2014 a factor analysis was conducted which yielded 9 factors, which were: social isolation, stress, body image, depression, health habits, expectations related to eating, overeating, motivation to lose weight and weight concern, all of them integrated in 30 items (Psihas, 2014). The questionnaire had acceptable validity and reliability ( $\geq .76$ ) in the Mexican sample, however the sample was very small compared to the original one, so it was decided to use the latter in the present investigation.

### **Neuropsychological battery of executive functions and frontal lobes (BANFE -2).**

The 15 sub-tests that integrate the battery are divided based on anatomical-functional criteria that evaluate complex functions of the orbitofrontal cortex (OFC), prefrontal cortex (PFC), through the sub-tests: stroop effect, card game and mazes; of the dorsolateral prefrontal cortex (DLPFC) through: self-directed signaling, spatial working memory, alphabetical word sorting, card sorting, maze and Hanoi tower; and of the anterior prefrontal cortex (aPFC) through: semantic classification, choice of sayings and metamemory. The qualification is carried out through the quantitative and qualitative analysis of successes and errors, based on the concept of the functional system postulated by Luria and has normative data for the Mexican population with 450 subjects between 6 and 80 years old with different schooling ranges. (Flores et al., 2014).

This analysis allows to obtain an overall performance index and a performance index of the three

pre-frontal areas evaluated, the standardized scores have an average of 100 and a standard deviation of 15. The interpretation of the total score and of each of the areas allows to classify the execution of a person as follows: high normal (116 onwards), normal (85-115) slight to moderate alterations (74-84) and severe alterations (less than 69). It is possible to obtain the standard scores for each of the sub-tests through the performance profile, the standard scores for the sub-tests have an average of 10 and a standard deviation of 3. (Flores et al., 2014). Therefore, subscales were chosen that evaluate inhibitory control, rule following and risk-benefit processing, cognitive processes and emotional regulation, which are related to deregulated behaviors related to fat percentage.

### **RENPHO ES-24M Smart Body Composition Scale.**

The RENPHO ES-24M Smart Body Composition scale was used for the measurement of body composition data; it sends a safe low-voltage electrical signal that runs through the body. It allows us to obtain data such as weight, BMI, BFP, and visceral fat level.

### **BFP Classification and values.**

Women: Low <21%, Normal 21 - 32.9%, High 33 - 38.9% and Very high > 39% Men: Low <8%, Normal 8.1 - 19.9%, High 20 - 24.9% and Very high > 25% (Cardozo, et al., 2016).

### **Ethical considerations**

This study was carried out following the criteria of the protocol and in accordance with the Ethical Principles for Medical Research Involving Human Subjects, this research had a minimal risk for participants. (AMM, 2013). This research was approved by the CICS-UST Ethics Committee (CEI-CICS-009).

### **Analysis and mathematical method.**

We used measures of central tendency, dispersion, percentages, and MLP ANN (Turban et al., 2007) with three feedforward connected layers to identify the hidden layer connected to the output layer. The output variables were the predicting variables among the variables of the stimuli or input layer, which are the

BANFE-2 and the OQ scales. For validation, we used 10-fold cross-validation. The learning algorithm used was backpropagation (García & Espinosa, 2013).

### Procedure.

A call for voluntary participation was made in the Centro Interdisciplinario de Ciencias de la Salud, Unidad Santo Tomás, Instituto Politécnico Nacional. We asked the participants to sign the informed consent and then we measured their BFP using the RENPHO ES-24M Smart Body Composition Scale. Once the data was collected, participants were divided into four groups according to their BFP: Low, Normal, High and Very High. None of the participants was classified in the Low Group.

The instruments were applied collectively at the Centro Interdisciplinario de Ciencias de la Salud, Unidad Santo Tomás, Instituto Politécnico Nacional by the authors.

We developed a MLP ANN whose output variables were BFP and metabolic age; and whose input variables were BANFE-2, OQ and gender. SPSS V 22.0 software was used for this analysis.

## Results

The sample was composed of 40 adults of whom 55% were women ( $n = 22$ ). The average age was 24.5 years ( $SD = 3.87$ ). Regarding anthropometric measurements, average BFP was 26.86 ( $SD = 7.23$ ), and average visceral fat was 8.49 ( $SD = 3.92$ ) (Table I).

**Table I. Means by groups of body fat percentage**

Table I			
Sex	Group	n	$\bar{x}$ (SD)
Woman	Normal	16	23.56 ± 4.55
	High	8	34.80 ± 1.65
	Very high	3	37.66 ± 4.61
	Total	27	28.46 ± 7.15
Man	Normal	6	14.93 ± 1.02
	High	3	24.80 ± 5.99
	Very high	13	29.54 ± 2.54

Note:  $\bar{x}$  = average, SD = Standar Deviation.

Table II describes the results of the means and SD. As described in the table, differences in fat percentages are observed, higher in women compared to men.

**Table II**

OQ Scales	$\bar{x}$ (SD)
Overeating	58.27 (10.98)
Defensiveness	51.20 (7.75)
Undereating	54.00 (7.25)
Craving	55.80 (7.12)
Expectations about eating	55.49 (8.96)
Rationalizations	54.47 (8.9)
Social isolation	52.92 (11.7)
Affectivity	60.57 (8.58)
Motivation	60.35 (9.59)
Healthy habits	49.45 (10.83)
Body image	53.96 (9.78)

Note:  $\bar{x}$  = average, SD = Standar Deviation.

Table III shows the applied subtests, which were: Stroop effect, card game and mazes, these tests evaluate the orbitofrontal cortex functions which estimates the ability to detect and avoid risk selections, to exercise inhibitory control, to respect limits and follow rules. For the analysis, the operating index of the orbitofrontal area was obtained based on the standardized scores allowing to group the participants according to the level of alteration of the evaluated cognitive functions.

**Table III**

BANFE-2 Scales	$\bar{x}$ (SD)
Orbitomedial Performance Score	81.27 (21.24)
Labyrinths	7.08 (3.713)
Percentage of Risk Cards	9.84 (2.6)
Total Card Game	9.10 (3.38)
Stroop A Errors	7.57 (4.38)
Stroop A Time	9.69 (2.87)
Stroop A Successes	6.33 (4.14)
Stroop B Errors	8.61 (4.23)
Stroop B Time	10.22 (2.17)
Stroop B Successes	8.86 (4.05)
Maintenance errors, Classification of cards	5.22 (3.16)


Note:  $\bar{x}$  = average, SD = Standar Deviation.

Graph 1 and Graph 2 show the results of average synaptic weights of input factors (OQ and BANFE-2 scales) for BFP output factor (synaptic weight percentage for each input factor).

The model shows a reduction in the error (see Graph 3) as the tests were run, with a low final error of 0.149 and an average error in the first test of 0.30 for the OQ variables and neuropsychological variables.


On the other hand, the model for BANFE-2 also shows a reduction in error (see lower graph) as the tests were run, with a low final error of 0.18 and an average error in the first test of 0.46. The percentages of synaptic weight of the input factors of OQ scales Healthy Habits, Rationalizations, Motivation and Body Image showed a greater synaptic weight in the ANN, with two input layers for the output factor BFP.

**Graph 1.**  
 Average synaptic weight of input factors (OQ).




Note: The averages of the neural weights of all the tests run for the OQ are presented. Health habits and Rationalizations for eating, have a high synaptic weight separately; later, Motivation, Body image and Cravings, with average weight.

**Graph 2.**  
 Average synaptic weights of input factors (BANFE-2).




Note: The averages of the neural weights of all the tests run for the BANFE-2 are presented. In this graph we observe a greater distribution in the importance of synapses separately, which goes from Hit Stroop A to Total card game.

**Graph 3.**  
Error per test of input factors (OQ).



Note: the average error of the ten tests is shown.

**Graph 4.**  
Error per test of input factors (BANFE).



Note: the average error of the ten tests is shown.

## Discussions

The main objective of this research was to identify the synaptic weights of the ANN, whose input variables were the executive functions (EF) and health lifestyles as predictors of Body Fat Percentage (BFP), in a group of adult subjects with low, normal, high, and very high body fat levels.

The implementation of ANN allows predictive models for the estimation of contextual intake variables and neuropsychological variables that contribute interactively to BFP, and that can be generalized for men and women aged between 18-40 years old.

Error correction by least squares allows the learning of patterns of neural networks, with these patterns, categories are generated (García & Espinosa, 2013).

These categories were confirmed with the application of two instruments, one for neuropsychological variables and another for healthy habits of overeating and undereating. We found that Healthy Habits and Rationalizations to Lose Weight, as well as Stroop A Successes, Stroop A Performance, Orbitomedial Performance, Stroop B Time, Stroop B Successes and Stroop B Errors have high synaptic weights for the output variable BFP. Therefore, these models can provide useful tools for the study of complex related variables in the context of OB, in addition to the development of more specific interventions that integrate healthy habits, as confirmed by Adavi, Salehi & Roudbari (2016), and intentional neurological interventions that integrate specially the orbitomedial area. As we can see, the average error decreases in test 10, which allows us to identify in both measurements an adequate adjustment of the ANN (synapse), both for the variables of healthy habits, and for the neuropsychological variables (variables of input) in relation to the percentage of fat (output variable), confirming what was stated by Huang (2019).

As Yang (2018) and Tsai (2019) suggest, Ob and therefore body fat, would generate lower neurocognitive performance mainly in inhibitory control tasks, however our findings transcend this hypothesis, since we found a synaptic weight of the subtest for the orbitofrontal area, related to both executive functions and emotional regulation.

In the multilayer neural network with architecture 9, 3, 3, 1, the specific finding of the synaptic weight of the OQ subscale Rationalizations to Continue Eating to BFP is related to the type of dysfunctional cognitive response of obese individuals, which allows them to disengage from the responsibility of maintaining a healthy body weight.

In the multilayer neural network with architecture 12, 3, 3, 1 with the BANFE-2 input variables and output variable BFP, we found that inhibitory control performance has the highest synaptic weight as the predictor of adiposity. Stroop is related to the processing speed and the ability to control the interference of other brain structures not congruent with a cognitive task, i.e., mechanisms of motor and behavioral inhibition (Golden, 2020; Lubrini et al., 2009 as cited in Blazquez et al. 2009). Therefore, obtaining meaningful

data in Stroop A and B Errors and Successes relates to the effectiveness of these mechanisms to control and coordinate simple cognitive processes.

This study shows the relevance of using ANN for simultaneous analysis of neuropsychological and healthy lifestyle data for the investigation of OB prevention and treatment, especially when considering EF performance regarding intake for the evaluation and modulation of hunger, satiation and satiety, craving and emotional changes in its corresponding behavioral response (García-Flores, 2017; Benelam, 2009), as environmental factors have been shown to impact biological processes and body weight.

As Ergün (2009) and Sarmiento-Ramos (2020) reports, the use of ANN allows health professionals to make more precise and effective decisions to improve well-being and quality of life.

## Conclusions

Maintenance of Healthy Habits subscale measures the regularity of behaviors considered as good health practices; therefore, such synaptic weight for the BFP would be related to an inability to modify healthy behaviors. The use of nonlinear mathematical models allows to identify open links between the dependent variables, in this case the percentage of fat, with differential weights (synaptic networks), and the independent variables (input variables), in this research neuropsychological factors, habits and cognitions related to overeating and undereating as measured by the OQ. Neural Networks are a novel method that can replace linear regressions or accompany these results, both to compare or even strengthen these findings. The orbitofrontal cortex and the prefrontal area are responsible for processing information and inhibiting responses, as well as habits, reasoning style, image perception and cravings which are factors that are related to the percentage of fat. These findings may redirect behavioral therapeutic goals, seeking greater specificity and effectiveness.

For subsequent research, we suggest expanding the sample in clusters by fat percentage, age groups and sociodemographic factors, and compare with linear regressions or binary logistic regressions.



## References

- Adavi, M., Salehi, M. & Roudbari, M. (2016). Artificial neural networks versus bivariate logistic regression in prediction diagnosis of patients with hypertension and diabetes. *Medical journal of the Islamic Republic of Iran*, 30, 312.
- Allan, J., McMinn, D. & Daly, M. (2016). A bidirectional relationship between executive function and health behavior: evidence, implications, and future directions. *Frontiers in neuroscience*, 10, 386.
- Arias, V., Salazar, J., Garicano, C., Contreras, J., Chacón, G., Chacín-González, M. & Bermúdez-Pirela, V. (2019). Una introducción a las aplicaciones de la inteligencia artificial en Medicina: Aspectos históricos. *Revista Latinoamericana de Hipertensión*, 14(5), 590-600.
- Asociación Médica Mundial (2013). Declaración de Helsinki de la AMM: Principios éticos para las investigaciones médicas en seres humanos.
- Bajo, S. & Ballesteros, M. (2002). Redes neuronales: concepto, aplicaciones y utilidad en medicina. *Atención primaria*, 30(2), 119.
- Benelam, B. (2009). Satiating, satiety and their effects on eating behavior. *Nutrition bulletin*, 34(2), 126-173.
- Brandan, N., Aguirre, M., Agolti, G. & Vila, M. (2014). Interrelaciones metabólicas. Interrelaciones metabólicas entre tejidos especializados. Ciclo Ayuno-Alimentación. Interrelaciones metabólicas en estados fisiológicos y patológicos. *Universidad Nacional del Nordeste Facultad de Medicina Cátedra de Bioquímica*.
- Calvo, D., Galioto, R., Gunstad, J. & Spitznagel, M. B. (2014). Uncontrolled eating is associated with reduced executive functioning. *Clinical obesity*, 4(3), 172-179.
- Cardozo, L., Cuervo, & Murcia, J. (2016). Porcentaje de grasa corporal y prevalencia de sobrepeso-obesidad en estudiantes universitarios de rendimiento deportivo de Bogotá, Colombia. *Nutrición clínica y dietética hospitalaria*, 36(3), 68-75.
- Carlson, N. (2014). *Fisiología de la conducta* (pp. 81-82). Madrid: Pearson.
- Carbine, K. A., Christensen, E., LeCheminant, J. D., Bailey, B. W., Tucker, L. A. & Larson, M. J. (2017). Testing food-related inhibitory control to high-and low-calorie food stimuli: Electrophysiological responses to high-calorie food stimuli predict calorie and carbohydrate intake. *Psychophysiology*, 54(7), 982-997.
- Cherbuin, N. & Walsh, E. (2019). Sugar in mind: untangling a sweet and sour relationship beyond type 2 diabetes. *Frontiers in neuroendocrinology*, 54, 100769.
- Ergün U. (2009). The classification of obesity disease in logistic regression and neural network methods. *Journal of medical systems*, 33(1), 67-72.
- García, F. & Espinosa, J. (2013). Estimation of body fat percentage using neural networks. *Revista vínculos*, 10(1), 308-318.
- García-Flores, C., Martínez, A., Beltrán C., Zepeda-Salvador, A. & Solano, L. (2017). Satiación vs saciedad: reguladores del consumo alimentario. *Revista médica de Chile*, 145(9), 1172-1178.
- Golden, C. (2020) *STROOP. Test de Colores y Palabras – Edición Revisada* (6a ed.) TEA Ediciones.
- Heydari, S. T., Ayatollahi, S. M., & Zare, N. (2012). Comparison of artificial neural networks with logistic regression for detection of obesity. *Journal of medical systems*, 36(4), 2449-2454.
- Huang, T., Chen, Z., Shen, L., Fan, X. & Wang, K. (2019). Associations of Cognitive Function with BMI, Body Fat Mass and Visceral Fat in Young Adulthood. *Medicina*, 55(6), 221.
- Lubrini, G., Periañez, J. A. & Rios-Lago, M. (2009). Introducción a la estimulación cognitiva y la rehabilitación neuropsicológica. *Estimulación cognitiva y rehabilitación neuropsicológica*, 13-16.
- Martínez-Mendoza, G. (2019). Funciones ejecutivas y consumo de alcohol en jóvenes universitarios: capacidad predictiva de las medidas de evaluación. *Revista de Psicología Clínica con Niños y Adolescentes*, 6(2), 22-29.
- Mármol, M., & Spano, R. (2018). Diferencias en el desempeño de las funciones ejecutivas en grupos de niños con y sin desnutrición de siete a diez años de La Vega, Antimano y Carapita. *Universidad Católica Andrés Bello, Facultad de Humanidades y Educación Escuela de Psicología*. <http://biblioteca2.ucab.edu.ve/anexos/biblioteca/marc/texto/AAT7115.pdf>
- Narimani, M., Esmailzadeh, S., Azevedo, L. B., Moradi, A., Heidari, B. & Kashfi-Moghadam, M. (2019). Association between weight status and executive function in young adults. *Medicina*, 55(7), 363.
- Psihas, E. (2014). Validación del cuestionario de sobreingesta alimentaria en población mexicana. [online] Repositorio. iberopuebla.mx. Available at: <<https://repositorio.iberopuebla.mx/bitstream/handle/20.500.11777/207/PSIHAS.pdf?sequence=1>> [Accessed 24 January 2022].
- Sarmiento-Ramos, J. (2020). Aplicaciones de las redes neuronales y el deep learning a la ingeniería biomédica. *Revista UIS Ingenierías*, 19(4), 1-18.
- Tirapu, J., García, A., Luna, P., Verdejo, A. & Ríos, M. (2012). Corteza prefrontal, funciones ejecutivas y regulación de la conducta. *Neuropsicología de la corteza prefrontal y las funciones ejecutivas*, 87-117.

- Turban, E., Aronsons, J. E., & Ting-Peng, L. (2007) *Decision Support and Business Intelligence Systems*, 8th Edition. United States Pearson.
- Tsai, C. L., Pan, C. Y., Chen, F. C., Huang, T. H., Tsai, M. C. & Chuang, C. Y. (2019). Differences in neurocognitive performance and metabolic and inflammatory indices in male adults with obesity as a function of regular exercise. *Experimental physiology*, 104(11), 1650-1660.
- World Health Organization. (2005). *WHO STEPS surveillance manual: the WHO STEPwise approach to chronic disease risk factor surveillance* (No. WHO/NMH/CHP/SIP/05.02). World Health Organization.
- Yang, Y., Shields, G., Guo, C. & Liu, Y. (2018). Executive function performance in obesity and overweight individuals: A meta-analysis and review. *Neuroscience & Biobehavioral Reviews*, 84, 225-244.