



Comparison of the most popular methods for predicting stone-free rate after percutaneous nephrolithotomy

Ugalde-Resano R, Villeda-Sandoval CI, Kobashi-Sandoval E, Rivera-Ramírez JA, Vargas-Robles MA, Méndez-Probst CE

Abstract

BACKGROUND: The Guy's stone score, S.T.O.N.E. nephrolithometry, CROES nomogram, and S-ReSC scoring system have recently been externally validated as effective predictors of stone-free rate after percutaneous nephrolithotomy.

OBJECTIVE: We describe herein the advantages of identifying the most accurate scale and propose its standardized use.

MATERIALS AND METHODS: We analyzed 188 patients that underwent percutaneous nephrolithotomy for kidney stones within the time frame of October 2010 and July 2015 at a tertiary care referral center. Preoperative and postoperative non-contrast computed tomography scans were used in all patients to compare the four scoring systems, using the strict criterion of absolute absence of residual stone as the stone-free rate.

RESULTS: The overall stone-free rate was 57.9%. All scoring systems presented a statistically significant stone-free rate ($p < 0.001$). Only the Guy's stone score had no correlation with postoperative complications. The four scoring systems had similar accuracy, none of them were more predictive for stone-free rate than the other, and there was no significant difference in the areas under the curve between them ($p = 0.2$). In addition, each scale had a correlation with operative time and length of hospital stay.

CONCLUSIONS: The four scales analyzed were excellent predictors for stone-free rate. They had similar receiver operating characteristic curves and areas under the curve, with no significant differences between them. However, the Guy's stone score presented the best predictive capacity and the S-ReSC scoring system was the best method for predicting complications. It is up to the urologic community to decide which evidence-based scale is the most suitable. Meanwhile, the need for a standardized method continues to grow.

KEYWORDS Stone-free rate; Percutaneous nephrolithotomy; CROES; S-ReSC, S.T.O.N.E.; Guy's stone score

Urology Department, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico city.

Received: marzo 2017

Accepted: julio 2017

Correspondence

Rodrigo Ugalde Resano
ugalde.resano@gmail.com

This article should be cited as

Ugalde-Resano R, Villeda-Sandoval CI, Kobashi-Sandoval E, y col. Comparison between the most popular methods for predicting the stone free rate after percutaneous nephrolithotomy. Rev Mex Urol. 2017 jul-agos;77(4):279-288.

DOI: <https://doi.org/10.24245/revmexurol.v77i4.1342>

Rev Mex Urol. 2017 jul;77(4):279-288.

Comparación de los métodos más populares para predecir la tasa libre de litiasis posterior a nefrolitotomía percutánea

Ugalde-Resano R, Villeda-Sandoval CI, Kobashi-Sandoval E, Rivera-Ramírez JA, Vargas-Robles MA, Méndez-Probst CE

Resumen

ANTECEDENTES: las escala de Guy y S.T.O.N.E., Nefrolitometría, CROES y el sistema de puntuación S-ReSC se validaron recientemente como predictores efectivos de la tasa libre de litiasis posterior a la nefrolitotomía percutánea.

OBJETIVO: describir las ventajas e identificar la escala más precisa y proponer su utilización.

MATERIALES Y MÉTODOS: estudio retrospectivo de pacientes a quienes se realizó nefrolitotomía percutánea para cálculos renales entre los meses de octubre de 2010 a julio de 2015 en un centro de referencia de atención terciaria. En todos los pacientes se indicaron estudios pre y posoperatorios de tomografía computada sin contraste para comparar las cuatro escalas, se utilizó el criterio estricto de ausencia absoluta de litiasis residual, que se consideró tasa libre.

RESULTADOS: se estudiaron 188 pacientes en quienes la tasa libre de litiasis fue de 57.9%. Todos los sistemas de puntuación reportaron una tasa libre de cálculos estadísticamente significativa ($p < 0.001$), excepto las escala de Guy y S.T.O.N.E no tuvieron correlación con las complicaciones posoperatorias. Los cuatro sistemas de puntuación tuvieron una precisión similar, ninguno fue más predictivo que otro para la tasa libre de litiasis y entre ellas no hubo diferencias significativas en las áreas bajo la curva ($p = 0.2$). Además, cada escala tuvo una correlación con el tiempo operativo y la duración de la estancia hospitalaria.

CONCLUSIONES: las cuatro escalas analizadas son excelentes predictores de la tasa libre de litiasis pues tuvieron receptores, curvas características de funcionamiento y áreas bajo la curva similares, sin diferencias significativas entre ellas. Sin embargo, las escalas de Guy y S.T.O.N.E tuvo la mejor capacidad predictiva y el puntaje S-ReSC fue el mejor método para predecir complicaciones. Depende de la comunidad urológica decidir cuál escala basada en la evidencia es la más adecuada. Mientras tanto, la necesidad de un método estandarizado sigue creciendo.

PALABRAS CLAVE: tasa libre de litiasis; nefrolitotomía percutánea; CROES; S-ReSC, S.T.O.N.E.; escala de Guy.

Departamento de Urología, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Ciudad de México, México.

Recibido: marzo 2017

Aceptado: julio 2017

Correspondencia

Dr. Roberto Patricio López Maguey
rpatloma@gmail.com

Este artículo debe citarse como

Ugalde-Resano R, Villeda-Sandoval CI, Kobashi-Sandoval E, y col. Comparación de los métodos más populares para predecir la tasa libre de litiasis posterior a nefrolitotomía percutánea. Rev Mex Urol. 2017 jul-agos;77(4):279-288.



INTRODUCTION

Percutaneous nephrolithotomy is now the first-line treatment option for large, complex stones and staghorn calculi, greatly reducing the need for open surgery.¹⁻²

Several studies have identified significant predictors of stone-free rate after percutaneous nephrolithotomy, with stone size, number, location, and pyelocaliceal system anatomy as the suggested predictors.³⁻⁴ Nevertheless, a significant predictor alone, is not a predictive tool. Some authors developed different scoring systems to standardize the terminology in relation to stone complexity. The Guy's stone score,⁵ S.T.O.N.E. nephrolithometry,⁶ CROES nomogram,⁷ and S-ReSC scoring system⁸ have recently been externally validated and they all effectively predicted stone-free rate after percutaneous nephrolithotomy.⁹⁻¹⁶ However, none of them has gained wide acceptance or implementation into clinical practice.¹⁷

There are many benefits to having a standardized method of predicting the stone-free rate after percutaneous nephrolithotomy. The primary aim of our study was to compare the most popular scoring systems, describe their advantages to identify the most accurate scale, and propose its standardized use. Our study is the first comparison of the four scoring systems in the same patient group.

MATERIALS AND METHODS

We analyzed a total of 188 patients that underwent percutaneous nephrolithotomy for kidney stones, within the time frame of October 2010 and July 2015 at a tertiary care referral center. Patients with incomplete data (n=36) were excluded from the study. All procedures were performed by a single experienced endourologist. The patients were placed in the prone

position and received general anesthesia, and the surgical technique was carried out according to previously published manuscripts.¹⁸⁻¹⁹

A preoperative non-contrast computed tomography scan was utilized in all patients to evaluate stone characteristics (stone burden, laterality, location, number, and density). Stone burden was estimated using the following formula: length x width x pi x 0.25, where pi is a mathematical constant equal to 3.14.²⁰ A junior urology resident from our institution reviewed all images and calculated the corresponding Guy's,⁵ S.T.O.N.E.,⁶ CROES,⁷ and S-ReSC⁸ scores. We compared and correlated the scores with preoperative and postoperative data. Each scoring system was categorized according to its original description, but the CROES nomogram score and stone burden were randomly categorized for better statistical analysis.

The demographic data and length of hospitalization were available from our prospective percutaneous nephrolithotomy database. Postoperative complication data were graded using the modified Clavien classification system,²¹ collected from the contemporaneous electronic patient records, radiologic imaging findings, and paper case notes. Patients had a non-contrast computed tomography scan between the first and third month follow-up visit. Postoperative stone-free rate was defined using the strict criterion of absolute absence of residual stone.

Statistical analyses were performed with the Statistical Package of Social Sciences version 20 (SPSS, Chicago, IL, USA). Categorical variables were presented as numbers and percentages and compared with the chi-square test. The Kruskal-Wallis test was used for the statistical analysis of the ordinal variables to assess the categories of the scoring systems. Continuous variables were presented as means and standard deviations and compared with an independent sample t test.

Correlation analyses were evaluated using the Pearson correlation coefficient (*r*). The area under the curve (AUC), calculated from the receiver operating characteristic (ROC) curve, was used to assess the predictive ability of the different scoring systems. The AUCs were compared using the online calculator of significance of difference between areas under two independent ROC curves from the website http://vassarstats.net/roc_comp.html accessed on November 30th, 2015. Statistical significance was considered at a two-tailed *p* value <0.05.

RESULTS

We identified 152 patients that underwent percutaneous nephrolithotomy between 2010 and 2015 at a single tertiary care referral center and that met the study inclusion criteria. **Table 1**

shows the demographic and preoperative characteristics of the patients. The overall stone-free rate was 57.9%, with the strict criterion of absolute absence of residual stone. The postoperative complication rate was 39.5%, with Clavien grade I in 37 patients, Clavien grade II in 14 patients, Clavien grade III in 3 patients, and Clavien grade IV in 6 patients. There were no deaths.

In patients that were stone-free and in those with residual stones, the mean Guy's stone score was 1.99 and 3.23, the mean S.T.O.N.E. score was 6.94 and 9.42, the mean CROES score was 177.34 and 127.28, and the mean S-ReSC score was 2.88 and 5.03, respectively (*p*<0.001 each). In patients that presented with any complications and in those with no complications, the mean Guy's stone score was 2.6 and 2.4 (*p*=0.38), the mean S.T.O.N.E. score was 8.3 and 7.6

Table 1. Demographic and preoperative characteristics

	Stone-free	Not stone-free	<i>p</i>
Number of patients (%)	88 (57)	64 (42.1)	
Mean age ± SD (years)	49.48 ± 14.1	47.65 ± 13	0.418 ^a
Sex (%)			
Male	34 (38.6)	20 (31.2)	0.34 ^b
Female	54 (61.4)	44 (68.8)	
Laterality (%)			
Right	43 (48.9)	25 (39.1)	0.23 ^b
Left	45 (51.1)	39 (60.9)	
Mean BMI ± SD (kg/m ²)	28.54 ± 5.6	27.21 ± 4.8	0.161 ^a
Mean Hounsfield Units ± SD	840.3 ± 335.3	893.7 ± 275.7	0.298 ^a
Mean number of stones ± SD	2.1 ± 1.6	3.3 ± 2.3	0.001 ^{*a}
Mean operative time ± SD (min)	167 ± 67.34	245.1 ± 100.08	<0.001 ^{*a}
Mean length of hospital stay ± SD (days)	3.4 ± 3.6	4.4 ± 3.4	0.081 ^a
Number of staghorn stones (%)	16 (18.2)	20 (54.7)	<0.0001 ^{*a}
Multiple locations (%)	36 (81.8)	8 (18.2)	<0.0001 ^{*a}
Mean stone burden ± SD (mm ²)	411.5 ± 362.7	895.6 ± 693.2	<0.0001 ^{*a}
Mean Guy's stone score ± SD	1.99 ± 1.07	3.23 ± 0.95	<0.001 ^{*a}
Mean S.T.O.N.E. score ± SD	6.94 ± 1.53	9.42 ± 1.79	<0.001 ^{*a}
Mean CROES score ± SD	177.34 ± 50.46	127.28 ± 59.76	<0.001 ^{*a}
Mean S-ReSC score ± SD	2.88 ± 1.69	5.03 ± 2.42	<0.001 ^{*a}

* Statistical significance <0.05; ^a Compared with an independent sample *t* test; ^b Compared with the chi-square test.



($p=0.069$), the mean CROES score was 146.2 and 164.8 ($p=0.061$), and the mean S-ReSC score was 4.2 and 3.33 ($p=0.016$), respectively.

Table 2 shows the stone-free rate of each of the four scoring systems. The S.T.O.N.E., CROES, and S-ReSC groups were significantly associated with the stone-free rate and complication rate. The Guy's stone score was associated with the stone-free rate, but not the complication rate. Each scale had a correlation with operative time and length of hospital stay: Guy's stone score ($r=0.41$, $p<0.001$) ($r=0.22$, $p=0.007$), S.T.O.N.E. score ($r=0.50$, $p<0.001$) ($r=0.33$, $p<0.001$), CROES score ($r=0.40$, $p<0.001$) ($r=0.27$, $p<0.001$), and S-ReSC score ($r=0.35$, $p<0.001$) ($r=0.20$, $p=0.012$), respectively. Stone burden also correlated with operative time and hospital stay duration ($r=0.41$, $p<0.0001$) ($r=0.41$, $p=0.022$).

Table 3 and **Figure 1** show the AUC and ROC curves for each scoring system and for stone bur-

den in relation to the stone-free rate. All scoring systems had similar accuracy and none was more predictive for stone-free rate than another. There was no significant difference in the AUC between the four scoring systems ($p=0.2$). However, the Guy's stone score had the greatest AUC for predicting the stone-free rate. **Table 4** and **Figure 2** show the AUC and ROC curves in relation to the complication rates. All the scoring systems had poor predictive capacity for complications and only the S-ReSC score had a statistically significant AUC ($p=0.007$).

DISCUSSION

Multiple attempts to identify significant predictors of stone-free rate after percutaneous nephrolithotomy have been made, since the procedure became the first-line surgical treatment for kidney stones.¹⁻² Tefekli et al. divided stones into simple or complex calculi, according to their location in the renal pelvis and calices.²²

Table 2. Association with stone-free rate and complication rate

Scoring system	No. stone free/ Total No. (%)	p^c	No. complication/ Total No. (%)	p^c
CROES				
0-100	3/29 (10.3)		18/29 (62)	
101-200	53/82 (64.6)	<0.0001*	27/82 (32.9)	0.020*
201 or greater	32/41 (78)		15/41 (36.5)	
S.T.O.N.E. (category)				
4-5	14/15 (93.3)		1/15 (6.6)	
6-8	56/74 (75.7)	<0.0001*	30/74 (40.5)	0.020*
9-13	18/63 (28.6)		29/63 (46)	
Guy's (grade)				
I	36/38 (94.7)		14/38 (36.8)	
II	32/49 (65.3)	<0.0001*	16/49 (32.6)	0.499
III	5/14 (35.7)		7/14 (50)	
IV	15/51 (29.4)		23/51 (45)	
S-ReSC (category)				
1-2	47/57 (82.4)		17/57 (29.8)	
3-4	27/47 (57.4)	<0.0001*	15/47 (31.9)	0.005*
5-9	14/48 (29.1)		28/48 (58.3)	
Stone burden (mm²)				
1-500	64/87 (73.5)		32/87 (36.7)	
501-1000	21/38 (55.2)	<0.0001*	10/38 (26.3)	0.004*
>1000	3/27 (11.1)		18/27 (66.6)	

* Statistical significance <0.05; ^c Compared with the Kruskal-Wallis test.

Table 3. ROC curve values for the stone-free rate

Scoring system	AUC	95% CI	p
Guy's	0.791	(0.71)-(0.86)	<0.0001*
S.T.O.N.E.	0.767	(0.69)-(0.84)	<0.0001*
CROES	0.722	(0.64)-(0.80)	<0.0001*
S-ReSC	0.746	(0.66)-(0.85)	<0.0001*
Stone burden (categorized)	0.724	(0.63)-(.80)	<0.0001*




Figure 1. Stone-free rate.

Table 4. ROC curve values for the complication rate

Scoring system	AUC	95% CI	p
Guy's	0.550	(0.45)-(0.64)	0.300
S.T.O.N.E.	0.591	(0.50)-(0.68)	0.058
CROES	0.579	(0.48)-(0.67)	0.100
S-ReSC	0.630	(0.53)-(0.72)	0.007*
Stone burden (categorized)	0.570	(0.47)-(.66)	0.147




Figure 2. Complication rate.

Nevertheless, it is not enough to standardize the terminology of the complexity of the procedure. Several scoring systems for predicting the stone-free rate have recently been published, but none of them has widespread acceptance or implementation in reported clinical practice.

The potential benefit of a standardized method for predicting the stone-free rate after percutaneous nephrolithotomy is reported across different case series.²³⁻²⁴ The clear advantages of a widely-accepted scoring system include more accurate preoperative patient counseling, surgical planning, and outcome evaluation, as well as uniform academic reporting. It could also aid in resource management, in referring complex cases to specialized centers,²⁵ or even in making the decisions to use adjunct techniques, such as combining the procedure with ureterorenoscopy. The ideal scoring system must be simple, repro-



ducible, and provide a high degree of accuracy to estimate the success of the procedure. The Guy's stone score, S.T.O.N.E. nephrolithometry, CROES nomogram, and S-ReSC scoring system have recently been externally validated as predictors of stone-free rate after percutaneous nephrolithotomy, using preoperative non-contrast computed tomography.^{9,15-17} However, our study provides the first comparison of the four scoring systems in the same cohort, with the strict criterion of absolute absence of residual stone in the non-contrast computed tomography study. As our results show, those four scoring systems were significantly associated with the stone-free rate after percutaneous nephrolithotomy ($p < 0.0001$), which has also been demonstrated by original articles describing the same scales.⁵⁻⁸ Those results have recently been replicated in several studies, but there is much discrepancy among them, regarding the predictive accuracy of each scoring system.^{5,15-16} This may be due to the fact that each system was constructed, based on the patient population analyzed, resulting in an intrinsic bias favoring predictive efficacy. In our study, the Guy's stone score showed the wider AUC, reaching 0.791, followed by the S.T.O.N.E., S-ReSC, and CROES scores, respectively. Sfoungaristos et al.⁹ described a very similar ROC curve for the Guy's stone score in their study, with a high AUC of 0.796 and excellent statistical significance ($p < 0.001$).

Another relevant factor that could be involved in the inter-study discrepancy is the absence of a standardized definition for stone-free rate after percutaneous nephrolithotomy. Many of the studies applied the criterion suggested by Opondo et al.²⁴ of no stones visible, or the presence of clinically insignificant residual fragments < 4 mm, for treatment success. Nonetheless, residual stone size does not always correlate with clinical significance.²⁶⁻²⁷ Moreover, different imaging methods were employed in some of the studies. Abdominal X-ray was the most commonly

used method for stone-free evaluation, which is inferior to non-contrast computed tomography imaging for residual fragment assessment.²⁸ Sensitivity was reported at close to 70%, and with a cut-off level of < 4 mm, it reached 85.7%, whereas non-contrast computed tomography had almost 100% sensitivity and has been accepted as the gold standard.²⁹ We used the strict criterion of absolute absence of residual stone, because our experience, together with the evidence acquired from several reports, has shown that a significant number of patients with residual fragments will experience a stone-related event during the postoperative period,²⁶⁻³⁰ reaching up to 46%. Of those events, renal colic is the most common, followed by stone regrowth, increasing the need for additional intervention. Gokce et al.²⁹ reported an absolute stone-free rate of 54.9%, using non-contrast computed tomography as the imaging control and applying the same strict criterion that we used. Their results were slightly lower than ours of almost 58%.

Despite all scoring systems having similar predictive accuracy, we found no significant difference between each AUC ($p = 0.2$) and none was more predictive than another for stone-free rate. Noureldin et al. showed the same results, comparing the AUCs of the Guy's stone score and the S.T.O.N.E. score ($p = 0.6$).³¹ It is up to the urologist to consider the specific characteristics of each method to decide which should be used as the standard in clinical practice and academic reporting.

All the scoring systems analyzed in the present work had significant correlation with operation time and length of hospital stay, concurring with the results of most of the published articles.^{8,31-32} However, the correlation between the S-ReSC score and length of hospital stay had not been described until now. Another important quality is the excellent inter-observer agreement of the four scales, which makes them very reliable in-

struments. The CROES nomogram was developed from a large prospective study that included 2,806 patients from 96 centers.⁷ Nevertheless, we feel it is a very complicated scoring system, compared with the others, making its every day application a challenge. Furthermore, its capacity to predict stone-free rate was no more accurate than that of stone burden alone,¹⁷ which has been described in several works as a significant predictor of stone-free rate.³³⁻³⁴ We, too, demonstrated its direct correlation with operative time and length of hospital stay. However, stone burden is commonly expressed as the largest diameter, which is potentially inaccurate, because of the complex shape of large kidney stones.

In a multi-center study of 850 patients, S.T.O.N.E. nephrolithometry was significantly associated with the complication rate,¹² as was also demonstrated by us, but unlike the study of Okhunov et al.,⁶ in which it could not be correlated with post-percutaneous nephrolithotomy complications ($p=0.09$). That scale is relatively easy to use and showed a high degree of accuracy in our study. However, it consists of variables that are obtained specifically from non-contrast computed tomography images and requires specific software to calculate the different variables,¹⁶ representing an important disadvantage for centers that do not have the necessary resources. On the other hand, the Guy's stone score was initially developed using preoperative plain abdominal X-ray, which is the most common assessment method. Moreover, said scale presented the best predictive ability for stone-free rate, when calculated using non-contrast computed tomography. We found no significant association with postoperative complications ($p=0.49$), results that are consistent with other studies.^{6,11,16,21} In contrast, Vicentini et al.¹³ reported a statistically significant positive association. The Guy's stone score was based on results in the literature and expert opinion, which we believe gives it a much more practical value, providing a simple, reproducible, and accurate method. Its main disadvantage

is the poor agreement among reviewers when grading patients with partial *versus* complete staghorn stones.⁵

In addition to all the above, the S-ReSC score is a recent, but not widely used, grading scale that has presented excellent stone-free rate predictive capacity. Our results showed no statistical difference between the different AUCs analyzed. This scoring system was developed under the hypothesis that complex stone distribution was the most powerful predictor of treatment success, counting the number of sites involved.⁸ However, stone distribution is closely related to stone size and number.¹⁵ The S-ReSC scoring system does not require any particular software and is extremely simple, requiring approximately 15 seconds to score, according to the original description.⁸ Additionally, it showed a significant ability to predict complications, something not previously described. Its greatest advantage is its almost perfect inter-observer reliability, with a correlation coefficient of 0.949 (95% CI: 0.922-0.969, $p<0.001$),¹⁵ the highest reported among all the assessment scales. In our opinion, these facts make it the ideal scoring system for standardizing its daily application and homogenizing criteria.

The main limitation of our study was its retrospective design, but the data were evaluated by an experienced endourologist to substantiate the clinical information. In addition, inter-observer reliability could not be analyzed, because the complexity grading was performed by a single junior urology resident from our institution. It is up to the urologic community to eventually decide which evidence-based scale is the most suitable. The need for a standardized method continues to grow, as experience increases and studies continue to be published.

CONCLUSION

The four scoring systems analyzed were significant predictors of stone-free rate and they had



similar ROC curves and AUCs, with no significant differences. However, the Guy's stone score demonstrated the best predictive capacity and the S-ReSC scoring system proved to be superior for predicting complications.

Conflict of Interest

The authors declare that they have no conflict of interest.

REFERENCES

1. Fernstrom I, Johansson B. Percutaneous pyelolithotomy: a new extraction technique. *Scand J Urol Nephrol*. 1976;10:257-259.
2. Preminger GM, Assimos DG, Lingeman JE, Nakada SY, Pearle MS, Wolf JS Jr. Chapter 1: AUA guideline on management of staghorn calculi: diagnosis and treatment recommendations. *J Urol*. 2005;173:1991-2000.
3. Binbay M, Akman T, Ozgor F, Yazici O, Sari E, Erbin A, Kezer C, Sarilar O, Berberoglu Y, Muslumanoglu AY. Does pelviccaliceal system anatomy affect success of percutaneous nephrolithotomy? *Urology* 2011;78:733-737.
4. Mishra S, Sabnis RB, Desai MR. Percutaneous nephrolithotomy monotherapy for staghorn: paradigm shift for 'staghorn morphometry' based clinical classification. *Curr Opin Urol* 2012;22: 148-153.
5. Thomas K, Smith NC, Hegarty N, Glass JM. The Guy's stone score-grading the complexity of percutaneous nephrolithotomy procedures. *Urology* 2011;78(2):277-281.
6. Okhunov Z, Friedlander JJ, George AK, Duty BD, Moreira DM, Srinivasan AK, Hillelsohn J, Smith AD, Okeke Z. S.T.O.N.E. nephrolithometry: novel surgical classification system for kidney calculi. *Urology* 2013;81(6):1154-1159.
7. Smith A, Averch TD, Shahrouf K, Opondo D, Daels FP, Labate G, Turna B, de la Rosette JJ. A nephrolithometric nomogram to predict treatment success of percutaneous nephrolithotomy. *J Urol* 2013; 190: 149.
8. Jeong CW, Jung J-W, Cha WH, Lee BK, Lee S, Jeong SJ, Hong SK, Byun SS, Lee SE. Seoul National University Renal Stone Complexity Score for Predicting Stone-Free Rate after Percutaneous Nephrolithotomy. 2013; *PLoS ONE* 8(6): e65888.
9. Sfoungaristos S, Lorber A, Gofrit EN, Yutkin V, Landau EH, Pode D, Duvdevani M. External Validation and Predictive Accuracy Assessment of Guy's Stone Score as a Preoperative Tool for Estimating Percutaneous Nephrolithotomy Outcomes. *J Endourol*. 2015 Oct;29(10):1131-5.
10. Mandal S, Goel A, Kathpalia R, Sankhwar S, Singh V, Sinha RJ, Singh BP, Dalela D. Prospective evaluation of complications using the modified Clavien grading system, and of success rates of percutaneous nephrolithotomy using Guy's Stone Score: a single-center experience. *Indian J Urol* 2012; 28: 392
11. Ingimarsson JP, Dagrosa LM, Hyams ES, Pais VM Jr. External validation of a preoperative renal stone grading system: reproducibility and inter-rater concordance of the Guy's stone score using preoperative computed tomography and rigorous postoperative stone-free criteria. *Urology* 2014; 83: 45.
12. Okhunov Z, Moreira D, George A. Multicenter validation of S.T.O.N.E. nephrolithometry. *J Urol, suppl.*, 2014; 191: e839, abstract PD32-09.
13. Vicentini FC, Marchini GS, Mazzucchi E, Claro JF, Srougi M. Utility of the Guy's stone score based on computed tomographic scan findings for predicting percutaneous nephrolithotomy outcomes. *Urology* 2014;83(6):1248-1253
14. Okhunov Z, Helmy M, Perez-Lansac A, Menhadji A, Bucur P, Kolla SB, Cho JS, Osann K, Lusch A, Landman J. Interobserver reliability and reproducibility of S.T.O.N.E. nephrolithometry for renal calculi. *J Endourol* 2013; 27(10):1303-1306
15. Choo MS, Jeong CW, Jung JH, Lee SB, Jeong H, Son H, Kim HH, Oh SJ, Cho SY. External Validation and Evaluation of Reliability and Validity of the S-ReSC Scoring System to Predict Stone-Free Status after Percutaneous Nephrolithotomy. 2014; *PLoS ONE* 9(1): e83628.
16. Labadie K, Okhunov Z, Akhavein A, Moreira DM, Moreno-Palacios J, Del Junco M, Okeke Z, Bird V, Smith AD, Landman J. Evaluation and comparison of urolithiasis scoring systems used in percutaneous kidney stone surgery. *J Urol* 2015;193:154-159.
17. Sfoungaristos S, Gofrit ON, Yutkin V, Landau EH, Pode D, Duvdevani M. External validation of CROES nephrolithometry as a preoperative predictive system for percutaneous nephrolithotomy outcomes. *J Urol*. 2016 Feb; 195 (2): 372-6.
18. Shahrouf W, Andonian S. Ambulatory percutaneous nephrolithotomy: initial series. *Urology* 2010;76:1288-1292.
19. Li R, Louie MK, Lee HJ, Osann K, Pick DL, Santos R, McDougall EM, Clayman RV. Prospective randomized trial of three different methods of nephrostomy tract closure after percutaneous nephrolithotripsy. *BJU Int* 2011; 107: 1660.
20. Tiselius HG, Andersson A. Stone burden in an average Swedish population of stone formers requiring active stone removal: how can the stone size be estimated in the clinical routine? *Eur Urol* 2003;43:275.
21. de la Rosette JJ, Opondo D, Daels FP, Giusti G, Serrano A, Kandasami SV, Wolf JS Jr, Grabe M, Gravas S. Categorisation of complications and validation of the Clavien score for percutaneous nephrolithotomy. *Eur Urol* 2012;62(2):246-255
22. Tefekli A, Ali-Karadag M, Tepeler K, Sari E, Berberoglu Y, Baykal M, Sarilar O, Muslumanoglu A. Classification of percutaneous nephrolithotomy complications using the modified Clavien grading system: looking for a standard. *Eur Urol*. 2008;53:184-190

23. Hyams ES, Bruhn A, Lipkin M, Shah O. Heterogeneity in the reporting of disease characteristics and treatment outcomes in studies evaluating treatments for nephrolithiasis. *J Endourol* 2010;24: 1411.
24. Opondo D, Gravas S, Joyce A, Pearle M, Matsuda T, Sun YH, Assimos D, Denstedt J, de la Rosette J. Standardization of patient outcomes reporting in percutaneous nephrolithotomy. *J Endourol* 2014; 28:767.
25. Rassweiler JJ, Renner C, Eisenberger F. The management of complex renal stones. *BJU Int.* 2000;86:919-928.
26. Raman JD, Bagrodia A, Gupta A, Bensalah K, Cadeddu JA, Lotan Y, Pearle MS. Natural history of residual fragments following percutaneous nephrostolithotomy. *J Urol.* 2009;181:1163-8.
27. Ganpule A, Desai M. Fate of residual stones after percutaneous nephrolithotomy: a critical analysis. *J Endourol.* 2009;23:399-403.
28. Heidenreich A, Desgrandschamps F, Terrier F. Modern approach of diagnosis and management of acute flank pain: review of all imaging modalities. *Eur Urol.* 2002;41:351-362.
29. Gokce MI, Ozden E, Suer E, Gulpinar B, Gulpinar O, Tangal S. Comparison of imaging modalities for detection of residual fragments and prediction of stone related events following percutaneous nephrolithotomy. *Int Braz J Urol.* 2015; 41: 86-90.
30. Altunrende F, Tefekli A, Stein RJ, Autorino R, Yuruk E, Laydner H, Binbay M, Muslumanoglu AY. Clinically insignificant residual fragments after percutaneous nephrolithotomy: medium-term follow-up. *J Endourol.* 2011;25(6):941-5.
31. Noureldin YA, Elkoushy MA, Andonian S. Which is better? Guy's versus S.T.O.N.E. nephrolithometry scoring systems in predicting stone-free status post-percutaneous nephrolithotomy. *World J Urol.* 2015;33(11):1821-5.
32. Bozkurt IH, Aydogdu O, Yonguc T, Yarimoglu S, Sen V, Gunlusoy B, Degirmenci T. Comparison of Guy and Clinical Research Office of the Endourological Society Nephrolithometry Scoring Systems for Predicting Stone-Free Status and Complication Rates After Percutaneous Nephrolithotomy: A Single Center Study with 437 Cases *J Endourol.* 2015;29(9):1006-10.
33. Zhu Z, Wang S, Xi Q, Bai J, Yu X, Liu J. Logistic regression model for predicting stone-free rate after minimally invasive percutaneous nephrolithotomy. *Urology.* 2011;78:32-36.
34. Desai M, De Lisa A, Turna B, Rioja J, Walfridsson H, D'Addessi A, Wong C. The clinical research office of the endourological society percutaneous nephrolithotomy global study: staghorn versus nonstaghorn stones. *J Endourol* 2011;25: 1263-1268.

AVISO PARA LOS AUTORES

Revista Mexicana de Urología tiene una nueva plataforma de gestión para envío de artículos: <https://www.revisionporpares.com/index.php/RMUrol> ahí podrá inscribirse a la base de datos administrada por el sistema *Open Journal System* (OJS) que ofrece las siguientes ventajas para los autores:

- Subir sus artículos directamente al sistema.
- Conocer, en cualquier momento, el estado de los artículos enviados, es decir, si ya fueron asignados a un revisor, aceptados con o sin cambios, o rechazados.
- Participar en el proceso editorial corrigiendo y modificando sus artículos hasta su aceptación final.